TYPE-SPECIMENS OF BIRDS

IN THE

BRITISH MUSEUM (NATURAL HISTORY)

Vol. 2
PASSERINES

RACHEL L. M. WARREN AND C. J. O. HARRISON



TRUSTEES OF
THE BRITISH MUSEUM (NATURAL HISTORY)
LONDON: 1971

© Trustees of the British Museum (Natural History) 1971



INTRODUCTION

This second volume lists in alphabetical order the name-bearing type specimens of passerine birds in the Department of Zoology of the British Museum (Natural History), following the format used for the non-passerines in volume I. The key numbers refer to passerine families or sub-families as follows:—

1.	Eurylaimidae	27.	Cinclidae	48.	Meliphagidae
2.	Dendrocolaptidae	28.	Troglodytidae		Emberizidae
3.	Furnariidae	29.	Mimidae	49.	Emberizinae
4.	Formicariidae	30.	Prunellidae	50.	Cardinalinae
5.	Conopophagidae		Muscicapidae		Richmondeninae
6.	Rhinocryptidae	31	Turdinge	51	Theoreign

e)

ιe

TI. EUSICIUDIUAC

70. Farauisaeiuae 71. Corvidae

The use of letters in addition to figures was necessary owing to taxonomic changes during the period of preparation.

Volume 2 differs from volume 1 in including drawings in the Department of Zoology accepted as type specimens (see below). The drawings of non-passerines accepted as types are listed among other amendments to volume 1 on pp. 613-622.

A third volume is in preparation which lists in systematic order all the names included in volumes 1 and 2.

The International Code of Zoological Nomenclature lays responsibilities upon the institutions in which types are deposited (recommendation 72). However, as was pointed out in the Introduction to Volume 1, they cannot

© Trustees of the British Museum (Natural History) 1971

ERRATA

- p. 113, line 26, chyula should be chyulu.
- p. 149, line 45, pallidae should be pallida.
- p. 175, line 20, ferreo rostris should be ferreorostris.
- p. 257, last entry, family number should be 35.
- p. 344, delete line 5.
- p. 345, fourth entry, family number should be 38.
- p. 345, fifth entry, family number should be 37A.
- p. 373, line 13, China Seas should be Malaya.
- p. 385, line 8, nigro rufa should be nigrorufa.
- p. 413, line 28, pallidicpes should be pallidiceps.



INTRODUCTION

This second volume lists in alphabetical order the name-bearing type specimens of passerine birds in the Department of Zoology of the British Museum (Natural History), following the format used for the non-passerines in volume I. The key numbers refer to passerine families or sub-families as follows:—

1.	Eurylaimidae	27.	Cinclidae	48.	Meliphagidae
2.	Dendrocolaptidae	28.	Troglodytidae		Emberizidae
3.	Furnariidae	29.	Mimidae	49.	Emberizinae
4.	Formicariidae	30.	Prunellidae	50.	Cardinalinae
5.	Conopophagidae		Muscicapidae		Richmondeninae
6.	Rhinocryptidae	31.	Turdinae	51.	Thraupinae
7.	Cotingidae	32A.	Orthonychinae	52.	Tersininae
8.	Pipridae	32B.	Timaliinae	53.	Geospizinae
9.	Tyrannidae	33.	Paradoxornithinae	54.	Parulidae
10.	Oxyruncidae	34.	Polioptilinae	55.	Drepanididae
11.	Phytotomidae	35.	Sylviinae	56.	Vireonidae
12.	Pittidae	36.	Malurinae	57.	Icteridae
13.	Acanthisittidae	37.	Muscicapinae		Fringillidae
14.	Philepittidae	37A.	Rhipidurinae	58.	Fringillinae
15.	Menuridae	38.	Monarchinae	59.	Carduelinae
16.	Atrichornithidae	39.	Pachycephalinae	60.	Estrildidae
17.	Alaudidae	40.	Aegithalidae	61.	Ploceidae
18.	Hirundinidae	41.	Remizidae	62.	Sturnidae
19.	Motacillidae	42.	Paridae	63.	Oriolidae
20.	Campephagidae	43.	Sittidae	64.	Dicruridae
21.	Pycnonotidae	44A.	. Certhiidae	65.	Callaeidae (none)
22.	Irenidae	44B.	Rhabdornithidae	66.	Grallinidae
23.	Laniidae	44C.	Climacteridae	67.	Artamidae
24.	Vangidae	45.	Dicaeidae	68.	Cracticidae
25.	Bombycillidae	46.	Nectariniidae	69.	Ptilinorhynchidae
26.	Dulidae	47.	Zosteropidae	70.	Paradisaeidae
				71.	Corvidae

The use of letters in addition to figures was necessary owing to taxonomic changes during the period of preparation.

Volume 2 differs from volume 1 in including drawings in the Department of Zoology accepted as type specimens (see below). The drawings of non-passerines accepted as types are listed among other amendments to volume 1 on pp. 613-622.

A third volume is in preparation which lists in systematic order all the names included in volumes 1 and 2.

The International Code of Zoological Nomenclature lays responsibilities upon the institutions in which types are deposited (recommendation 72). However, as was pointed out in the Introduction to Volume 1, they cannot

fulfil these responsibilities unless authors notify them of the publication of new names based on specimens in their institutions. Labelling of the specimens before publication is no substitute; see the entry "meinertzhageni" in the additions and corrections to volume 1 on p. 618.

We hope that our colleagues in other museums will find this catalogue to be a valuable, accurate and systematic record of the bird holotypes and selected syntypes, numbering about 8,500, in the Department of Zoology.

We apologise for any minor inconsistencies of presentation, due to the fact that the original entries were prepared in systematic order, over a period of years, and that more than one person was involved. Some small differences in the wording of entries which only became apparent in the final proofs were not considered of sufficient importance to merit alteration.

Drawings as Type Specimens

The library of the Museum holds a number of original drawings of birds, on which, in some instances, the descriptions and names of new species were based, and the models of which were not preserved. These drawings must, therefore, be considered as the holotypes, and all those accepted as such are now listed in this volume. They are all of Australian birds, by four artists of whom Forster, Parkinson and Ellis have been dealt with by Lysaght (1959). The fourth was *Thomas Watling* [1762–?c. 1810].

Watling arrived at Port Jackson, Sydney, in October 1792, as a convict transported for forgery; and was employed by Surgeon J. White to make drawings 'of landscapes and natural objects'. Latham (1801, and in Sharpe 1906) may have based his descriptions in whole or in part on Watling's drawings or on copies of them by himself and others (see references on p. vi) although he also had access to the paintings by various artists collected by A. B. Lambert. These were drawn in New South Wales and sent to Latham in about 1800, and are now in Lord Derby's library at Knowsley, Lancashire. The two series of drawings were compared by H. W. England, Librarian of the department of Zoology at the Museum (Mathews 1942), and he numbered the drawings in the sequence in which they are mounted in the volumes. Both the Watling and the Lambert drawing numbers are given in the entries here. number of the drawings listed by Sharpe (1906) cannot be accepted as types of Latham's descriptions; some being now considered indeterminate, while in other instances Latham used other material. descriptions were wrongly attributed to Latham by Sharpe. Four of Latham's English names, although listed by Sharpe, were not latinized by him and are not types. Since they are in English they are not referred to in the main sequence of this volume; they are :- no. 270 Crested or Caspian Tern, no. 274 New Holland Tern, no. 277 New Holland Crimsonbilled Gull, and no. 291 New Holland Penguin.

Additional notes on Authors and Collections

Johann Natterer [1787–1843] and August von Pelzeln [1825–1891]

The collection made by Natterer in Brazil between 1817 and 1835 went to the Imperial Zoological Cabinet, Vienna, and was described by von Pelzeln (1871) using some names proposed by Natterer. Von Pelzeln would refer to several specimens that had been collected from various localities on different dates, and only rarely selected a holotype. P. L. Sclater received a number of the specimens in exchange with Vienna, and his collection was purchased by the Museum between 1884 and 1890. Those specimens referred to by Sclater (1888–1890) as "authentic specimens", the dates and localities of which agree with the original description, are recognised here as syntypes.

N. A. Vigors [1785–1840] and T. Horsfield [1773–1859]

"Australian birds in the collection of the Linnean Society" (Trans. Linn. Soc. Lond., 15: 170-331) was published on 9 February 1827 (minutes of the Council of the Society). The first four volumes of Peters' Checklist (1931-1939) cite the year incorrectly as 1826, while subsequent volumes cite 1827. A few entries in volume 1 of the present work may give the incorrect date.

E. L. Layard [1824-1900] and his son L. C. Layard (dates not known)

Sharpe (1906) gives information on the Layards, father and son, and my note on H. B. Tristram in volume one mentions the former. L. C. Layard would sometimes send home to his father further specimens from the same locality soon after the original description was published. Some of these have been incorrectly labelled "type" or described as such in the Catalogue of Birds. Where this has occurred such invalid "types" have entries in the present work with the initial name not in boldface print.

Acknowledgements

Although in the first volume I expressed my thanks to the Bird Section staff as a whole, three names deserve special mention. When head of the section Sir Norman Kinnear did much work on the selection of type specimens from the older collections of Indian birds. Under the supervision of Mr. C. J. O. Harrison, Mr. S. A. Parker checked and revised the layout of every entry in Volume 1, as well as a number in this volume. For the last three years Mr. Harrison has continued this work in collaboration with me, and so enabled the task to be completed.

References

- GLADSTONE, H. S., 1937. Thomas Watling, limner of Dumfries. *Trans. Dumfriesshire and Galloway nat. Hist. Antiq. Soc.*, (3) **20**: 70–133.
- GRAY, G. R., 1843. Some rectification of the nomenclature of Australian birds. Ann. Mag. nat. Hist., 11: 189-194.
- GMELIN, J. F., 1789. Systema Naturae . . . Editio Decimo Tertia . . . cura J. F. Gmelin. 3 vols . . . Lipsiae.
- HINDWOOD, K. A., 1931. Historical associations and early records of the Emu-Wren. *Emu*, 31: 99-110.
- HINDWOOD, K. A., 1965. Sydney bird paintings. Austr. Zool., 13:86. LATHAM, J., 1801. Index Ornithologicus, 2nd. Supplement. London.
- LYSAGHT, A., 1959. Some Eighteenth Century bird paintings in the library of Sir Joesph Banks (1743–1820). Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1: 251–372.
- MATHEWS, G. M., 1942. The Lambert drawings. Enu. 42: 123-124.
- MATHEWS, G. M. & IREDALE, T., 1922. Thomas Watling, artist. Austral Avian Record, 5: 22-32, 69-73.
- PELZELN, A. von, 1871. Zur Ornithologie Brasiliens. Wien.
- Peters, J. L. et al., 1951-1968. Check-list of birds of the world. Cambridge, Mass.
- RIENITZ, R & RIENITZ, T., 1963. Early artists of Australia. Sydney.
- SAWYER, F. C., 1949–1950. Notes on some original drawings of birds used by John Latham. J. Soc. Bibliog. nat. Hist., 2: 177–179, 190.
- SCLATER, P. L., 1888–1890. Catalogue of the birds in the British Museum, vols. 14–15. London.
- SHARPE, R. B., 1906. Section 3, Birds. in, History of the collections contained in the Natural History Departments of the British Museum, 2:79-515.
- Vigors, N. A. & Horsfield, T., 1827. Australian birds in the collection of the Linnean Society. *Trans. Linn. Soc. Lond.*, 15:170-331.
- Watling, T., 1792. Original water-colour drawings of natives, animals, plants and scenery in the neighbourhood of Port Jackson, New South Wales. [In the Library of the Department of Zoology, British Museum (Natural History)].

Corrections and Additions to Vols. 1 and 2.

The corrections and additions to the preceding volume, and those relating to the present volume which were noted too late for inclusion in the main body of the text, are included as appendices on pp. 613-622 and 623 - 628.

aberdare—Cisticola robusta aberdare Lynes, 1930 .		. 35
Ibis, (12) 6, supplement: 426. HOLOTYPE, Adult male.	Reg.	no.
1923.8.7.3673. Aberdare Range, 9,000-11,000', Kenya,	26 M	arch
1912. Collected and presented by Col. Stephenson Clar	ke.	

- aberrans-Drymoica aberrans Smith, 1843 . Illustrations of the Zoology of South Africa, Aves, pl. 78. SYNTYPE, Adult female (male, vide Lynes, Ibis, (12)6, Supplement, (1930): 615). Reg. no. 1843.2.28.90 (?error in register, =.97). Near Port Natal, (South Africa, on label; Transvaal, vide White, Occ. Pap. Nat. Mus. Southern Rhodesia, 262 (1962): 660). Collected by A. Smith. Purchased of the South African Museum. See also Lynes, Ibis, (12) 6, Supplement, (1930): 615.
- abingdoni—Cactornis abingdoni P. L. Sclater and Salvin, 1870 Proc. zool. Soc. Lond., 1870: 326, fig. 5. SYNTYPE, Adult male. Reg. no. 1885.12.14.561. Abingdon Island, Galapagos Archipelago, (Dec. 1868, on label). Collected by A. Habel. Salvin-Godman Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 20.

The second syntype is also in the collection.

ablutum—Dicaeum sanguinolentum ablutum Robinson and Kloss, J.F.M.S. Mus., 11:57. SYNTYPE, Adult male. Reg. no. 1927. 4.18.691. Tamansari, 16,000', near Banjoewnagi, East Java, 20 Jan. 1920. Collected by C. B. Kloss. Presented by H. C. Robinson.

The female syntype is also in the collection.

abuensis—Dumetia albigularis abuensis Harington, 1915 J. Bombay nat. Hist. Soc., 23:429. SYNTYPE, Adult male. Reg. no. 1886.10.1.4702. Mount Abu, Deesa, India, 27 April 1875. Collected by E. A. Butler. Hume Collection. There are four other syntypes in the collection, two adult and

two immature.

- abuensis—Otocompsa jocosa abuensis Whistler, 1931 Bull. Br. Orn. Cl., 52:40. HOLOTYPE, Adult male. Reg. no. 1886.9.1.2519. Mount Abu, Rajputana, 29 April 1868. Hume Collection.
- abyssinica—Eremomela elegans abyssinica Bannerman, 1911. Bull. Brit. Orn. Cl., 29:38. HOLOTYPE, Adult male. Reg. no. 1912.10.15.840. Omo River, 2,000', Kullo, South-western Abyssinia, 24 June 1905. Collected by P. Zaphiro. Presented by W. N. Macmillan.

- abyssinica—Lusciniola abyssinica Blundell and Lovat, 1899 . . . 35

 Bull. Brit. Orn. Cl., 10: 19. Holotype, Adult male. Reg. no. 1900.1.3.109. Lake Chercher, near Harrar, South Abyssinia, 12 July 1899. Collected and presented by H. W. Blundell and Lord Lovat.
- abyssinica—Salpornis salvadorii abyssinica Ogilvie-Grant, 1913 . 44a Ibis, (10) 1:597. Syntype, Adult male. Reg. no. 1912.10.15.1148. Limmu, (c. 150 miles west of Addis Ababa, vide op. cit. 552, pl. 12, map.), 2,300', Urguessa River, Abyssinia, 30 April 1905. Collected by P. C. Zaphiro. Presented by W. N. Macmillan. The female syntype is also in the collection.
- abariensis—Pachysylvia thoracicus abariensis Chubb, 1921 . . . 56
 Birds of British Guiana, 2:395. HOLOTYPE, Adult male. Reg. no. 1922.3.5.3629. Abary River, British Guiana, July 1906. Collected by F. V. McConnell. McConnell Collection.
- abariensis—Planesticus fumigatus abariensis Chubb, 1918 31

 Bull. Br. Orn. Cl., 38: 53. Holotype, Adult. Reg. no. 1905.6.28.

 124. Abary River, British Guiana, Sept. 1906. Collected by F. V. McConnell. McConnell Collection.

There are several other syntypes in the collection. This specimen was registered as the type, but another was listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 134.

- acholiensis—Sheppardia aequatorialis acholiensis Macdonald, 1940. 31 Ibis, (14) 4:670. HOLOTYPE, Adult male. Reg. no. 1939.10.3.21. Kitibol, 6,000', Acholi Hills, Imatong Mountains, Sudan, 16 Nov. 1938. Collected and presented by F. O. Cave.
- acuminata—Haemophila acuminata Salvin and Godman, 1886. 49
 Biol. Centr.-Amer., Aves, 1:397. Syntype, Adult male. Reg.
 no. 1885.2.10.558. Yauntepec, Mexico. Collected by Deppe.
 Sclater Collection, from the Berlin Museum.

Salvin and Godman took the name from a Lichtenstein nomen nudum, Aimophila acuminata, vide Nomencl. Av. Musei Zool. Berol. (1854): 43.

- Proc. zool. Soc. Lond., 1873: 187. SYNTYPE, Adult. Reg. no. 1888.1.13.320. Bogota, Colombia. Collected for Verreaux. Sclater Collection.

 Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 95.

 The second syntype is also in the collection.
- acutirostris—Brachypteryx acutirostris Eyton, 1845 . . . 32B Ann. Mag. nat. Hist., (1) 16:228. HOLOTYPE, Adult. Reg. no. 1846.3.4.36. Malacca, Malaya. Collected and presented by A. Charlton.
- acutirostris—Calandrella acutirostris Hume, 1873 17

 In Henderson and Hume, Lahore to Yarkand: 265. HOLOTYPE,
 Adult male. Reg. no. 1887.7.1.3739. Balakchi, Yarkand, 5 Aug.
 1870. Collected by Dr. G. Henderson. Hume Collection.

There are six other syntypes in the collection.

adamsi—Montifringilla adamsi Adams, 1859 61

Proc. zool. Soc. Lond., 1858 (pub. 1859): 482. ex Moore MS.

SYNTYPE, Adult. Reg. no. 1860.1.6.1. Ladakh and Little
Tibet, July 1852 (vide Adams, Proc. zool. Soc. Lond., 1859: 178).

Collected and presented by A. L. Adams.

The single Adams specimen received from the Honourable East India Company Museum was listed by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 262, as the type, but Adams (vide Proc. zool. Soc. Lond., 1859: 178) refers to several syntypes and the above specimen presented by Adams has been selected.

- adjacens—Apalis melanocephala adjacens Clancey, 1969 . . . 35 Bull. Br. Orn. Cl., 89:93. HOLOTYPE, Adult male. Reg. no. 1946.5.3383. Njesi Plateau, c.5000', Unangu, North Mozambique, Lat. 12°45' S., Long. 35°20' E., 11 August 1945. Collected and presented by C. W. Benson.

Birds of Jamaica: 253, pl. 65 (1849). HOLOTYPE, Immature (relaxed mount). Reg. no. 1847.6.8.17. Grand Vale, Jamaica, 9 Aug. 1845. Collected by and purchased of P. H. Gosse.
adusta—Synallaxis adusta Salvin and Godman, 1884 3 Ibis, (5) 2: 450. SYNTYPE, Adult male. Reg. no. 1889.5.14.209. 6,000', Mount Roraima, British Guiana, 7 Sept. 1883. Collected by H. Whitely. Salvin-Godman Collection. There are four other syntypes in the collection.
aemodius—Conostoma aemodius Hodgson, 1841
aemodius—Parus aemodius Blyth, 1844
aeneigularis—Nectarinia aeneigularis Sharpe, 1891 46 Ibis, (6) 3:444. HOLOTYPE, Adult male. Reg. no. 1893.12.1.12. Sotik, Kenya, 6 Oct. 1889. Collected and presented by F. J. Jackson.
aequatorialis—Amaurospiza aequatorialis Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:157. HOLOTYPE, Not fully adult male. Reg. no. 1885.2.10.103. Pallatanga, Ecuador, 1859. Collected by L. Fraser. Sclater Collection.
aequatorialis—Callene aequatorialis Jackson, 1906 31 Bull. Br. Orn. Cl., 16: 46. HOLOTYPE, Adult. Reg. no. 1906.1.5.1. Kericho, 7,500', Lumbwa, Kenya, 15 June 1903. Collected and presented by F. J. Jackson.
aequatorialis—Megabias aequatorialis Jackson, 1904 38 Bull. Brit. Orn. Cl., 15:11. Syntype, Adult male. Reg. no. 1905.1.10.3. Entebbe, Uganda (4,000', 6 Oct. 1901, on label). Collected and presented as the type by F. J. Jackson. A female syntype, not in the collection, was also described.
aequatorialis—Quiscalus aequatorialis P. L. Sclater, 1862 57 Catalogue of a collection of American birds: 140. Syntype, Adult male. Reg. no. 1885.7.12.334. Babahoyo, west Ecuador. Aug. 1859. Collected by L. Fraser. Sclater Collection. The female syntype is also in the collection.

Proc. zool. Soc. Lond., 1961: 380. SYNTYPE, Adult. Reg. no. 1889.9.20.178. Rio Napo, Ecuador (type locality restricted by Chapman, Bull. Amer. Mus. nat. Hist., 36 (1917): 369). Collected for Verreaux. Sclater Collection. The other syntype is in the collection. aequinoctialis—Cyclorhynchus aequinoctialis P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858: 70. HOLOTYPE, Adult. Reg. no. 1888.1.13.694. Napo River, eastern Ecuador, Collected for Verreaux. Sclater Collection. 37 aestigma—Muscicapa aestigma J. E. Gray, 1846. Catalogue of mammalia and birds of Nepal and Thibet presented by Hodgson to the British Museum: 90, 155. SYNTYPE, Adult (male, vide, Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 206). Reg. no. 1843.1.13. 684. (Collector's no. 141). Nepal. Collected and presented by B. H. Hodgson. Listed as the type by Sharpe (op. cit.). There are several other syntypes in the collection. aethiopica—Hirundo aethiopica Blanford, 1869 18 Ann. Mag. nat. Hist., (4) 4:329. HOLOTYPE, Adult (female on label). Reg. no. 1869.10.16.4. (Barakit, 7,500', Tigré, Abyssinia, 24 May 1868 on label). Collected by W. T. Blanford and presented by the Governor and Council of the Bombay Government. aethiops—Oryzoborus aethiops P. L. Sclater, 1860 Proc. zool. Soc. Lond., 1860: 88. HOLOTYPE, Adult male. Reg. no. 1888.2.10.95. Nanegal, West Ecuador, Feb. 1859. Collected by L. Fraser. Sclater Collection. æthiops—Poecilodryas aethiops P. L. Sclater, 1880 31 Proc. zool. Soc. Lond., 1880: 66, pl. 7. HOLOTYPE, Adult male. Reg. no. 1881.3.29.19. Kabakadai, New Britain, Jan. 1879. Collected by the Rev. G. Brown and purchased of Gerrard. aethiops—Thamnophilus aethiops P. L. Sclater, 1858. Proc. zool. Soc. Lond., 1858: 65. SYNTYPE, Adult male. Reg. no. 1889.9.20.37. Rio Napo, Ecuador. Probably collected by Villavicencio. Sclater Collection; obtained by Sclater from Verreaux. There is a female syntype in the collection. affinis—Abrornis affinis Moore, 1854. in Horsfield and Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 1: 341. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.3042. Nepal. Collected by B. H.

Hodgson. Presented by the Honourable East India Company.

aequatorialis—Thannistes aequatorialis P. L. Sclater, 1861.

Bull. Brit. Orn. Cl., 16:116. SYNTYPE, Adult male. Reg. no. 1906.12.23.1022. (Mubuku Valley, 11 Jan. 1906, on label), Eastern Ruwenzori. Collected by R. E. Dent. Presented by the subscribers to the Ruwenzori Expedition. The female syntype is also in the collection. affinis—Cimyris affinis Horsfield, 1821	offinis Anglis affinis Ogilvio Grant 1006	25
Trans. Linn. Soc. Lond., 13:166. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4786. (Allocated in 1961). Java. Collected by T. Horsfield. Presented by the Honourable East India Company Museum. In very bad condition. Although Gadow, Cat. Bds. Brit. Mus., 9 (1884): 108, refers to two specimens, this is the only one that can be traced. affinis—Drymoica affinis Smith, 1843	Bull. Brit. Orn. Cl., 16:116. SYNTYPE, Adult male. Reg. no. 1906.12.23.1022. (Mubuku Valley, 11 Jan. 1906, on label), Eastern Ruwenzori. Collected by R. E. Dent. Presented by the subscribers to the Ruwenzori Expedition.	,
Trans. Linn. Soc. Lond., 13:166. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4786. (Allocated in 1961). Java. Collected by T. Horsfield. Presented by the Honourable East India Company Museum. In very bad condition. Although Gadow, Cat. Bds. Brit. Mus., 9 (1884): 108, refers to two specimens, this is the only one that can be traced. affinis—Drymoica affinis Smith, 1843	affinis—Cimyris affinis Horsfield, 1821	46
In very bad condition. Although Gadow, Cat. Bds. Brit. Mus., 9 (1884): 108, refers to two specimens, this is the only one that can be traced. affinis—Drymoica affinis Smith, 1843	Trans. Linn. Soc. Lond., 13: 166. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4786. (Allocated in 1961). Java. Collected by T. Horsfield. Presented by the Honourable East India Company	
Illustrations of the Zoology of South Africa, Aves, pl. 77. SYNTYPE, female (immature, vide Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 195). Reg. no. 1843.2.28.109. South Africa. (=Rustenberg; vide Roberts, Ann. Transvaal Mus., (1922); 242). Collected by A. Smith. Purchased of the South Africa Society. Listed as the type by Sharpe, op. cit., but there is no information regarding the number of specimens available. affinis—Garrulax affinis Blyth, 1843	In very bad condition. Although Gadow, Cat. Bds. Brit. Mus., 9 (1884): 108, refers to two specimens, this is the only one that can	
female (immature, vide Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 195). Reg. no. 1843.2.28.109. South Africa. (=Rustenberg; vide Roberts, Ann. Transvaal Mus., (1922); 242). Collected by A. Smith. Purchased of the South Africa Society. Listed as the type by Sharpe, op. cit., but there is no information regarding the number of specimens available. affinis—Garrulax affinis Blyth, 1843	affinis—Drymoica affinis Smith, 1843	35
Reg. no. 1843.1.13.553. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection. The author used Hodgson's MS name. affinis—Hapalophus affinis G. R. Gray, 1837	female (immature, vide Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 195). Reg. no. 1843.2.28.109. South Africa. (=Rustenberg; vide Roberts, Ann. Transvaal Mus., (1922); 242). Collected by A. Smith. Purchased of the South Africa Society. Listed as the type by Sharpe, op. cit., but there is no information	
Reg. no. 1843.1.13.553. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection. The author used Hodgson's MS name. affinis—Hapalophus affinis G. R. Gray, 1837	official Commutation official Direction 1942)D
Mag. nat. Hist., (n.s.) 1:489. Holotype (Adult male, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883):142). Reg. no. 1837.6.10.407. Zanzibar. Purchased at Baron Laugier's sale in Paris. affinis—Hemignathus affinis Rothschild, 1893	Reg. no. 1843.1.13.553. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection. The author	7
Mag. nat. Hist., (n.s.) 1:489. Holotype (Adult male, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883):142). Reg. no. 1837.6.10.407. Zanzibar. Purchased at Baron Laugier's sale in Paris. affinis—Hemignathus affinis Rothschild, 1893	- Control Hamilton of Control 1927	22
 Ibis, (6) 5: 112. SYNTYPE, Adult. Reg. no. 1939.12.9.142. Collector's number 1698). Mauai Island, Sandwich Islands, (9 Aug. 1892, on label). Collected by H. C. Palmer. Rothschild Bequest. No holotype was designated. The specimen listed as the type by Hartert, Novit. Zool., 26 (1919): 170 was one of a series of syntypes which included the above specimen. affinis—Mirafra affinis Blyth, 1845	Mag. nat. Hist., (n.s.) 1:489. HOLOTYPE (Adult male, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883):142). Reg. no. 1837.6.10.407.	,
 Ibis, (6) 5: 112. SYNTYPE, Adult. Reg. no. 1939.12.9.142. Collector's number 1698). Mauai Island, Sandwich Islands, (9 Aug. 1892, on label). Collected by H. C. Palmer. Rothschild Bequest. No holotype was designated. The specimen listed as the type by Hartert, Novit. Zool., 26 (1919): 170 was one of a series of syntypes which included the above specimen. affinis—Mirafra affinis Blyth, 1845	affinis—Hemignathus affinis Rothschild, 1893	55
J Asiat Soc. Bengal, 13 (1844, pub. 1845): 959. The two Jerdon specimens listed as types by Sharpe, Cat. Bds.	Ibis, (6) 5:112. SYNTYPE, Adult. Reg. no. 1939.12.9.142. Collector's number 1698). Mauai Island, Sandwich Islands, (9 Aug. 1892, on label). Collected by H. C. Palmer. Rothschild Bequest. No holotype was designated. The specimen listed as the type by Hartert, Novit. Zool., 26 (1919): 170 was one of a series of syntypes	
J Asiat Soc. Bengal, 13 (1844, pub. 1845): 959. The two Jerdon specimens listed as types by Sharpe, Cat. Bds.	affinis—Mirafra affinis Blyth, 1845	17
Brit. Mus., 13 (1890): 615 are presumably those described by Jerdon	J Asiat Soc. Bengal, 13 (1844, pub. 1845): 959.	

as Mirafra	affinis in the	Madras J.	Lit. Sci., 1	3, vol. 3	1 (1844,	pub.
1846): 136.						
known from	what specin	nens the for	mer made	his desc	ription.	

- male. Reg. no. 1873.5.12.1284. Gagie Island (off western New Guinea). Collected by A. R. Wallace. Wallace Collection.
- affinis—Myiothera affinis Horsfield, 1821 12 Trans. Linn. Soc. Lond., 13:145. SYNTYPE, Adult. Reg. no. 1880 1.1.4782. (allocated at a later date). Java. Presented by the Honourable East India Company Museum.

The specimen is a partly-relaxed mount almost completely devoid of feathers but is the specimen referable to the period of the descrip-

tion

- affinis—Phoenicornis affinis McClelland, 1840 20 in Horsfield, Proc. zool. Soc. Lond., 1839 (pub. 1840): 157. Syntype, Adult female (relaxed mount, tail incomplete). Reg. no. 1880.1.1.2757. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum. There is another syntype in the collection.
- affinis—Phyllopneuste affinis Hodgson, 1844 in J. E. Gray, Zoological Miscellany; 82; and also J. E. Gray, List of Mammalia and birds of Nepal and Thibet presented by B. H. Hodgson . . . to the British Museum (1846): 65. These are nomina nuda and the specimen listed as the type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 65, has no type status.
- affinis—Poecile affinis Prievalski, 1876 42 Mongol. i strana Tangut., 2:52. SYNTYPE, Adult male. Reg. no. 1901.5.4.75. Kansu, Mongolia, (February, on label). Collected by N. Prjevalski. Purchased of M. Menzbier as the type. There is a translation of the description by F. C. Craemers in Rowley, Ornithological Miscellany 2 (1877): 188.
- affinis—Poocaetes gramineus affinis Miller, 1888. Auk, 5:404. HOLOTYPE, Reg. no. 1906.12.7.3737. Salem, Oregon, 29 May 1887. Collected by and purchased of G. S. Miller. The specimen was not received until 1937 but was given the registration number of the original collection.
- affinis—Salicaria affinis J. E. Gray, 1846 Catalogue of Mammalia and Birds of Nepal and Thibet presented by B. H. Hodgson to the British Museum: 64, 151. SYNTYPE, Adult. Reg. no. 1845.1.9.836. Nepal. Collected by a native collector for B. H. Hodgson. Presented by B. H. Hodgson. There are other syntypes in the collection.

affinis—Xanthopygia affinis Ogilvie-Grant, 1906
This is the author's selected female. His selected male and other syntypes are also in the collection.
africana—Mirafra africana Smith, 1836
africanoides—Mirafra africanoides Smith, 1836 1834; Report of the Expedition for exploring Central Africa 1834; Appendix: 47. SYNTYPE, Adult male (relaxed mount). Reg. no. 1838.6.9.89. Eastern province of the colony and country towards Latakoo (in text; type locality restricted to Colesberg by Macdonald, Contributions to the Ornithology of western South Africa, 1957: 95, to Litakun, near Kuruman, by White, Bull. Br. Orn. Cl., 80 (1960): 11, and back to Colesberg by Clancey, Bull. Br. Orn. Cl., 80 (1960): 103). Collected by A. Smith and purchased of the South African Museum. There are two other syntypes in the collection; other syntypes are at the Academy of Natural Sciences, Philadelphia.
agilis—Anthus agilis Sykes, 1832
agilis—Euscarthmus agilis P. L. Sclater, 1856
agnatus—Furnarius agnatus P. L. Sclater and Salvin, 1873
agricola—Sylvia (Acrocephalus) agricola Jerdon, 1844 35 Madras J. Lit. Sci., 13, pt. 2.: 131. SYNTYPE, Adult. Reg. no. 1846.4.30.15. Near Nellore. Collected and presented by T. C. Jerdon.

This specimen was labelled as the type but there is another syntype in the collection.
agricolensis—Jerdonia agricolensis Hume, 1870
 alakulensis—Budytes thumbergi alakulensis C. H. B. Grant and Mackworth-Praed, 1949
alberti—Menura alberti Bonaparte, 1850
albescens—Pachyramphus albescens Gould, 1839
albicaudata—Muscicapa albicaudata Jerdon, 1840

A second specimen from the same region was received at a later

date.

albiceps—Turdus albiceps Swinhoe, 1864 . Ibis, (1) 6:363. SYNTYPE, Adult male. Reg. no. 1896.6.1.656. (Tamsuy, Formosa, 15 March 1864, vide Swinhoe, Ibis, (2) 2 (1866): 135, pl. 5). Collected by R. Swinhoe. Seebohm Collection. Listed as type by Swinhoe, 1866 (loc. cit.). The two other syn-

types cannot be traced in the collection. (See also appendix, Corrections and Additions, vol. 2.)

albicilius—Buglodytes albicilius Verreaux, 1854 . C. r. Acad. Sci., Paris, 38:57. SYNTYPE, Adult. Reg. no. 1854.1. 17.59. Santa Marta, New Granada. Collected for and purchased of Verreaux.

- albicollis—Ixulus flavicollis albicollis Ticehurst and Whistler, 1924. 32B Bull. Br. Orn. Cl., 44:71. HOLOTYPE, Adult male. Reg. no. 1924.3.13.2. Dharmsala, 4,000', Kangra, north-west Himalayas, 20 Nov. 1921. Collected and presented by H. Whistler.
- albicrissalis—Bradypterus alfreda albicrissalis Neumann, 1914 . 35 Orn. Monats., 22:10. Holotype, Adult female. Reg. no. 1906.12.23.1008. Mubuku Valley, (5,000', on label), East Ruwenzori Mountains, Africa, 5 April 1907. Collected by G. Legge on the Ruwenzori Expedition.
- albidiventris—Cinclodes albidiventris P. L. Sclater, 1860. . . . 3

 Proc. zool. Soc. Lond., 1860: 77. SYNTYPE, Adult (male, vide Sclater, Cat. Bds. Brit. Mus., 15 (1890): 24). Reg. no. 1889.5.20.74. Panza, Chimborazo, Ecuador, 1859. Collected by L. Fraser. Sclater Collection.
 - There are three other syntypes in the collection.
- albifrons—Chlorospingus albifrons Salvin and Godman, 1889 . 5.

 Ibis, (6) 1:237. SYNTYPE, Adult male. Reg. no. 1894.7.1.913.

 Omilteme, 8,000', Sierra Madre del Sur, Guerrero, Mexico, July 1888. Collected by Mrs H. H. Smith. Salvin-Godman Collection.

 There are several other syntypes in the collection.
- albifrons—Machaerirhynchus albifrons G. R. Gray, 1861 . . . 3

 Proc. zool. Soc. Lond., 1861: 429, pl. 43. SYNTYPE, Adult male.

 Reg. no. 1873.5.12.692. Waigiou Island, off North-west New Guinea, 1860. Collected by A. R. Wallace. Wallace Collection.

 Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 392.

 This is the only specimen from this locality but three other syntypes from Mysol, also part of the type locality, are in the Collection.

- albifrons—Pericrocotus albifrons Jerdon, 1862 Ibis, (1) 4:20. Syntype, Adult male. Reg. no. 1862.1.17.14. Thavetmyo, Upper Burma, Collected and presented by T. C. Ierdon The female syntype is also in the collection. albifrons—Platystira albifrons Sharpe, 1873 38 Ibis, (3) 3:159. SYNTYPE, Adult. Reg. no. 1873.12.10.61. Loge River, Angola. Collected by and purchased of J. J. Monteiro. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 147. There is a second syntype in the collection. albifrons—Setophaga albifrons P. L. Sclater and Salvin, 1871 Proc. zool. Soc. Lond., 1870 (publ. 1871): 784. SYNTYPE, Adult male. Reg. no. 1884.5.15.497. Upper wood—region of Merida, Venezuela. Collected by A. Goering. Sclater Collection.

 Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 423. There is another syntype in the collection. albigularis—Empidonax albigularis P. L. Sclater and Salvin, 1859 . 9 Ibis, (1) 1:122. HOLOTYPE, Adult male. Reg. no. 1888,1.1.1448. Dueñas, Guatamala, 1 Feb. 1858. Collected by O. Salvin, Salvin-Godman Collection. 1886.9.15.79. Rio Napo, Quixos Province, east Ecuador. Collected for Verreaux, probably by Cillavicencio. Sclater Collection. albigularis—Leptasthenura striata albigularis Morrison, 1938. Ibis, (14) 2:775. HOLOTYPE, Adult female. Reg. no. 1939.6.17.1. Lircay, 10,660', Huancavelica Dept., central Peru, 29 Nov. 1937. Collected by and purchased of A. Morrison. albigularis—Nemosia albigularis P. L. Sclater, 1855 Proc. zool. Soc. Lond., 1855: 109, pl. 99. SYNTYPE, Adult male. Reg. no. 1885.6.12.816. Bogota. Collected by E. Parzudaki. Sclater Collection. Listed as the type by Sclater, Catalogue of a collection of American birds. (1862): 87. The other syntypes are in the collection. albigularis—Philydor albigularis Salvin and Godman, 1884 Ibis, (5) 2:450. SYNTYPE, Adult male. Reg. no. 1889.5.14.348. 6,000', Roraima, British Guiana, 5 Sept. 1883. Collected by
- albigularis—Sclerurus albigularis P. L. Sclater and Salvin, 1869. . 3

 Proc. zool. Soc. Lond., 1868 (publ. 1869): 630. HOLOTYPE, Adult.

H. Whitely. Salvin-Godman Collection.

There are several other syntypes in the collection.

- Reg. no. 1889,5.20,416. Venezuela, 1868 (Cumbre de Valencia, vide Hellmayr, Catalogue of Birds of the Americas, 4 (1925): 247). Collected by A. Goering. Sclater Collection. The authors used Swainson's MS, name. albigularis—Synallaxis albigularis P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858: 63. Schater, Catalogue of a collection of American Birds, 1862: 151, referred to specimen B.M. Reg. no. 1889.5.20.135 from Zamora, Ecuador, as the type. This is in fact the specimen described by Sclater in Proc. zool. Soc. Lond., 1858: 456, and has no type status. albigularis—Xenocichla albigularis Sharpe, 1881. Cat. Bds. Brit. Mus., 6: 103, pl. 7. HOLOTYPE, Adult. Reg. no. 1876.5.23.649. Fantee, West Africa. Collected by Ussher. Sharpe Collection. albiloris—Grallaria albiloris Taczanowski, 1880 . Proc. zool. Soc. Lond., 1880: 201. SYNTYPE, Adult male. Reg. no. 1889.6.20.635. Callacate, 4,800-5,200', northern Peru, 2 May 1879. Collected by Stolzmann. Sclater Collection. Listed as "authentic specimen" by P. L. Sclater, Cat. Bds. Brit. Mus., 15 (1890): 321. albiloris—Pachyrhamphus albiloris Chubb, 1920. Bull. Brit. Orn. Cl., 40:73. HOLOTYPE, Adult male. Reg. no. 1888.1.13.1515. San Estaban, Carabobo, Venezuela (1871, on label). Collected by A. Goering. Sclater Collection. albiloris—Polioptila albiloris P. L. Sclater and Salvin, 1860 Proc. zool. Soc. Lond., 1860: 298. HOLOTYPE, Adult. (male, on label). Reg. no. 1885.3.2.612. (Chuacus, vide Salvin, Ibis, (1) 2 (1860): 397). Vera Paz, Montagua Valley, Guatemala, Feb. 1860. albinuchalis—Thannophilus albinuchalis P. L. Sclater, 1855 . Proc. zool. Soc. Lond., 1855: 18. SYNTYPE, Adult male (relaxed mount). Reg. no. 1850.1.31.56. Guayaquil, south-west Ecuador. Collected and presented by Capt. H. Kellett and Lt. Wood of H.M.S. There is another syntype in the collection.

drawing number on the label and is therefore likely to have been

before him when the description was made.

- albispecularis—Tachyphorus albispecularis Leotaud, 1866 Oiseaux de l'Ile de la Trinidad: 300. SYNTYPE, Adult male. Reg. no. 1886.6.12.734. Trinidad. Sclater Collection.

 Labelled "one of the types". See also Sclater, Ibis, (2) 3 (1867): 108
- albitorquata—Amadina fasciata albitorquata Vincent 1933 Bull. Brit. Orn. Cl., 53: 148. HOLOTYPE, Adult male. Reg. no. 1933.3.1.18. Fort Johnston, 1,700', Nyasaland, 1 May 1932. Collected and presented by J. Vincent.
- albitorques—Philemon albitorques P. L. Sclater, 1877 Proc. zool. Soc. Lond., 1877: 553. HOLOTYPE, Adult male. Reg. no. 1880.11.18.99. Wild Island, Admiralty Islands, South-west Pacific. Collected by J. Murray, from the voyage of H.M.S. Challenger.
- albitorquis—Spermophila albitorquis Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:120. SYNTYPE, Adult male. Reg. no. 1885.12.14.402. Capulalpam, South Mexico, Oct. 1860. Collected by A. Sallé. Salvin-Godman Collection.

The second syntype is also in the collection, but the measurements of the selected syntype are more accurate (vide A. R. Phillips, in prep. 1970).

albiventer—Tesia albiventer Hodgson, 1837. J. Asiat. Soc. Bengal, 6:102. SYNTYPE, Adult (relaxed mount). Reg. no. 1843.1.13.418. Nepal. Collected and presented by B. H. Hodgson.

There are other syntypes in the collection.

- albiventris—Callene albiventris Blanford, 1867 . Proc. zool. Soc. Lond., 1867: 833, pl. 39. SYNTYPE, Adult male. Reg. no. 1869.9.15.2. Pulney Hills, 6,000-7,000', south India. Collected and presented by the Rev. S. Fairbank. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 16. Blanford used Fairbank's MS name.
- albiventris—Camaroptera brevicaudata albiventris Granvik, 1934 . 35 Revue Zool. Bot. afr., 25:101. HOLOTYPE, Adult male. Reg. no. 1887.11.3.84. Manda Islands, Kenya. Collected and presented by F. J. Jackson.
- albiventris—Cyclorhis albiventris P. L. Sclater and Salvin, 1873 Nomencl. Av. Neotrop.: 156. HOLOTYPE, Adult. Reg. no. 1886. 9.15.344. Bahia, Brazil. Collected by Wucherer. Sclater Collec-
- albiventris—Hyloterpe albiventris Ogilvie-Grant, 1894 Bull. Brit. Orn. Cl., 3:49. SYNTYPE, Adult male. Reg. no. 1897. 5. 13. 110. La Trinidad, Benguet District, north-west Luzon, Philippines, 5 March 1894. Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. There is a second syntype in the collection.

albiventris—Loxia albiventris Swinhoe, 1870
albiventris—Minnocichla ardesiaca albiventris P. L. Sclater, 1889 . 31 Proc. zool. Soc. Lond., 1889: 326. SYNTYPE, Adult male. Reg. no. 1889.6.10.4. Batalie, Dominica, West Indies, 8 March 1889. Collected by G. A. Ramage and presented by the West Indies Committee of the Royal Society. The second syntype is also in the collection.
albiventris—Monarcha albiventris Gould, 1866
albiventris—Niornis albiventris Godwin-Austen, 1876 32B J. Asiat. Soc. Bengal, 45, pt. 2:199. Holotype, Adult. Reg. no. 1895.7.14.1949. Sengmai, Manipur Valley, North-east Frontier, India. Collected by W. Robert. Godwin-Austen Collection.
albiventris—Parus albiventris Shelley, 1881
albiventris—Rhamphocaenus albiventris P. L. Sclater, 1883 34 Ibis, (5) 1:95. SYNTYPE, Adult. Reg. no. 1889.9.20.388. Surinam, Guiana. Collected by C. Bartlett. Sclater Collection. Listed as one of the types by Sclater, Cat. Bds. Brit. Mus., 15 (1890):261.
albocinctus—Acridotheres albocinctus Godwin-Austen and Walden, 1875
albocinctus—Turdus albocinctus Royle, 1839 31 Illustrations of Botany and natural History of the Himalayan Mountains: pl. 8. Holotype, Adult male (relaxed mount). Reg. no. 1880.1.1.1988. (Bengal on label). Collected by J. F. Royle and presented by the Honourable East India Company Museum.
albocinereus—Syristes albocinereus P. L. Sclater and Salvin, 1880. 9 Proc. zool. Soc. Lond., 1880: pl. 156. SYNTYPE, Adult. Reg. no. 1888.1.13.765. Bogota, Colombia. Collected for Verreaux. Salvin-Godman Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 183. There are two other syntypes in the collection.

albocoronata—Serpophaga albocoronata Gould, 1839 Zoology of the voyage of H.M.S. Beagle, 3, Birds: 49. SYNTYPE. Adult. Reg. no. 1837.2.21.344. Maldonado. Collected by Capt. Fitzroy, R.N., of *H.M.S. Beagle*. Presented by Sir W. Burnett and Capt. Fitzrov. Listed as the type by Hellmayr, Catalogue of birds of the Americas.

5 (1927) : 382.

- albofasciatus—Dryoscopus albofasciatus Sharpe, 1891. Ibis, (6) 3:598. SYNTYPE, Adult male. Reg. no. 1893.12.1.16. Mount Elgon, 6,000', East Africa, 12 Feb. 1890. Collected and presented by F. J. Jackson.
- albofrontata—Gervgone? albofrontata J. E. Grav. 1844 The Zoology of the voyage of H.M.S. Erebus and Terror, 1, Birds (1844): 5, pl. 4. HOLOTYPE, Adult (relaxed mount). Reg. no. 1842.9.29.25. "Brought from New Zealand" (Chatham Island, in British Museum Register). Collected by Dieffenbach. Presented by the New Zealand Company.
- albofrontata—Hetaerornis albofrontata Layard, 1854 62 Ann. Mag. nat. Hist., (2) 13:217. SYNTYPE, Adult (relaxed mount). Reg. no. 1853.5.3.34. Cevlon. Collected by Thwaites. Purchased of Cumming.

This specimen was in the collection but omitted from Sharpe,

Cat. Bds. Brit. Mus., 13 (1890): 56.

- albogrisea—Serphophaga albogrisea P. L. Sclater and Salvin, 1880. Proc. zool. Soc. Lond., 1880: 156. HOLOTYPE, Adult. Reg. no. 1888.1.1.473. Sarayacu, Ecuador. Collected by C. Buckley. Salvin-Godman Collection.
- albogriseus—Pachyrhamphus albogriseus P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1857: 78. SYNTYPE, Adult male. Reg. no. 1888.1.13.1511. Bogota, Colombia. (Collected by E. Parzudaki, on label). Sclater Collection. The female syntype is also in the collection.
- albogularis—Abrornis albogularis Moore, 1854 in Horsfield and Moore, Catalogue of birds in the Museum of the Honourable East India Company, 1: 340. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.3059. Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company Museum.

The second syntype is also in the collection.

albogularis—Crithagra albogularis Smith, 1833 South African Quarterly Journal, 2:48. HOLOTYPE, Adult (relaxed mount). Reg. no. 1838.6.9.76. South Africa. Collected by A. Smith. Purchased of the South Africa Society.

albogularis—Platyrhynchus albogularis P. L. Sclater, 1860 9 Proc. zool. Soc. Lond., 1860: 68. Holotype, Adult male. Reg. no. 1888.1.13.223. Pallatanga, Ecuador (Oct. 1858, on label). Collected by L. Fraser. Sclater Collection.
albogularis—Prinia albogularis Walden, 1870
albogularis—Suya albogularis Hume, 1873
alboides—Motacilla alboides Hodgson, 1836
alboniger—Saxicola alboniger Hume, 1873
albonotatus—Trochocercus albonotatus Sharpe, 1891
alboolivacea—Cyornis alboolivacea Hume, 1877
alboplagatus—Dryoscopus alboplagatus Jackson, 1906 23 Bull. Br. Orn. Cl., 16:90. Holotype, Adult (female on label). Reg. no. 1906.6.3.2. Doingo Narok, Kenya, 26 March 1906. Collected and presented by F. J. Jackson.
albosuperciliaris—Garrulax albosuperciliaris Godwin-Austen, 1874. 321 Proc. zool. Soc. Lond., 1874: 45. HOLOTYPE, Adult male. Reg. no. 1895.7.14.2257. Munipur Valley, near Kaibi, north-east Bengal. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.
albosuperciliaris—Suya albosuperciliaris Hume, 1871 35 in Handerson and Hume, Ibis, (3) 1:409. Holotype, Adult (female, on label). Reg. no. 1873.6.30.12. Koshtak, plains of Yarkand, 10 Sept. 1870. Collected by G. Henderson. Purchased of Gerrard.

aldabranus—Nesillas aldabranus Benson and Penny, 1968 . . . 35
Bull. Brit. Orn. Cl., 88: 102. HOLOTYPE, Adult female. Reg. no. 1968.43.1. ½ mile from western extremity of Middle Island (Johnny Channel or Gionnet), Aldabra Atoll, Indian Ocean, 11 Dec. 1967. Collected by M. J. Penny. From the Royal Society Aldabra Expedition. alexanderi-Eremomela flaviventris alexanderi W. L. Sclater and 35 Bara, Kordofan, Sudan, 27 April 1904. Collected and presented by A. L. Butler. alexanderi—Nigrita alexanderi Ogilvie-Grant, 1907 60 Bull. Brit. Orn. Cl., 21:16. SYNTYPE, Adult male. Reg. no. 1904.6.22.325. Fernando Po. (Banterbari, Bubu Town, on label), 6 Feb. 1904. Collected by E. Seimund. Presented by Mrs P. Sladen, the Duke of Bedford, and the Hon. W. Rothschild. The author labelled this specimen as the type but there are several other syntypes in the collection. alexanderi—Ploceus (Othyphantes) baglafecht alexanderi Sclater, 1925 61 Bull. Br. Orn. Cl., 46:15. HOLOTYPE, Adult male. Reg. no. 1911.12.23.3682. Gudima, Iri River, north-east Belgian Congo, 5 Sept. 1906. Collected and presented by Boyd Alexander. alexanderi-Poliolais alexanderi Bannerman, 1915 35 Bull. Brit. Orn. Cl., 35:53. HOLOTYPE, Adult male. Reg. no. 1911.12.23.4196. Cameroon Mountains, West Africa, 6 May 1909. Collected by W. Boyd Alexander. Boyd Alexander bequest. 31 no. 1901.10.29.27. Efulen, Cameroons, 6 June 1901. Collected by and purchased of G. L. Bates. alexandri—Lobornis alexandri Sharpe, 1874 Ann. Mag. nat. Hist., (4) 14:64. HOLOTYPE, (immature, vide Sclater, Systema avium Aethiopicarum, (1930):773). Reg. no. 1874.5.18.3. Old Calabar, Nigeria. Collected by A. Goldie. Presented by Dr. A. Smith. alfredi—Bleda alfredi Shelley, 1903 . Bull. Br. Orn. Cl., 13:61. SYNTYPE, Adult male. Reg. no. 1903.4. 20.8. Mwenembe, 7,500', northern Nyasaland, Sept. 1902. Collected by McClounie and presented by A. Sharpe. There are four other syntypes in the collection. alfredi-Cyanomitra olivacea alfredi Vincent, 1934 Ibis, (13) 4:90. HOLOTYPE, Adult male. Reg. no. 1933.3.1.26. Namuli Mountains, 4,600', Quelimane Province, Portuguese East Africa, 25 July 1932. Collected and presented by J. Vincent.

10 algerrensis	
algeriensis—Ammomanes algeriensis Sharpe, 1890 16 Cat. Bds. Brit. Mus., 13:645. SYNTYPE, Adult male. Reg. no. 1889.9.13.186. Algerian Sahara, 1867. Collected by Loche. Shelley Collection; obtained by Shelley from Schneider of Basel. There are several other syntypes in the collection.	7
aliena—Rhodopechys aliena Whitaker, 1897	9
aliena—Sitagra aliena Sharpe, 1902	1
alinae—Cyanomitra alinae Jackson, 1904	
alleni—Coereba chloropyga alleni P. R. Lowe, 1912 5. Bull. Brit. Orn. Cl., 29:86. SYNTYPE, Adult male. Reg. no. 1905.1.30.100. Chapade, Matto Grosso, Brazil. 22 April 1885. Collected by H. H. Smith. Sclater Collection. The female syntype is also in the collection.	4
almorae—Sitta castaneoventris almoare Kinnear and Whistler, 1930 4. Bull. Brit. Orn. Cl., 51:27. HOLOTYPE, Adult male. Reg. no. 1886.11.1.1917. Valley of Ramganga, between Almorah and Pethora, Himalayas, 29 Oct. 1866. Hume Collection.	3
alopex—Mirafra alopex Sharpe, 1890	7
alpina—Cryptolopha alpina Ogilvie-Grant, 1906	5
alpina—Taenioptera alpina Jardine, 1849	9

	•
alpinus—Coccothraustes alpinus Sharpe, 1888 Cat. Bds. Brit. Mus., 12:38. Sharpe referred to a specimen by this name which had been inscribed on the label by M. Parzudaki of whom it was purchased. It is obvious that Sharpe was relegating it to synonymy but in doing so he quoted the name and description and listed it as "s.n.". The specimen reg. no. 1852.5.27.5. cannot now be traced.	
altera—Alaemon hamertoni altera Witherby, 1905	
altera—Pyrrhula altera Rippon, 1906	
Alterum—Edoliosoma alterum R. G. W. Ramsay, 1881	
Alterum—Trichastoma abbotti alterum Sims, 1957	В
Stray Feathers, 7:60. SYNTYPE, Adult male. Reg. no. 1886.7.8. 156. Jhansie, Kashmir, (29 Sept. 1868, on label). Hume Collection. This specimen was labelled as the type. There are four other syntypes in the collection.	35
Alticola—Catharus alticola Salvin and Godman, 1879	31

alticola—Chamaeospiza alticola Salvin and Godman, 1889 . . . 49 Ibis, (6) 1:381. Syntype, Adult male. Reg. no. 1899.2.1.2872. Sierra Nevada de Colima, 10,000', western Mexico, 4 May. Collected by W. Lloyd and W. B. Richardson. Salvin-Godman Collection.

There are two other syntypes in the collection.

Listed as the type by Hellmayr and Conover, <i>Catalogue of birds of the Americas</i> , 11 (1938): 453. There are several other syntypes in the collection.
alticola—Cisticola alticola Shelley, 1899
alticola—Cistothorus alticola Salvin and Godman, 1883
alticola—Junco alticola Salvin, 1863
alticola—Poospiza alticola Salvin, 1895
altirostre—Chrysomma altirostre Jerdon, 1862 32B Ibis, (1) 4: 22. HOLOTYPE, Adult. Reg. no. 1862.1.17.18. Islands of Burrampootra River, Thayetmyo, Upper Burma. Collected and presented by T. C. Jerdon.
altirostris—Cyclorhis altirostris Salvin, 1880
amabilis—Cotinga amabilis Gould, 1857
amabilis—Malurus amabilis Gould, 1850

- that the type is in the British Museum, but the Gould specimens from Cape York in the collection are of a later date.
- amadoni—Hirundo aethiopica amadoni White, 1956 18

 Bull. Br. Orn. Cl., 76: 160. Holotype, Adult male. Reg. no. 1923.8.7.4014. Geloher, 3,000′, Somaliland, 17 March 1919. Collected by Sir G. Archer and presented by Col. Stephenson Clarke.
- amadonis—Coracina caledonica amadonis Cain and Galbraith, 1955 20 Bull. Br. Orn. Cl., 75:90. HOLOTYPE, Adult female. Reg, no. 1959.21.434. Turipava, above 3,500', Guadacanal, Solomon Islands, 16 Aug. 1953. Collected by a native for the Oxford University (Department of Zoology) Expedition to the Solomon Islands.
- amani—Dioptrornis fischeri amani W. L. Sclater, 1931 37 Bull. Brit. Orn. Cl., 51:112. HOLOTYPE, Adult female. Reg. no. 1931.5.7.5. Near Amani, 1,300', Tanganyika, 28 Jan. 1931. Collected and presented by R. E. Moreau.
- amani—Oriolus chlorocephalus amani Benson, 1946 63

 Bull. Brit. Orn. Cl., 67: 26. HOLOTYPE, Adult male. Reg. no. 1931.12.19.242. Amani, Usambara Mountains, 1,600', Tanganyika, 21 March 1931. Collected by R. Moreau. Presented by the Salvin-Godman Fund.
- amauropteryx—Hypochaera amauropteryx Sharpe, 1890 . . . 61 Cat. Bds. Brit. Mus., 13:309. SYNTYPE, Adult male. Reg. no. 1875.10.7.32. Rustenberg, East Transvaal, Jan. 1875. Collected by and purchased of F. A. Barratt.

The specimen is labelled by the author as the described specimen and type, but there are five other syntypes in the collection.

Listed as the type by Hellmayr, Catalogue of birds of the Americas, 11 (1938): 212. The three described female syntypes are also in the collection.

amzonicus—Thannophilus amazonicus P. L. Sclater, 1858 . . . 4

Proc. zool. Soc. Lond., 1858: 214, pl. 139. Syntype, Adult male.

Reg. no. 1889.9.20.89. Rio Javarri, Upper Amazon (vide Hellmayr.

Catalogue of birds of the Americas, 3 (1924): 98). Collected by

H. W. Bates. Sclater Collection.

There are three other syntypes in the collection.

amazonicus—Thryothorus amazonicus Sharpe, 1881 28 Cat. Bds Brit. Mus., 6: 235, pl. 15. HOLOTYPE, Adult male. Reg. no. 1869.4.10.2. Sarayacu, Ucayali River, east Peru. Collected by and purchased of E. Bartlett.
amazonum—Heteropelma amazonum P. L. Sclater, 1860 8 Proc. zool. Soc. Lond., 1860: 466. HOLOTYPE, Adult (female, on label). Reg. no. 1888.1.13.1370. Chamicuros (Huallago or Uceyale River, on label), Peru. Collected by Hauxwell. Sclater Collection; purchased by him from J. Gould.
amazonum—Thlypopsis amazonum P. L. Sclater, 1886 51 Cat. Bds. Brit. Mus., 11: 229. HOLOTYPE, Adult male. Reg. no. 1884.6.12.825. Lower Ucayali, Upper Amazonia (30 May 1865, on label). Collected by E. Bartlett. Sclater Collection.
ambigua—Cisticola ambigua Sharpe, 1900
ambiguus—Propasser ambiguus Hume, 1874
amboinensis—Calornis amboinensis G. R. Gray, 1858 62 Proc. zool. Soc. Lond., 1858: 182. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1842.1.19.49. Amboyna. Possibly collected by S. Muller (vide Sharpe, in Hist. Coll. Brit. Mus., 2 (1906): 411). Purchased of J. Leadbeater from the Leiden Museum.
amboinensis—Dicrurus amboinensis G. R. Gray, 1860 64 Proc. zool. Soc. Lond., 1860: 354. HOLOTYPE, Adult female. Reg. no. 1873.5.12.1969. Amboyna, Moluccas, 1857. Collected by A. R. Wallace. Wallace Collection.
amboynensis—Leucocirca leucophys amboynensis Mathews, 1928. 37A Bull. Brit. Orn. Cl., 48:92. HOLOTYPE, Adult male. Reg. no. 1913.6.10.118. Pokka, Amboyna, Moluccas, 12 April 1911. Collected by C. B. Kloss. Presented by the subscribers to the British Ornithologists' Union Expedition to Dutch New Guinea.
ambrynensis—Petroeca ambrynensis Sharpe, 1900
amicorum—Turdus torquatus amicorum, see appendix, Corrections and

ammodendri—Passer ammodendri Gould, 1872 The Birds of Asia, 5: pl. 15. SYNTYPE, Adult male. Reg. no. 1881.5.1.2043. Turkestan, (Djulek on the Syr-Darya, vide Hartert, Vogel palaearctische Fauna, (1909): 158), 27 Nov. 1866. Collected by N. A. Severtzoff. Gould Collection. Gould used nomen nudum of Dode, Proc. zool. Soc. Lond., 1871: 481, (see also Sharpe, Cat. Bds. Brit. Mus., 12 (1887): 337, footnote). amoenus—Turdus amoenus Horsfield, 1821. Trans. Linn. Soc. Lond., 13: 147. Syntype, Adult male. Reg. no. 1880.1.1.3001. Java, between 1811 and 1816. Presented by the Honourable East India Company Museum. There are several other syntypes in the collection. 20.288. Warar Bum, 6,000', 30 miles east of Bhamo, Shan States. Collected and presented by G. Rippon. ampotekae—Zosterops maderaspetana ampotekae Salomonsen, 1934 47 Proc. zool. Soc. Lond., 1934: 222. HOLOTYPE, Adult female. Reg. no. 1931.8.18.3413. Ampoteka, South-west Madagascar, 15 March 1930. Collected by J. Delacour or his party. From the French-Anglo-American Expedition. 20.415. Gulf of Abrek, Sea of Japan, 14 April 1876. Seebohm Collection: purchased by Seebohm of W. Schlüter. amurensis—Sitta amurensis Swinhoe, 1871. Proc. zool. Soc. Lond., 1871: 350. HOLOTYPE, Adult (female, on label. Reg. no. 1898.9.20.870. Amur-land (2 August 1855, on label). Collected by Dr. Schrenk. Seebohm Bequest. amyae—Acrocephalus stentoreus amyae E. C. S. Baker, 1905. . 35 Bull. Brit. Orn. Cl., 43:17. HOLOTYPE, Adult male. Reg. no. 1922.8.9.1. Hessamara, North Lakhimpur, Eastern Assam, 12 April 1905. Collected and presented by H. Stevens. amyae—Turdinulus epilepidotus amyae Kinnear, 1925 . Bull. Br. Orn. Cl., 45:73. HOLOTYPE, Adult male. Reg. no. 1924.12.21.185. Bao Ha, Tonkin, 5 Jan. 1924. Collected by H. Stevens and presented by the Percy Sladen Trust and Godman Exploration Fund. anabatinus—Thamnistes anabatinus P. L. Sclater and Salvin, 1860 . 4 Proc. zool. Soc. Lond., 1860: 299. SYNTYPE, Adult. Reg. no. 1889.9.20.176. Choctum, Vera Paz. Collected by O. Salvin. Sclater Collection. There are three other syntypes in the collection.

- anak—Cvornis anak Robinson and Kloss, 1922 J. Fed. Malay Sts. Mus., 10:261. HOLOTYPE, Adult male. Reg. no. 1936.4.12.2122. Krongmun, 2,000', Trang, Siam Peninsula, 16 Feb. 1910. Collected by H. C. Robinson and C. B. Kloss. Presented by the Raffles Museum, Singapore.
- 1860.4.16.226. Java. Presented by the Honourable East India Company Museum.

There are two other syntypes in the collection.

- analoga—Zosterops maderaspatana analoga Salomonsen, 1934 . 47 Novit. Zool., 39: 208. HOLOTYPE, Adult male. Reg. no. 1931.8. 18.3409. Manjakatompo, Ankarata Mountains, Madagascar, 3 May 1929. Collected by J. Delacour or party. French-Anglo-American Expedition.
- anatoliae—Garrulus atricapillus anatoliae Seebohm, 1883 Ibis, (5) 1:7. SYNTYPE, Adult female. Reg. no. 1897.6.1.328. Gosna, Taurus, on label), Asia Minor, (two labels, dated 14 Dec. 1875 and 12 Dec. 1874 respectively). Collected by C. G. Danford. Seebohm Bequest.

There are two other syntypes in the collection.

andamanensis—Eulabes andamanensis Beavan, 1867 Ibis, (2) 3:331. SYNTYPE, Adult male. Reg. no. 1888.9.20.307. Andaman Islands, June 1865. Collected by R. C. Beavan. Tweeddale Collection.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 108. Beavan used a Tytler MS name.

- andamanensis—Geocichla andamanensis Walden, 1874. 31 Ann. Mag. nat. Hist., (4) 14:156. SYNTYPE, Adult male. no. 1888.2.20.449. South Andaman Islands, 25 April 1873. Collected by R. G. Wardlaw-Ramsay. Tweeddale Collection. There is also a female syntype in the collection.
- andamanensis—Graucalus javensis andamanensis Whistler, 1940 Ibis, (14) 4:731. HOLOTYPE, Adult male. Reg. no. 1887.12.30.106. South Andaman Islands, 3 Jan. 1873. Collected by W. R. Wardlaw-Ramsay. Tweeddale Collection.
- andamanica—Arachnechthra andamanica Hume, 1873 . Stray Feathers, 1:404. SYNTYPE, Adult male. Reg. no. 1886.12.1. 1898. Aberdeen, Andaman Islands, (26 Dec. 1872, on label). Collected by W. Davison. Hume Collection. There are a number of syntypes in the collection.
- andapae—Oxylabes zosterops andapae Salomonsen, 1934. . 21 Ann. Mag. nat. Hist., (10) 14:69. HOLOTYPE, Adult male. Reg. no. 1931.8.18.2043. One day west of Andapa, north-east Madagas-

car, 22 Aug. 1930.	Collected by A. L.	. Rand and	d Du Mon	t on the
Archbold-Vernay E	xpedition to Madag	gascar.		

- andarvae—Chlorophoneus andarvae Jackson, 1919 23 Bull. Br. Orn. Cl., 39:94. HOLOTYPE, Adult male. Reg, no. 1919.7.5.1. Kisubi, 4,000', Entebbe, Uganda, 8 Oct. 1916. Collected by Andarya for Sir F. J. Jackson and presented by the latter.
- anderseni—Cinnyris regius anderseni Williams, 1950 46 Ibis, 92:644. HOLOTYPE, Adult male. Reg. no. 1950.65.2. Mahari Mountains, eastern shore of Lake Tanganyika, 16 Oct. 1949. Collected for T. Andersen. Presented by the Coryndon Museum, Nairobi.
- andersoni—Aethopyga andersoni Oates, 1890 Fauna of British India, Birds, 2:349. SYNTYPE, Adult male. Reg. no. 1876.4.7.38. Sawaddy, East of Bhamo, Yunnan, Burma, 28 Jan. 1875. Collected by J. Anderson. Purchased from the Yunnan Expedition.

The two other syntypes are also in the collection.

andersoni—Ixus andersoni Swinhoe, 1870 21 Ann. Mag. nat. Hist., (4) 5:175. SYNTYPE, Adult male. Reg. no. 1898.10.2.123. Ichang, 1,000 miles up the Yangtse, (Kweichoo on label), 18 April 1869. Collected by R. Swinhoe. Seebohm Collection.

There are two female syntypes in the collection.

- anderssoni—Campephaga anderssoni Sharpe, 1870 20 Proc. zool. Soc. Lond., 1870: 70, pl. 4. HOLOTYPE, (Adult female, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 30). Reg. no. 1875. 4.9.318. Ovaquenyama, Damaraland, South-West Africa, 25 June 1867. Collected by C. J. Andersson. Sharpe Collection.
- anderssoni-Cotile anderssoni Sharpe and Wyatt, 1887. A Monograph of the Hirundinidae, 1:119. HOLOTYPE, Adult (female on label). Reg. no. 1874.5.1.277. Damaraland (in text: Daviep, 27 Oct. 1860, on label). Collected by C. J. Andersson and purchased of R. B. Sharpe.
- anderssoni—Saxicola anderssoni Sharpe, 1877 in Layard, Birds of South Africa: 249. SYNTYPE, Adult male. Reg. no. 1876.5.23.426. "The Reeds", Swakop River, Damaraland, 8 Dec. 1860. Collected by C. J. Andersson. Sharpe Collection. There are several other syntypes in the collection.
- anderssoni—Zosterops anderssoni Shelley, 1892 47 Bull. Brit. Orn. Cl., 1:5. SYNTYPE, Adult male. Reg. no. 1876.5.23.167. Elephant Vley, Damaraland, (14 Nov. 1859, on label). Collected by C. J. Andersson. Sharpe Collection. The author refers to two specimens (vide Shelley, Birds of Africa,

2 (1900): 177) and both will be syntypes.

Agriornis andicola P. L. Sclater, 1860
Collected by L. Fraser. Sclater Collection. Renamed A. pollens P. L. Sclater, 1869 on the grounds of its preoccupation by A. andecola d'Orbigny, (1839).
andicola—Leptasthenura andicola P. L. Sclater, 1869
andrewi—Hirundo andrewi J. G. Williams, 1966
anglica—Loxia curvirostra anglica Hartert, 1904
anglorum—Regulus regulus anglorum Hartert, 1905 35 Bull. Brit. Orn. Cl., 16:11. Holotype, Adult male. Reg. no. 1936.10.15.15. Tring, Hertfordshire, England, 31 Oct. 1900. Collected by E. Hartert. Presented by the American Museum of Natural History, New York; from the Rothschild Collection.
angolensis—Cercomela familiaris angolensis Lynes, 1926 31 Ibis, (12) 2:393. HOLOTYPE, Adult male. Reg. no. 1906.12.4.119. Huxe, Benguella, Angola, 10 Oct. 1905. Collected by and purchased of Dr. W. J. Ansorge.
angolensis—Dryoscopus angolensis Hartlaub, 1860 23 Proc. zool. Soc. Lond., 1860: 111. HOLOTYPE, Adult. Reg. no. 1873.12.10.10. Bembe, Loanda Province, Angola. 5 May 1858. Collected by and purchased of J. J. Monteiro.
angolensis—Eremomela scotops angolensis Bannerman, 1937 . 35 Bull. Brit. Orn. Cl., 57:111. HOLOTYPE, Adult male. Reg. no. 1910.5.6.848. Lalange, North Angola, 12 Feb. 1909. Collected by and purchased of W. J. Ansorge.
angolensis—Euprinodes rufigularis angolensis Bannerman, 1922 . 35 Bull. Brit. Orn. Cl., 43: 30. HOLOTYPE, Adult female. Reg. no. 1910.5.6.918. Ndala Tando, Northern Angola, 29 Oct. 1908. Collected and purchased of W. J. Ansorge.

- angolensis—Macrospherus flavicans angolensis Bannerman, 1920 . 35

 Bull. Brit. Orn. Cl., 41:6. HOLOTYPE, Adult male. Reg. no.
 1910.5.6.670. N'Dala Tando, northern Nigeria, 26 Oct. 1909.
 Collected by W. J. Ansorge. Ansorge Collection.

 angolensis—Nigrita canicapilla angolensis Bannerman, 1921 . . 60

 Bull. Brit. Orn. Cl., 41:126. HOLOTYPE, Adult female. Reg. no.
- angolensis—Nigrita canicapilla angolensis Bannerman, 1921 . . . 60 Bull. Brit. Orn. Cl., 41:126. HOLOTYPE, Adult female. Reg. no. 1910.5.6.1432. Ndala Tando, northern Angola, 14 Sept. 1908. Collected by and purchased of W. J. Ansorge.

- ankafanae—Oxylabes zosterops ankafanae Salomonsen, 1934. 21
 Ann. Mag. nat. Hist., (10) 14:70. Holotype, Adult male. Reg.
 no. 1882.2.27.59. Fianarantsoa, Ankafana Forest, south-east
 Madagascar, Feb. 1881. Collected by and purchased of the Rev.
 W. D. Cowan.
- ankaratrae—Saxicola torquata ankaratrae Salomonsen, 1934 . 31 Novit. Zool., 39: 210. HOLOTYPE, Adult male. Reg. no. 1931. 8.18.2556. Manjakatompo, Ankaratra Mountains, central Madascar, 14 May 1929. Collected by J. Delacour on the Archbold-Vernay Expedition.
- ankole—Cisticola brachyptera ankole Lynes, 1930 . . . 35

 Ibis, (12) 6, supplement: 489. HOLOTYPE, Adult male. Reg. no. 1928.10.25.10. Ankole District, Uganda, 17 Feb. 1917. Collected by Sir F. Jackson. Presented by Admiral H. Lynes.
- annae—Cisticola cisticola annae R. Meinertzhagen, 1922 . . . 35

 Ibis, (11) 4:12. HOLOTYPE, Adult. Reg. no. 1903.8.2.99. Kuklia

 Marsh, Famagusta, Cyprus, 27 Oct. 1901. Collected and presented
 by Miss D. Bates.

- annamensis—Aegithaliscus annamensis Robinson and Kloss, 1919 Ibis. (11) 1:606. SYNTYPE, Adult male. Reg. no. 1919.12.20.425. Langbian Peaks, 6,500', southern Annam, 17 April 1918. Collected by C. B. Kloss, and presented by H. C. Robinson and C. B. Kloss. This is the authors' selected male, their selected female syntype is also in the collection annamensis—Aethopyga gouldiae annamensis Robinson and Kloss. Ibis. (11) 1:621, pl. 18. SYNTYPE, Adult male. Reg. no. 1919. 12.20.508. Langbian Peaks, 6,500', South Annam, 19 April 1918. Collected and presented by C. B. Kloss. The described female syntype is also in the collection. annamensis—Alcippe nepalensis annamensis Robinson and Kloss, 1919 Ibis. (11) 1:582. SYNTYPE, Adult female. Reg. no. 1919.12.20.296. Dalat, 5,000', south Annam, 5 April 1918. Collected and presented by C. B. Kloss. This is the authors' selected male. Their selected female syntype is also in the collection. annamensis—Corythocichla annamensis Delacour and Jabouille, 1928 32B Bull. Br. Orn. Cl., 48:131. HOLOTYPE, Adult male. Reg. no. 1928.6.26.7. Phuqui, 150 metres, north Annam, 23 Feb. 1928. Collected by J. Delacour and P. Jabouille on the Fourth Franco-British Expedition to Indochina. annamensis—Cryptolopha castaneiceps annamensis Robinson and Ibis, (11) 1:447. SYNTYPE, Adult male. Reg. no. 1919.12.20.181. Langbian Peaks, 5,500-5,600', South Annam, 27 April 1818. Collected and presented by C. B. Kloss. Kloss, 1919 35 This is the described male. The described female syntype is also in the collection. annamensis—Dendrobiastes hyperythra annamensis Robinson and Kloss, 1919 Langbian Peaks, 6,000-7,000', South Annam, 20 April 1918. Collected and presented by H. C. Robinson and B. B. Kloss. This is the described male, the described female syntype is also in the collection.
 - annamensis—Drymocataphus tickelli annamensis Delacour, 1926. 32B Bull. Br. Orn. Cl., 47:17. Syntype, Adult male. Reg. no. 1926. 9.8.19. Col des Nuages, 1,250', Annam, 29 Nov. 1925. Collected by J. Delacour on the Second Franco-British Expedition to Indochina.

The collector's number is 910, not 906 as printed. This is the author's selected male. His selected female syntype is also in the collection.

- annamensis—Leptocoma flammaxillaris annamensis Hachisuka, 1926 46 Bull. Brit. Orn. Cl., 47:68. HOLOTYPE, Adult male. Reg. no. 1906.12.5.69. Nhatrang, Annam, 2 Oct. 1905. Collected by and purchased of J. J. Vassal.
- annamensis-Pericrocotus brevirostris annamensis Robinson and Kloss, 1923 Bull. Br. Orn. Cl., 43:119. HOLOTYPE, Adult female. Reg. no. 1959.3.1. Dalat, 5,000', Langbian Plateau, south Annam, 8 April 1918. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore.
- annamensis—Pnoepyga pusilla annamensis Robinson and Kloss, 1919 32B Ibis, (11) 1:591. SYNTYPE, Adult female. Reg. no. 1919.12.20. 342. Langbian Peaks, 6,000-7,500', south Annam, 28 April 1918. Collected and presented by C. B. Kloss.

 This is the authors' selected female. Their selected male syntype

is also in the collection.

- annamensis—Poliopsar leucocephalus annamensis Wells, 1919. . . . 62 Bull. Br. Orn. Cl., 39:77. HOLOTYPE, Adult male. Reg. no. 1910.5.8.120. Nhatrang, Annam, Cochin China, Jan. 1908. Collected and purchased of J. J. Vassall.
- annamensis—Pomatorhinus olivaceus annamensis Robinson and Kloss 1919 276. Dran, 3,000', south Annam, 13 May 1918. Collected and presented by C. B. Kloss.
- annamensis-Pterythius aeralatus annamensis Robinson and Kloss, Langbian Peaks, 6,000-7,000', south Annam, 19 April 1918. Collected and presented by C. B. Kloss.

This is the author's selected male. Their selected female is also

in the collection.

annamensis-Stactocichla merulina annamensis Robinson and Kloss, *Ibis*, (11) 1:577, pl. 13. SYNTYPE, Adult male. Reg. no. 1919.12. 20.529. Dran, 3,000', south Annam, 12 May 1918. Collected and presented by C. B. Kloss.

This is the authors' selected male. Their selected female syntype

is also in the collection.

Reg. no. 1843.1.13.326. Nepal. Collected and presented by B. H. Hodgson.

30	annectens
	Received with his main collection, the specimen has Hodgson's drawing number on the label and is therefore likely to have been before him when the description was made.
r	Rectens—Ixus annectens Walden, 1875
<i>J</i>	ectens—Tephrodornis pelvica annectens Robinson and Kloss, 1918 20 E.F.M.S. Mus., 8: 222. HOLOTYPE, Adult male. Reg. no. 1936. 12.140. Lamra, Trang, Federated Malay States, 12 Jan. 1910. Presented by the Raffles Museum, Singapore.
I I	mala—Callene anomala Shelley, 1893
J	nyma—Drymoeca anonyma von Müller, 1855
/ c k	phrys—Aegithaliscus anophrys Swinhoe, 1868 40 bis, (2) 4:64. SYNTYPE, Adult. Reg. no. 1888.5.30.164. Tinghow Mountains, (120 miles north-east of Amoy, on label), Fohcien. (Aug. 1867, on label). Collected for R. Swinhoe. Tweeddale Request. There are three other syntypes in the collection.

anoxantha—Spermophila anoxantha Gosse, 1847. Birds of Jamaica: 247, pl. 62. Syntype, Adult male (relaxed mount). Reg. no. 1847.6.8.20. Mount Edcecumbe, Jamaica. Collected by and purchased of P. H. Gosse. Two other syntypes, a female and a male, are also in the collection.

ansorgei—Anthus leucophrys ansorgei White, 1948 Ibis, 90: 549. HOLOTYPE, Adult male. Reg. no. 1910.5.6.1250. Bissao, Portuguese Guinea, 18 Aug. 1909. Collected by and purchased of Dr. W. J. Ansorge.

Ndalla Tando, Northern Angola, 15 Sept. 1908. Collected by W. J. Ansorge. Ansorge Collection.

ansorgei—Dryoscopus bocagei ansorgei W. L. Sclater, 1911 Bull. Br. Orn. Cl., 29:37. HOLOTYPE, Adult male. Reg. no. 1910.5.6.950. N'Dalla Tando, north Angola, 5 Aug. 1908. Collected by and purchased of Dr. W. J. Ansorge.

- ansorgei—Mirafra sabota ansorgei W. L. Sclater, 1926. Bull. Br. Orn. Cl., 47:29. HOLOTYPE, Adult female (in text; male on label). Reg. no. 1905.11.22.173. Catumbella, Benguella, Angola, 6 Aug. 1905. Collected by and purchased of W. J. Ansorge.
- ansorgei—Ortygospiza ansorgei Ogilvie-Grant, 1910 Bull. Brit. Orn. Cl., 25:84. SYNTYPE, Adult male. Reg. no. 1910.5.6.1337. Gunnel, Portuguese Guinea, 30 July 1909. Collected by and purchased of W. J. Ansorge. The two other syntypes are also in the collection.
- ansorgei—Prinia ansorgei W. L. Sclater, 1927 Bull. Brit. Orn. Cl., 48:18. HOLOTYPE, Adult male. Reg. no. 1906.12.4.149. Huxe, Benguella, Angola, 5 Sept. 1905. Collected and purchased by W. L. Ansorge.
- ansorgii—Bradvornis infuscatus ansorgii Ogilvie-Grant, 1913 Ibis, (10) 1:636. SYNTYPE, Adult male. Reg. no. 1906.12.4.9. Huxe, Benguella, Angola, (6 Sept. 1905, on label). Collected by and purchased of W. J. Ansorge.

This is the described male, the described female syntype is also in the collection

- ansorgii—Erythropygia ansorgii Ogilvie-Grant, 1914 Bull. Br. Orn. Cl., 33:134. SYNTYPE, Adult male. Reg. no. 1910.5.6.712. Malange, north Angola, 15 Feb. 1909. Collected by and purchased of Dr. W. J. Ansorge. The female syntype is also in the collection.
- antarcticus—Anthus antarcticus Cabanis, 1884 J. Orn., 1884: 254. SYNTYPE, Adult male. Reg. no. 1884.5.15.133. South Georgia, 1885. Collected by Dr. von Steinen. Sclater Collection; obtained by Sclater in exchange with the Hamburg Museum.
- antisiensis—Synallaxis antisiensis P. L. Sclater, 1858. Proc. zool. Soc. Lond., 1858: 457. HOLOTYPE, Adult (female, vide Sclater, Catalogue of American Birds, 1862: 158). Reg. no. 1889.5. 20.219. Cuenca, Ecuador, Nov. 1857. Collected by L. Fraser. Sclater Collection.
- antonii—Mirafra angolensis antonii Hall, 1958 17 Bull. Br. Orn. Cl., 78: 153. HOLOTYPE, Adult male. Reg. no. 1957.35.2. 5 miles east of Luacano, 3,500', east Angola, 31 Aug. 1957. Collected by A. L. Archer and presented by the B. P. Hall Angola Expedition.
- apicalis—Myiarchus apicalis P. L. Sclater and Salvin, 1881 . Ibis, (4) 5:269. SYNTYPE, Adult. Reg. no. 1888.1.13.1114. Bogota, Colombia. Collected for Boucard. Sclater Collection. There are three other syntypes in the collection.

annroximans—Cercomacra annroximans Pelzeln 1869 4
approximans—Cercomacra approximans Pelzeln, 1869 4 Zur Ornithologie Brasiliens: 85, 158. Syntype, Adult male. Reg. no. 1889.9.20.421. Matto Grosso, Brazil, 23 Oct. 1826. Collected by J. Natterer. P. L. Sclater Collection. There is a female syntype in the collection. Both are listed as "authentic specimens" by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 266.
approximans—Pachycephala poliosoma approximans Ogilvie-Grant, 1911
approximans—Xenops approximans Pelzeln, 1859 3 Sher. Akad. Wiss. Wien, 34:113. Syntype, Adult. Reg. no. 1889.5.20.405. Rio Negro, Brazil, 30 Sept. 1830. Collected by J. Natterer. P. L. Sclater Collection. Listed as "authentic specimen" by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 111.
aquatica—Sayornis aquatica P. L. Sclater and Salvin, 1859 9 Ibis, (1) 1:119. HOLOTYPE, Adult. Reg. no. 1888.1.1.126. Dueñas, Guatamala, 4 Feb. 1858. Collected by O. Salvin. Salvin- Godman Collection. This is the only known specimen collected at the time that the type description was made. Sclater's list of three syntypes (vide, Cat. Bds. Brit. Mus., 14 (1888): 34) appears to be an error.
arabicus—Cinnyricinclus leucogaster arabicus C. H. B. Grant and Mackworth-Praed, 1942
arabicus—Lanius arabicus Ogilvie-Grant, 1905 23 Bull. Br. Orn. Cl., 15:78. SYNTYPE, Adult male. Reg. no. 1902.12.12.24. Gerba, Amiri District, South Arabia. Collected by and purchased of G. W. Bury. There are several other syntypes in the collection.
arcaei—Buthraupis arcaei P. L. Sclater and Salvin, 1869 51 Proc. zool. Soc. Lond., 1869: 439, pl. 31. Syntype, Adult male. Reg. no. 1885.6.7.54. Cordillera del Chucu, Veragua, 1869. Collected by E. Arcé. Salvin-Godman Collection. The female syntype is also in the collection.
archeri—Cossypha archeri Sharpe, 1902

archeri—Eremomela griseoflava archeri W. L. Sclater, 1927 35 Bull. Brit. Orn. Cl., 48:14. HOLOTYPE, Adult male. Reg. no. 1908.12.22.54. Burao, Somaliland, 9 Jan. 1906. Collected by G. W. Bury. Received in exchange with the Transvaal Museum.
archeri—Heteromirafra archeri Clarke, 1920
archetes—Cecropsis archetes Hume, 1877
arcticincta—Hirundo arcticincta Sharpe, 1891
arctitorquis—Pachycephala arctitorquis P. L. Sclater, 1883 39 Proc. zool. Soc. Lond., 1883:55, pl. 13. SYNTYPE, Adult female. Reg. no. 1883.5.30.62. Larat Island, Timor Laut, Tenimber Group, 3 August 1882. Collected by H. O. Forbes. Presented by the "Timor Laut" Committee of the British Association. The male syntype cannot be traced.
arcuata—Emberiza arcuata Sharpe, 1888
ardens—Psaropholus ardens Swinhoe, 1862
ardesiacus—Cinclus ardesiacus Salvin, 1867
ardesiacus—Dysithamnus ardesiacus P. L. Sclater, 1867 4 Proc. zool. Soc. Lond., 1867: 756. SYNTYPE, Adult male. Reg. no. 1889.9.20.214. Rio Napo, eastern Ecuador. Collected for Verreaux. Sclater Collection. There are two other syntypes in the collection.

- aremoricus—Aegithalos caudatus aremoricus Whistler, 1929 . . . 40 Bull. Brit. Orn. Cl., 49:87. HOLOTYPE, Adule female. Reg. no. 1929.3.9.1. Loudiac, Brittany, France, 21 Oct. 1928. Collected and presented by H. Whistler.
- arenacea—Ochthoeca arenacea P. L. Sclater and Salvin, 1877. 9

 Proc. zool. Soc. Lond., 1877: 20. Holotype, Adult. Reg. no. 1888.1.1.1321. Bogota, Colombia. (Collected by G. Crowther, on label.) Salvin-Godman Collection.
- arequipae—Synallaxis arequipae P. L. Sclater and Salvin, 1869 . 3

 Proc. zool. Soc. Lond., 1869: 417. SYNTYPE, Adult male. Reg.
 no. 1889.5.20.242. Arequipa, south-west Peru, 1867. Collected
 by H. Whitely. Sclater Collection.
 There are five other syntypes in the collection.
- argentauris—Mesia argentauris Hodgson, 1838 32B India Review, 2:88. Syntype, Adult (relaxed mount). Reg. no. 1843.1.13.437. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
- argentea—Mixornis gularis argentea Chasen and Kloss, 1930 . 32B Bull. Raffles Mus., 4:82. HOLOTYPE, Adult male. Reg. no. 1947.60.58. Mallewallé Island, North Borneo, 9 Aug. 1927. Collected by F. N. Chasen and C. B. Kloss and presented by the Raffles Museum, Singapore.

In the original description reference was made only to this subadult specimen, but an adult was described. Subsequently Gould, *Birds of Australia*, 2 (1848): 51, thanked Bynoe "for one of the specimens from which my description was taken." The latter, an adult, must also be a syntype and would account for the description, and later illustration, of an adult. It is in the Academy of Natural Sciences

of Philadelphia and De Schaunsee, <i>Proc. Acad. nat. Sci. Philadelphia</i> , 109 (1957): 223 acknowledges it as a cotype in discussion but calls it a holotype in the major reference.
argentina—Guiraca argentina Sharpe, 1888
of the Americas, 11 (1938): 102. There are several other syntypes in the collection.
argyrofenges—Calliste argyrofenges P. L. Sclater and Salvin, 1876. 51 Proc. zool. Soc. Lond., 1876: 354, pl. 30. HOLOTYPE, Adult male. Reg. no. 1888.6.4.628. Tilotilo, Yungas Province, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.
aridula—Cisticola aridula Witherby, 1900
Bull. Brit. Orn. Cl., 11:13. HOLOTYPE, Adult male (immature, vide Lynes, Ibis, (12) 6, Supplement, (1930):130). Reg. no. 1901. 12.12.1. Gerazi, 60 miles south of Khartoum, White Nile, (1 May 1900, on label). Collected and presented by H. F. Witherby.
arizanica—Pyrrhula arizanica Ogilvie-Grant, 1912
armillatus—Cyanocorax armillatus G. R. Gray and Mitchell, 1845. 71 The Genera of Birds, 2:307, pl. 74. HOLOTYPE, Adult. Reg. no. 1845.1.2.3. ("Bogota", Colombia, vide Sharpe, Cat. Bds. Brit. Mus., 3 (1877):137). "Purchased of Warwick", in register; "purchased of Argent" on label.
chased of Argent", on label.
arquata—Pitta (Phoenicocichla) arquata Gould, 1871 12 Ann. Mag. nat. Hist., (4) 7: 340. HOLOTYPE, Adult, probably female. Reg. no. 1881.5.1.4603. Borneo. Gould Collection.
arthuri—Planesticus arthuri Chubb, 1914
arturi—Nectarinia arturi P. L. Sclater, 1906
Bull. Brit. Orn. Cl., 19:30. SYNTYPE, Adult male. Reg. no. 1907.6.25.2. Wolverhampton, South Melsetter, Southern Rhodesia, 13 Feb. 1906 (13 April, on label). Collected by W. M. Stanley for

A. L. Sclater. Purchased of A. L. Sclater.

The second syntype with head much damaged is also in the collection.

aruensis—Myiolestes aruensis G. R. Gray, 1858
aruensis—Myzomela obscura aruensis Kinnear, 1924
aruensis—Ptilotis aruensis Sharpe, 1884
aruensis—Rectes aruensis Sharpe, 1877
ashantiensis—Cinnyris superbus ashantiensis Bannerman, 1922 . 46 Bull. Brit. Orn. Cl., 43:7. Holotype, Adult male. Reg. no. 1911.4.26.2. Ashanti, Gold Coast. Collected and presented by H. G. F. Spurrell.
asirensis—Pica pica asirensis Bates, 1936
asmaraensis—Calandrella cinerea asmaraensis K. G. Smith, 1951. 17. Bull. Br. Orn, Cl., 71:55. HOLOTYPE, Adult Male. Reg. no. 1951. 48.2. Near Asmara, Eritrea, 12 Oct. 1951. Collected and presented by K. D. Smith.
aspasioides—Nectarinia aspasioides G. R. Gray, 1860
assamensis—Cinnyris assamensis McClelland, 1840

assamensis—Drymocataphus assamensis Sharpe, 1883 32B Cat. Bds. Brit. Mus., 7:557. Syntype, Adult male. Reg. no. 1895.7.14.3398. Dikrang Valley, Assam, Dec. 1874. Collected and presented by H. H. Godwin-Austen. The second syntype is also in the collection.
assamensis—Pomatorhinus assamensis Blyth, 1847 32B J. Asiat. Soc. Bengal, 16: 451. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4776. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum. Blyth used McClelland's MS name.
assamensis—Psaroglossa spiloptera assamensis E. C. S. Baker, 1924 62 Bull. Br. Orn. Cl., 45:14. HOLOTYPE, Adult male. Reg. no. 1888.9.20.469. North Khasia Hills, Assam, Dec. 1876. Collected by A. W. Chennell, Tweeddale Collection.
assamica—Mirafru assamica, McClelland, 1840 17 in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840): 162. Holotype, Adult (relaxed mount). Reg. no. 1880.1.1.4732. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.
assamica—Suya crinigera assamica E. C. S. Baker, 1924 Bull. Brit. Orn. Cl., 44:80. HOLOTYPE, Adult male. Reg. no. 1895.7.14.1560. Shillong, Assam, (22 May, on label). Collected by H. H. Godwin-Austen. Godwin-Austen Collection.
assimilis—Cactornis assimilis Gould, 1837
assimilis—Dendrocitta assimilis Hume, 1877
assimilis—Dicrurus assimilis G. R. Gray, 1858
assimilis—Eremiornis carteri assimilis Montague, 1913 35 Austral Avian Record, 1:181. SYNTYPE, Adult. Reg. no. 1913.5. 20.24. Hermite Island, Monte Bello Group, North-west Australia,

Monte Bello Expedition. Presented by the Royal Society. There are two other syntypes in the collection.
assimilis—Horornis assimilis J. E. Gray, 1863 35 Catalogue of mammalia and birds of Nepal and Thibet presented by B. H. Hodgson to the British Museum, 2nd Edn.: 30. ?SYNTYPE, Adult (relaxed mount). Reg. no. 1848.6.4.58. Nepal. Collected and presented by B. H. Hodgson. This specimen was relaxed from mounting in 1882 as the type, and listed as such by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 137. There do not appear to be other available specimens but it cannot be proved that in this case the author described a mounted specimen in the public gallery of the museum.
assimilis—Mionectes assimilis P. L. Sclater, 1859
assimilis—Oriolus assimilis Tweeddale, 1877 60 Proc. zool. Soc. Lond., 1877: 760, pl. 76. Holotype, Adult male. Reg. no. 1887.11.20.525. Cebu Island, Philippines, Mar. 1877. Collected by A. H. Everett. Tweeddale Collection.
assimilis—Passer assimilis Walden, 1870
assimilis—Psarisomus assimilis Hume, 1873. Stray Feathers, 3:53. SYNTYPE, Adult male. Reg. no. 1887.5.1. 507. West Pegu Hills, Burma, 12 Apr. 1873. Collected by E. W. Oates. Hume Collection. The female syntype is in the collection.
assimilis—Quiscalus assimilis P. L. Sclater, 1862
assimilis—Rhipidura assimilis G. R. Gray, 1858 37A Proc. zool. Soc. Lond., 1858 : 176. HOLOTYPE, Adult male. Reg. no. 1858.3.10.79. Ké Island, South-west Pacific. Collected by A. R. Wallace. Wallace Collection.
assimilis—Strachyris assimilis Walden, 1875

1148. Karen-Nee, 2,800', Burma, 14 Feb. 1874. Collected by R. G. Wardlaw-Ramsay. Tweeddale Collection.
Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 602. The male syntype is also in the collection.
Istigma—Hirundo astigma Shelley, 1893
Bull. Br. Orn. Cl., 41:115. SYNTYPE, Adult male (described from life). Reg. no. 1921.9.2.1. China. Presented by the Zoological Society of London, in whose gardens it died on 28 Oct. 1921. The female syntype is also in the collection.
Interrima—Ceblepyris aterrima Blyth, 1866
Athensis—Spizocorys athensis Sharpe, 1900
Itlantica—Coereba atlantica P. R. Lowe, 1912
Atlantis—Sylvia atricapilla atlantis Williamson, 1964 35 British Trust for Ornithology Field Guide no. 9. Identification for Ringers, 3:14. HOLOTYPE, Adult male. Reg. no. 1904.12.31.134. Ponte Delgada, San Migel, Azores, 19 March 1903. Collected by W. R. Ogilvie-Grant. Presented by the Hon. W. Rothschild.
Bull. Br. Orn. Cl., 40: 32. SYNTYPE, Adult male. Reg. no. 1919. 12.11.206. (Azrou, 4,800', Lesser Atlas, central Morocco, 19 May 1919, on label.) Collected and presented by H. Lynes. There are four other syntypes in the collection.
ntlas—Otocorys atlas Whitaker, 1898
Bull. Br. Orn. Cl., 7:47. HOLOTYPE, Adult male. Reg. no. 1939. 12.14.10. Glani (lapsus for Glaoui, vide Whitaker, Ibis, (7) 4 (1898): 604), Atlas Mts., Morocco. Collected and presented by E. Dodson. The author also had a female but only referred to the male in the description.
Atlas—Parus atlas Meade-Waldo, 1901

This is the author's selected male, his selected female syntype is also in the collection.

The female syntype is also in the collection.

atricapilla—Meliphaga atricapilla Jardine and Selby, 1833 . . . 48
Illustrations of Ornithology: pl. 134. and text. HOLOTYPE, Adult.
Reg. no. 1886.6.24.346. Tasmania. (Collected by P. Forster, on label.) Jardine Collection; purchased of Gerrard.

The original label is inscribed "The authority for the fig.". The name is apparently preoccupied by both *Meliphaga atricapilla* Temminck and Languer, 1825, and *Certhia atricapilla* Latham, 1801; Gould, *Proc. zool. Soc. Lond.*, 1845: 62 provided a new name, *Melithreptus melanocephalus*, believing it to be a new species description, but *Eidopserus affinis* Lesson, 1839 is now accepted as the earliest valid name.

Listed as "authentic specimen" by Sclater, Cat. Bds. Brit. Mus. 15 (1890): 247.

The second syntype is also in the collection.

atriceps—Pseudominla atriceps Robinson and Kloss, 1919 Ibis, (11) 1:583, pl. 14. SYNTYPE, Adult male. Reg. no. 1919.12. 20.300. Langbian Peaks, 5,500–7,500', south Annam, 16 April 1918. Collected and presented by C. B. Kloss. This is the authors' selected male. Their selected female syntype

is also in the collection. Renamed Alcippe castaneicens Klossi Delacour and Jabouille, 1931 on the grounds of its preoccupation by A. atriceps Jerdon, 1839.

- atriceps—Turdinus atriceps Sharpe, 1902 . 32B Bull. Br. Orn. Cl., 13:10. HOLOTYPE, Adult male. Reg. no. 1902.12.8.16. Ruwenzori Mountains, Uganda, 14 Feb. 1902. Collected and presented by F. J. Jackson.
- atriceps—Zosterops atriceps G. R. Gray, 1860 Proc. zool. Soc. Lond., 1860: 350. SYNTYPE, Adult (male, vide Gadow, Cat. Bds. Brit. Mus., 9 (1884): 201). Reg. no. 1873.5.12. 897. Batchian, Molucca Group, 1858. Collected by A. R. Wallace, Wallace Collection.

Listed as the type by Gadow, op. cit., p. 201. There is another possible syntype in the collection.

- atrifrons—Tyrannus atrifrons P. L. Sclater, 1857. Proc. zool. Soc. Lond., 1857: 274. HOLOTYPE, Adult. Reg. no. 1888.1.13.781. Guayaquil, Ecuador. Collected for T. C. Eyton. Sclater Collection: Sclater obtained the specimen from Eyton.
- atrifrons—Zosterops atrifrons Wallace, 1864 Proc. zool. Soc. Lond., 1863 (publ. 1864): 493. HOLOTYPE, Adult (male, vide Museum catalogue). Reg. no. 1873.5.12.113. Menado. North Celebes, 1859. Collected by A. R. Wallace. Wallace Collection.
- atrigula—Ploceus atrigula Sharpe, 1890 . . Cat. Bds. Brit. Mus., 13:491. SYNTYPE, Adult male. Reg. no. 1843.1.13.881. Nepal Terai. Collected and presented by B. H. Hodgson.

This is the only syntype now available. Sharpe used a Hodgson nomen nudum from J. E. Gray, Zoological Miscellany (1844): 84. Renamed Ploceus passerinus Reichenow, 1886 on the grounds of

alleged preoccupation.

atrinucha—Thamnophilus atrinucha Salvin and Godman, 1892 Biol. Centr.-Amer., 2:200. SYNTYPE, Adult male. Reg. no. 1889.7.10.107. Lion Hill, Panama (vide Hellmayr. Proc. zool. Soc. Lond., 1911: 1158). Collected by M'Leannan. Salvin-Godman Collection.

There is a female syntype in the collection.

no. 1888.1.13.122. Babahoyo, north-east of Guayaquil, Ecuador, Aug. 1859. Collected by L. Fraser. Sclater Collection. There are two other syntypes in the collection.

atripennis—Rhipidura atripennis G. R. Gray, 1858 37A Proc. zool. Soc. Lond., 1858: 175. SYNTYPE, Adult female. Reg. no. 1858.3.10.50. Aru Islands, 1857. Collected by A. R. Wallace. Wallace Collection.

The author stated that a syntype of each sex was in the collection in 1858, but only the female one remains. Since a male specimen (register no. 55) was sent in exchange to the Calcutta Museum at an unknown date this was probably the male syntype.

- atrocaeruleus—Dicrurus atrocaeruleus G. R. Gray, 1860 . . . 64

 Proc. zool. Soc. Lond., 1860 : 354. SYNTYPE, Adult male. Reg.
 no. 1873.5.12.1967. Batchian, Moluccas, 1858. Collected by
 A. R. Wallace. Wallace Collection.

There is an obvious printer's error in the length of wing in the description. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 3 (1877): 240. There is another syntype in the collection.

atrochalybeia—Tchitrea atrochalybeia Thomson, 1842 38

Ann. Mag. nat. Hist., (1) 10: 204. Holotype, Adult male. Reg.
no. 1842.9.24.8. Fernando Po, ("more probably Sao Thomé Island", P. L. Sclater, Systema Avium Aethiopicarum (1930): 436).
Collected and presented by T. R. H. Thomson (an old label gives Fraser as the collector), from the River Niger Expedition.

A note in the British Museum Register states "described by G. R. Gray for Thomson".

This specimen, together with three other syntypes in the collection were listed as "types" by Sharpe, Cat. Bds. Brit. Mus., 12 (1888):

367.

atrogularis—Suya atrogularis Moore, 1854	5
pany Museum as the type. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 181. There are three other syntypes in the collection.	
atrogularis—Thryothorus atrogularis Salvin, 1864	3
atronitens—Aplonis atronitens G. R. Gray, 1859 62 Proc. zool. Soc. Lond., 1859: 164. HOLOTYPE, Immature (vide E. L. and L. C. Layard, Ibis, (4) 4 (1880): 229). Reg. no. 1854.5.31.8. Loyalty Islands. Presented by Sir G. Grey.	2
atrosuperciliaris—Chleuasicus ruficeps var. atrosuperciliaris Godwin-Austen, 1877	3
atrovirens—Campephaga (Lalage) atrovirens G. R. Gray, 1861 . 20 Proc. zool. Soc. Lond., 1861: 430. SYNTYPE, Adult male. Reg. no. 1861.12.11.48. Mysol. Collected by A. R. Wallace and purchased of S. Stevens. There are two other syntypes in the collection.)
aucheri—Lanius aucheri Bonaparte, 1853	3
aucklandicus—Anthus aucklandicus G. R. Gray, 1862 19 Ibis, (1) 4:224. SYNTYPE, Adult. Reg. no. 1842.12.16.35. Auckland Islands. Collected on the voyage of H.M.S. Erebus and Terror and presented by the Lords of the Admiralty. There are two other syntypes in the collection.)
aurantiaca—Chiromachaeris aurantiaca Salvin, 1870 8 Proc. zool. Soc. Lond., 1870 : 200. SYNTYPE, Adult male. Reg. no. 1888.1.20.294. Bugaba, Chiriqui, Veragua, 1869. Collected by E. Arcé. Salvin-Godman Collection. There are several other syntypes in the collection.	}
aurantiaca—Pyrrhula aurantiaca Gould, 1857)

aurantiacus—Metopothrix aurantiacus P. L. Sclater and Salvin, 1866 Proc. zool. Soc. Lond., 1866: 190. SYNTYPE, Adult female. Reg. no. 1888.1.13.1243. Sarayacu, lower Ucayali River, Peru, 30 May 1865. Collected by E. Bartlett. Sclater Collection. There is another syntype in the collection.
aurantiacus—Pheucticus aurantiacus Salvin and Godman, 1891 . 50 Ibis, (6) 3:272. Syntype, Adult male. Reg. no. 1899.2.1.33. Volcan de Santa Maria, Quezaltenango, Guatemala, 22 Aug. 1890. Collected by W. B. Richardson. Salvin-Godman Collection. The immature male syntype is also in the collection.
aurantiiventris—Pachycephala aurantiiventris Seebohm, 1891 39
Ibis, (6) 3:96. HOLOTYPE, Adult male. Reg. no. 1898.9.20.1018. Bua Island, Vanua Levu, Fiji Islands, 28 April 1876. Collected by E. L. Layard. Seebohm Bequest.
aurantiivertex—Heterocercus aurantiivertex P. L. Sclater and Salvin,
1880
The state of the s
aurata—Geocichla aurata Sharpe, 1888
aureifrons—Zosterops aureifrons Wallace, 1864
Proc. zool. Soc. Lond., 1863 (publ. 1864): 493. SYNTYPE, Adult. Reg. no. 1863.5.12.898. Flores, 1862. Collected by A. R. Wallace. Wallace Collection. There are two other syntypes in the collection (see also appendix,
Corrections and Additions, vol. 2).
aureiloris—Zosterops aureiloris Ogilvie-Grant, 1895
collection, and two in the Cambridge University Zoological Museum.
aureiventris—Myiobius aureiventris P. L. Sclater, 1873
and the two other syntypes in the concention.

- auricollis 45 aureoflavus—Ploceus aureoflavus Smith, 1839 61 Illustrations of the zoology of South Africa, Aves: text to pl. 30. HOLOTYPE, Adult (relaxed mount). Reg. no. 1845,7.6.20. Western Africa (probably Zanzibar, vide Peters, Checklist of the Birds of the World, 15 (1962): 40). Collected by and purchased of A. Smith. aureola—Sylvicola aureola Gould, 1839 HOLOTYPE, Adult (relaxed mount). Reg. no. 1856.3.15.4. Galapagos Islands. September (the date 4 Jan. 1837 on the specimen label appears to refer to the date when it was received by Gould). Collected by C. Darwin and presented by him, probably through the Zoological Society of London. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 283. aureolimbatus—Prionochilus aureolimbatus Wallace, 1865 Proc. zool. Soc. Lond., 1865.: 477, pl. 29. HOLOTYPE, Adult male.
- Reg. no. 1873.5.12.1035. (Tondana, on label), North Celebes. 1859. Collected by A. R. Wallace. Wallace Collection.
- aureum—Malacopteron aureum Eyton, 1845 21 Ann. Mag. nat. Hist., (1) 16:228. HOLOTYPE, Adult. Reg. no. 1846.3.4.34. Malacca. Collected and presented by Capt. A. Charlton.
- aureus—Anthoscopus parvulus aureus Bannerman. 1939 41 Bull. Brit. Orn. Cl., 59:41. HOLOTYPE, Adult female. Reg. no. 1935.7.17.98. Bole, Northern Territories, Gold Coast, 10 Jan. 1935. Collected and presented by W. P. Lowe and Miss F. Waldron.
- 21 no. 1888.4.1.448. Togian Islands, Gulf of Tomini, Celebes (6,000', Aug. 1871, vide Meyer, Ibis, (4) 3 (1879): 130. Collected by A. B. Meyer. Tweeddale Collection.
- auriceps—Myiobius auriceps Gould, 1839 Zoology of the voyage of H.M.S. Beagle, 3, Birds: 47. HOLOTYPE, Adult (male, vide Sclater, Cat. Bds. Brit. Mus., 14 (1888): 210). (relaxed mount). Reg. no. 1855.12.19.172. Buenos Ayres, August. Collected by C. Darwin. Purchased of the Zoological Society of London.
- auriceps—Nectarinia auriceps G. R. Gray, 1860 . 46 Proc. zool. Soc. Lond., 1860: 348. SYNTYPE, Adult male. Reg. no. 1860.2.4.36. Batchian, Moluccas. Collected by A. R. Wallace. Wallace Collection.

There are several other syntypes in the collection.

auricollis—Nemosia auricollis P. L. Sclater, 1856. Proc. zool. Soc. Lond., 1856: 111. SYNTYPE, Adult male. Reg. no. 1857.11.18.5. Cayenne. Presented by P. L. Sclater. The second syntype is also in the collection.

auricomis—Muscicapa auricomis Latham, 1801
auricrissa—Dubusia auricrissa P. L. Sclater, 1856
auricularis—Basileuterus auricularis Sharpe, 1885 54 Cat. Bds. Brit. Mus., 10: 386. SYNTYPE, Adult. Reg. no. 1885. 3.8.666. Colombia (restricted to Bogota, vide Todd, Proc. U.S. Nat. Mus. 74, Art. 7 (1929): 60) 1869. Collected by Truebner. Salvin-Godman Collection. The other syntype from Bogota is in the collection.
auricularis—Geothlypis auricularis Salvin, 1883
auricularis—Kittacincla auricularis Swinhoe, 1864 32B Ibis, (1) 6:361. HOLOTYPE, Adult. Reg. no. 1898.10.3.563. Formosa, March 1864. Collected for R. Swinhoe. Seebohm Bequest.
aurifrons—Serinus mozambicus aurifrons W. L. Sclater and Mackworth-Praed, 1918
auriventer—Zosterops auriventer Hume, 1878
aurora—Cyrtostomus aurora Tweeddale, 1878

- aussae—Cercomela melanura aussae Thesiger and Meynell, 1934. 31 Bull. Br. Orn. Cl., 55:79. HOLOTYPE, Adult male. Reg. no. 1934.8.9.2. Aussa, Danakil, Abyssinia, 5 April 1934. Collected and presented by W. Thesiger.
- austeni—Franklinia rubescens austeni E. C. S. Baker, 1923 . . . 35

 Bull. Brit. Orn. Cl., 44: 39. Holotype, Adult male. Reg. no. 1895.7.14.1989. Lhota, Naga Hills, Assam, 19 Dec. 1875. (Collected by A. W. Channell, on label). Shelley Collection.

There are several other syntypes in the collection.

Listed as type by Ogilvie-Grant, *Ibis*, (7) 2 (1896): 61. The three other syntypes are also in the collection.

austeni—Trochalopteron austeni Godwin-Austen, 1870 32B J. Asiat. Soc. Bengal, 39, pt. 2:105. SYNTYPE, Adult. Reg. no. 1895.7.14.2253. Hengden Peak, 7,000', Jhiri River, Khasi Hills, Assam. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.

There is a second syntype in the collection, from the same locality.

australis—Ammodromus australis Maynard, 1887 49 American Exchange and Mart, 3, no. 3:33. Syntype, Adult male. Reg. no. 1906.12.7.2701. Nassau, New Providence, Bahamas,
21 Feb. 1884. Collected by C. J. Maynard. G. S. Miller Collection; the two syntypes being obtained by him from Maynard. The second syntype is also in the collection.
australis—Anthus australis Vigors and Horsfield, 1827 19 Trans. Linn. Soc. Lond., 15:229. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.30. South coast of Australia. Collected by R. Brown and presented by the Linnean Society.
australis—Cryptospiza australis Shelley, 1896
australis—Dasyornis australis Vigors and Horsfield, 1827 36 Trans. Linn. Soc. Lond., 15:232. HOLOTYPE (immature, vide Sharpe, Cat. Bds. Brit. Mus., 7 (1883):105). North of Parramatta, New South Wales. Collected by G. Caley. Presented as the type by the Linnean Society of London.
australis—Drymocataphus tickelli australis Robinson and Kloss, 1921
australis—Hyliota australis Shelley, 1886
australis—Junco phaeonotus australis van Rossem, 1938 49 Bull. Brit. Orn. Cl., 58: 133. HOLOTYPE, Adult male. Reg. no. 1899.2.1.2169. Omilteme, 8,000 ft., Guerrero, Mexico, 1 Aug. 1888. Collected by D. W. Smith. Salvin-Godman Collection.
australis—Malaconotus australis Smith, 1836
australis—Megalopterus australis Smith, 1836 62 Report of the Expedition for exploring Central Africa 1834 : 52.

SYNTYPE, Adult (relaxed mount). Reg. no. 1838.6.9.38. Beyond Kurrichane, South Africa. Collected by A. Smith. Purchased of the South African Museum, London.

Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 159.

australis—Megalotis australis Smith, 1836 Report of the Expedition for exploring Central Africa . . . 1834; Appendix: 49. SYNTYPE, Adult male (relaxed mount). Reg. no. 1843.2.28.57. Orange River (in text; type locality restricted to Colesberg, east-central Cape Province by Macdonald, *Contributions* to the Ornithology of western South Africa, 1957: 107). Collected by A. Smith and purchased of the South African Society. There are two other syntypes at the Academy of Natural Sciences,

Philadelphia.

- australis—Nannorchilus leucogaster australis Van Rossem, 1938 Bull. Br. Orn. Cl., 59:15. HOLOTYPE, Adult. Reg. no. 1890,12.20. 681. Orange Walk, Belize, British Honduras, Dec. 1887. Collected by G. F. Gaumer. Salvin-Godman Collection.
- australis—Nectarinia australis Gould, 1850. in Jardine, Contributions to Ornithology: 106. (Also in Proc. zool. Soc. Lond., 1850 (publ. Feb. 1851): 201). HOLOTYPE, Adult male. Reg. no. 1882.12.10.33. Oomaga Island, Torres Straits, Northeast Australia. Collected by Commander R. Ince, R.N. (vide Gould, Birds of Australia, 1 (1865): 584). Presented by Mrs. Ince after the death of her husband.
- australis—Stelgidocichla latirostris australis Moreau, 1941 . Bull. Br. Orn. Cl., 62:29. HOLOTYPE, Adult male, not breeding. Reg. no. 1944.8.1.2. Mbisi Forest, 8,000', near Sumbawanga, Ufipa District, Tanganyika, 2 June 1941. Collected by Salimu Asmani and presented by R. E. Moreau.
- australoabyssinicus—Anthus caffer australoabyssinicus Benson, 1942 19 Bull. Br. Orn. Cl., 63:12. HOLOTYPE, Adult male. Reg. no. 1946.5.2750. 30 miles south of Yavello, 4,500', south Abyssinia, 12 Sept. 1941. Collected and presented by C. W. Benson.
- australoabyssinicus—Mirafra poecilosterna australoabyssinicus Benson, Bull. Br. Orn. Cl., 63:13. HOLOTYPE, Adult male. Reg. no. 1946.5.2797. 20 miles south of Mega, 3,000', south Abyssinia, 15 Feb. 1942. Collected and presented by C. W. Benson.
- australoabyssinicus—Pseudonigrita arnaudi australoabyssinicus Benson, Bull. Br. Orn. Cl., 63:19. HOLOTYPE, Adult male. Reg. no. 1946.5.3141. Yavello, 4,500', south Abyssinia, 15 June 1941. Collected and presented by C. W. Benson.

30 australoabyssmicus
australoabyssinicus—Turdus tephronotus australoabyssinicus Benson, 1942
australoabyssinicus—Zosterops senegalensis australoabyssinicus Benson, 1942
awemba—Cisticola robusta awemba Lynes, 1933
axillaris—Amaurospiza axillaris Sharpe, 1888
axillaris—Arremon axillaris P. L. Sclater, 1855
axillaris—Pachycephalopsis hattamensis axillaris Mayr, 1931 37 Bull. Brit. Orn. Cl., 51:59. Holotype, Adult male. Reg. no. 1916.5.30.245. Camp 6a, 2,900', Utakwa River, Dutch New Guinea, 6 Jan. 1913. Collected by C. B. Kloss. Presented by the Committee of the Dutch New Guinea Expedition.
axillaris—Pratincola axillaris Shelley, 1884
aximensis—Pedilorhynchus comitatus aximensis W. L. Sclater, 1924 37 Bull. Brit. Orn. Cl., 45:45. HOLOTYPE, Adult male. Reg. no. 1911.12.18.219. Axim, Gold Coast, West Africa, 27 Feb. 1911. Collected by W. P. Lowe. Presented by Capt. E. C. Hardy, R.N.
aylmeri—Argya aylmeri Shelley, 1885

- azizi—Ammomanes deserti azizi Ticehurst and Cheesman, 1924. 17 Bull. Br. Orn. Cl., 45: 20. Holotype, Adult male. Reg. no. 1924.11.30.29. Jabal Ghanima, Hasa Province, Arabia, 12 Jan. 1924. Collected by R. E. Cheesman. Cox-Cheesman Collection.
- azoricus—Regulus cristatus var. azoricus Seebohm, 1883 . . . 35

 A history of British birds, 1: 451. Lectotype, Adult female. Reg.
 no. 1879.4.5.307. St. Michael's, Azores, April 1865. Collected by F. D. Godman. Presented by F. D. Godman and O. Salvin.

A male and a female syntype were listed by Seebohm, Cat. Bds. Brit. Mus., 8 (1883): 82, but the male cannot be traced. Murphy and Chapin, Amer. Mus. Novit. 384 (1929): 15 designated the female as the (lecto)type.

azureocapilla—Myiagra azureocapilla Layard, 1875 38

Ibis, (3) 5:434. Syntype, Adult male. Reg. no. 1879.4.5.1829.

Taviani, Fiji Islands. Collected by Liardet. Salvin-Godman Collection.

This specimen, now tail-less, is inscribed on the label "Mr Layard told me this was a new species". The labels of two other specimens in the collection, collected by Layard in Aug. 1875, have been marked "type" in a later hand but may not have been before the author at the time of the description.

- babaulti—Scotocichla fuscicapilla babaulti Wells, 1919. . . . 32B Bull. Br. Orn. Cl., 39:69. Holotype, Adult male. Reg. no. 1898.10.3.725. Trincomalee, north Ceylon, 16 Nov. 1872. Collected by W. V. Legge, Seebohm Collection.
- babylonicus—Acrocephalus babylonicus Ticehurst, 1920 . . . 35
 Bull. Brit. Orn. Cl., 41:12. HOLOTYPE, Adult male. Reg. no. 1923.6.19.3. Basra, Lower Mesopotamia, 22 April 1917. Collected by R. E. Cheesman and Sir Percy Cox. Presented by the Bombay Natural History Society.

This specimen was not listed by Sharpe, Cat. Bds. Brit. Mus., 7 (1883) but the mounting was relaxed at that date and it was probably not available.
baetica—Calendrella baetica Dresser, 1873
bafirawari—Bradornis bafirawari Bannerman, 1924 37 Bull. Brit. Orn. Cl., 45:41. HOLOTYPE, Adult male. Reg. no. 1924.11.21.1. Wajheir, 3,000′, 100 miles north-east of the Lorian Swamp, Juba Land Province, Kenya, 7 Nov. 1910. Collected by Bafirawar for F. J. Jackson. Presented by F. J. Jackson.
bahiae—Sclerurus mexicanus bahiae Chubb, 1919
baicalensis—Motacilla baicalensis Swinhoe, 1871
baicalensis—Poecile baicalensis Swinhoe, 1871
baileyi—Ixulus flavicollis baileyi E. C. S. Baker, 1914 32B Bull. Br. Orn. Cl., 35: 17. Holotype, Adult (male on label). Reg. no. 1914.10.5.23. Tembang (on label), 7,000', Mishmi Hills, north-east India, 10 Oct. 1913. Collected and presented by F. M. Bailey.
baileyi—Leioptila capistrata baileyi Kinnear
bairdi—Empidonax bairdi P. L. Sclater, 1858

bairdi—Setophaga bairdi Salvin, 1878
bairdi—Thryothorus bairdi Salvin and Godman, 1880 28 Biol. CentrAmer.; Aves, 1:95. Holotype, Adult male. Reg. no. 1885.3.6.382. Oaxaca, Mexico. Collected by A. Sallé. Salvin-Godman Collection.
bakeri—Pomatorhinus ruficollis bakeri Harington, 1914 32B J. Bombay nat. Hist. Soc., 23:336. Holotype, Adult male. Reg. no. 1886.10.1.3610. Shillong, Assam, Nov. 1877. Collected by J. Cockburn. Hume Collection.
bakeri—Turdinulus epilepidotus bakeri Harington, 1913 32B Bull. Br. Orn. Cl., 33:44. HOLOTYPE, Adult. Reg. no. 1900. 12.16.416a. Na Noi, South Shan States, 25 March 1899. Collected and presented by Col. G. Rippon.
bakeri—Turdinus macrodactylus bakeri Hachisuka, 1926 32B Bull. Br. Orn. Cl., 47:54. HOLOTYPE, Adult male. Reg. no. 1910.12.27.295. Lam ra, Trang, north Malay Peninsula, 19 Jan. 1910. Presented by the Kuala Lumpur Museum, Malaya.
balfouri—Cinnyris balfouri P. L. Scalter and Hartlaub, 1881
balstoni—Sericornis balstoni Ogilvie-Grant, 1909
balstoni—Zosterops balstoni Ogilvie-Grant, 1909
baltistanicus—Acanthis flavirostris baltistanicus R. and A. Meinertzhagen, 1926

- bambarae—Thamnolaea cinnamomeiventris bambarae Bates, 1928 31 Bull. Br. Orn. Cl., 49:32. Holotype, Adult male. Reg. no. 1928.11.21.2. Kulikoro, below Bamako on River Niger, French Sudan, 7 July 1928. Collected and presented by G. L. Bates.
- bambuluensis—Apalis jacksoni bambuluensis Serle, 1949 . . . 35

 Bull. Brit. Orn. Cl., 69:55. HOLOTYPE, Adult male. Reg. no. 1949.14.16. Near Bambulue Lake, 6,000′, Bamenda, Cameroons, 5°50′ N. 10°10′ E., 14 Sept. 1948. Collected and presented by W. Serle.
- bamendae—Andropadus tephrolaemus bamendae Bannerman, 1923 21 Bull. Br. Orn. Cl., 44: 4. HOLOTYPE, Adult male. Reg. no. 1923. 10.26.264. Near Bamenda, 5,500', Cameroon Highlands, 23 Aug. 1922. Collected by and purchased of G. L. Bates.

- bamingui—Psalidoprocne bamingui Boyd Alexander, 1908 . . . 18

 Bull. Br. Orn. Cl., 21:88. SYNTYPE, Adult male. Reg. no.
 1911.12.23.1315. Bamingui River, Upper Shari River, 13 Aug.
 1905. Collected and bequeathed by Boyd Alexander.

 This is the author's selected male. His selected female and two immature syntypes are in the collection.
- banguey—Dissemurus paradiseus banguey Chasen and Kloss, 1929. 64
 J. Orn. 1929 (2): 120. Holotype, Adult male. Reg. no. 1947.
 60.27. Banguey Island, North Borneo, 3 Sept. 1927. Collected by F. N. Chasen and C. B. Kloss. Presented by the Raffles Museum, Singapore.
- bannermani—Anthus richardi bannermani Bates, 1930 19
 Bull. Br. Orn. Cl., 51:48. HOLOTYPE, Adult male. Reg. no. 1930.12.3.1. Birwa Peak, 4,800', Kono District, Sierra Leone, 18 July 1930. Collected by and purchased of G. L. Bates.

Mackworth-Praed, 1943
bannermani—Hirundo abyssinica bannermani C. H. B. Grant and Mackworth-Praed, 1942
bannermani—Plocepasser superciliosus bannermani C. H. B. Grant and Mackworth-Praed, 1943 61 Bull. Br. Orn. Cl., 64:18. HOLOTYPE, Adult male. Reg. no. 1927. 11.5.865. Gomit River, 4,000', Big Abbai, 60 miles south of Lake Tana, northern Abyssinia, 21 Jan. 1927. Collected by and purchased of R. E. Cheesman.
bansoensis—Cyanomitra oritis bansoensis Bannerman, 1922 46 Bull. Brit. Orn. Cl., 43:8. HOLOTYPE, Adult male. Reg. no. 1922.11.25.224. Banso Mountains, 6,000', north of Kumbo, Nigerian-Cameroon Highlands, 26 Sept. 1921. Collected by and purchased of G. L. Bates.
bansoensis—Nesocharis shelleyi bansoensis Bannerman, 1923 . 60 Bull. Brit. Orn. Cl., 44:3. HOLOTYPE, Adult male. Reg. no. 1923. 10.26.422. Banso Mountains, 6,000', north of Kumbo, Cameroon, 13 Sept. 1922. Collected by and purchased of G. L. Bates.
bansoensis—Riparia rufigula bansoensis Bannerman, 1923 18 Bull. Br. Orn. Cl., 44:5. HOLOTYPE, Adult male. Reg. no. 1923.10.26.146. Bamenda, 4,750', Cameroon Highlands, 25 Aug. 1922. Collected by and purchased of G. L. Bates.
banyumas—Muscicapa banyumas Horsfield, 1821 37 Trans. Linn. Soc. Lond., 13:146. SYNTYPE, Adult male (relaxed mount). Reg. no. 1880.1.1.873. Province of Banyumas, Java. Collected by T. Horsfield. Presented by the Honourable East India Company Museum. A second syntype is also in the collection.
baraka—Merula baraka Sharpe, 1903
baraka—Sylviella baraka Sharpe, 1897

barakae—Bradypterus barakae Sharpe, 1906
barakae—Nectarinia barakae Sharpe, 1902
barakae—Parus barakae Jackson, 1899
barakae—Turdinus barakae Jackson, 1906
barat—Pycnonotus bimaculatus barat Robinson and Kloss, 1920 21 J. Straits Branch, roy. Asiat. Soc., 81:103. SYNTYPE, Adult male. Reg. no. 1947.60.45. Siolak Daras, 3,000', west Sumatra, 15 March 1914. Collected by H. C. Robinson and C. B. Kloss and presented by the Raffles Museum, Singapore. The female syntype did not come to the British Museum.
barbata—Muscicapa barbata Latham, 1801
barnesi—Parisoma layardi barnesi Vincent, 1948
barnesi—Saxicola barnesi Oates, 1890
baroica—Saxicola baroica Smith, 1840

baroni—Buarremon baroni Salvin, 1895
baroni—Siptornis baroni Salvin, 1895
barratti—Bradypterus barratti Sharpe, 1876
bartletti—Dendrocincla bartletti Chubb, 1919
bartletti—Malimbus bartletti Sharpe, 1890
bartletti—Paradisea bartletti W. Goodwin, 1860
bartoni—Microeca griseiceps bartoni Ogilvie-Grant, 1915 37. Ibis, (10) 2, Supplement : 174. SYNTYPE. Adult male. Reg. no. 1913.3.6.304. Owen Stanley Range, 5,000'. Dutch New Guinea, April 1908. Collected by W. Goodfellow. Purchased of E. J. Johnstone. This is the described male, the described female syntype is also in the collection.
bartoni—Pachycephala soror bartoni Ogilvie-Grant, 1915 39 Ibis, (10) 2, Jubilee Supplement: 89. Syntype, Adult male. Reg. no. 1913.3.6.324. Owen Stanley Mountains, 5,000', British New

A Smith specimen in the collection was listed as type by Seebohm,

Cat. Bds. Brit. Mus., 5 (1881): 399.

Guinea, April 1908. Collected by W. Goodfellow. Purchased of E. Johnstone.

This is the author's selected male.

- basilanica—Dendrobiastes basilanica Sharpe, 1877 37

 Trans. Linn. Soc. Lond., (2) Zool., 1:332. An adult female, Reg. no. 1896.6.6.400, collected by J. B. Steere on Isabella de Basilan, Philippines, on 18th Nov., was listed as the type by Hachisuka, Birds of the Philippines, 2 (1934–5):302, but from the date when it was received, and from the fact that Steere, List of birds and mammals collected by the Steere Expedition to the Philippines (1890), states that on the second expedition he visited Basilan in November, and lists this species as collected on that island, it seems certain that this specimen has no type status.
- basilanica—Ptiocichla basilanica Steere, 1890 32B

 List of the Birds and Mammals collected by the Steere Expediton
 to the Philippines: 18 (reprinted in Ibis, (6) 3 (1891): 312, pl. 7.).

 SYNTYPE, Adult male. Reg. no. 1896.6.6.397. Basilan, Philippines, 7 Nov. Collected by J. B. Steere. Steere Collection.
- basilanica—Rhinomyias ruficauda basilanica Hachisuka, 1932 . 37

 Bull. Brit. Orn. Cl., 52:110. HOLOTYPE, Adult male. Reg. no. 1887.12.30.1387. (not 1889 as stated in the description). Basilan, Philippines, June 1878. Collected by A. H. Everett. Tweeddale Collection.
- basilanicus—Anthreptes malacensis basilanicus Hachisuka, 1926. . 46

 Bull. Brit. Orn. Cl., 47:68. Holotype, Adult male. Reg. no. 1896.6.6.483. Basilan Island, Philippines, 19 Nov. Collected by J. B. Steere. Steere Collection.

basilanicus—Oriolus basilanicus Ogilvie-Grant, 1896 63 Ibis, (7) 2:532. SYNTYPE, Adult male. Reg. no. 1896.6.6.39. Basilan, Philippines, 6 Nov. Collected by J. B. Steere. Steere Collection. There are five other syntypes in the collection. Ogilvie-Grant labelled this specimen as the type but did not designate it in the text.
basilanicus—Poliolophus basilanicus Steere, 1890 21 List of Birds and Mammals collected by the Steere Expedition to the Philippines: 19 (reprinted in Ibis, (6) 3 (1891): 313). SYNTYPE, Adult male. Reg. no. 1896.6.6.324. Basilan, Philippines, 14 Nov. 1887. Collected by and purchased of J. B. Steere. The female syntype is also in the collection.
bassetti—Zosterops bassetti Sharpe, 1894
basutoensis—Fringillaria capensis basutoensis Vincent, 1950 49 Bull. Brit. Orn. Cl., 70: 14. HOLOTYPE, Adult male. Reg. no. 1950.12.2. Source of the Lekhalabaletsi River, 9,600', Basutoland, 29°18' S., 29°25' E., 15 July 1949 (25 July, on label). Collected and presented by J. Vincent.
batakana—Munia atricapilla batakana Chasen and Kloss, 1929 . 60 Bull. Raffles Mus., 2:23. Holotype, Adult female. Reg. no. 1947.60.69. Brastagi, 1,390 metres, Simeloengan, Batak Highlands, North-east Sumatra, 28 Jan. 1919. Collected by A. C. van Heyst. Presented by the Raffles Museum, Singapore.
batchianensis—Anthus batchianensis G. R. Gray, 1869 19 Handlist of Birds, 1: 251 (nomen nudum). A Wallace specimen was listed as type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 614.
batesi—Batis bella batesi Bannerman, 1923
batesi—Bledi batesi Sharpe, 1903
batesi—Callene batesi Bannerman, 1922

batesi—Chloropeta batesi Sharpe, 1904
batesi—Chlorophoneus batesi Sharpe, 1908
batesi—Cinnyris batesi Ogilvie-Grant, 1908
batesi—Euplectes axillaris batesi Delacour, 1933 61 Bull. Br. Orn. Cl., 53: 181. HOLOTYPE, Adult male. Reg. no. 1932.8.6.8. Niger River, between Tillabery and Ansonge, French Sudan, 30 July 1931. Collected by and purchased of G. L. Bates.
batesi—Geocichla batesi Sharpe, 1905
batesi—Lanius excubitor batesi C. H. B. Grant and Mackworth-Praed, 1951
batesi—Melirrhophetes batesi Sharpe, 1886
batesi—Mirafra batesi Bannerman, 1923
batesi—Othyphantes batesi Sharpe, 1908

- bayleyi—Leioptila capistrata bayleyi Kinnear, 1939 32B Ibis, (14) 3:752. HOLOTYPE, Adult female (on label; male in text). Reg. no. 1935.4.5.156. Taktoo, near Sakdan, east Bhutan, 27 Oct. 1934. Collected and presented by F. Ludlow and G. Sherriff.

The subspecific name was corrected to *baileyi* by Ludlow, *Ibis*, 86 (1944) : 82–3.

The second syntype is also in the collection.

- beccarii—Dicaeum beccarii Robinson and Kloss, 1916. 45

 J. Straits Branch Roy. Asiat. Soc., 73: 278. HOLOTYPE, Adult male. Reg. no. 1947.60.63. Korinchi Peak, 7,300', West Sumatra, 26 April 1914. Collected by H. C. Robinson and C. B. Kloss. Presented by the Raffles Museum, Singapore.
- bechuanae—Prinia superciliosa bechuanae Macdonald, 1941 . . . 35

 Bull. Brit. Orn. Cl., 62:28. Holotype, Adult male. Reg. no. 1911.12.16.102. Mababe Flats, (3,000', on label), North-western Bechuanaland, 8 Aug. 1909. Collected by R. B. Woosman. Presented by the R. B. Woosman and Hon. C. Legge's Ngamiland Expedition.
- bedfordi—Pholidornis bedfordi Ogilvie-Grant, 1904 60
 Bull. Brit. Orn. Cl., 14:56. SYNTYPE, Adult male. Reg. no. 1904.6.22.207. Santa Isabel, north Fernando Po, 25 Dec. 1903. Collected by E. Seimund. Presented by the Duke of Bedford, Mrs P. Sladen, and the Hon. W. Rothschild.

The female syntype is also in the collection.

bedfordi—Sitta bedfordi Ogilvie-Grant, 1909
bedfordi—Trochocercus bedfordi Ogilvie-Grant, 1907 38 Bull. Brit. Orn. Cl., 19:40. SYNTYPE, Adult male. Reg. no. 1906.12.23.1532. Mawambi, 3,000', Eastern Congo Forest, 17 Oct. 1906. Collected by R. B. Woosnam. Presented by the subscribers to the Ruwenzori Expedition. The described female syntype is also in the collection.
beesleyi—Chersomanes albofasciata beesleyi Benson, 1966 17 Bull. Br. Orn. Cl., 86:77. HOLOTYPE, Adult male. Reg. no. 1965.37.1. Masai Plains, 30 miles north of Arusha, northern Tanzania, 2 Nov. 1965. Collected by J. S. S. Beesley and presented by C. W. Benson.
bella—Aethopyga bella Tweeddale, 1877
bella—Loxia bella Latham, 1801
belli—Cisticola belli Ogilvie-Grant, 1908
belli—Pytelia belli Ogilvie-Grant, 1907
belli—Eremomela griseoflava belli C. H. B. Grant and Mackworth-Praed, 1946

Proc. zool. Soc. Lond., 1862:111. SYNTYPE, Adult. Reg. no. 1888.1.13.860. Bogota, Columbia. Sclater Collection. The second syntype is in the collection. bengalensis—Timalia bengalensis Godwin-Austen, 1872 J. Asiat. Soc. Bengal, 41, pt. 2:143. HOLOTYPE, Adult. Reg. no. 1895.7.14.1894. Head of Jhiri River, Khasia Hills, Assam. Collected by H. H. Godwin-Austen. Godwin-Austen Collection. benguelae—Parus afer benguelae Hall and Traylor, 1959 Bull. Brit. Orn. Cl., 79:45. HOLOTYPE, Adult male. Reg. no. 1957.35.3. 12 miles south-east of Benguela, West Angola. 12 Sept. 1957. Collected by G. W. Lathbury. Presented by B. P. Hall. benguelensis—Alaemon benguelensis Sharpe, 1904 17 Ibis, (8) 4:361. HOLOTYPE, Adult. Reg. no. 1873.12.10.130. Benguela, Angola. Collected by and purchased of J. J. Monteiro. benguellensis—Amydrus nabouroup benguellensis Neumann, 1903 Orn. Mber. 11:184. HOLOTYPE, Adult. Reg. no. 1873.12.10.53. Benguella, Angola. Collected and presented by J. J. Monteiro. benguellensis—Bradypterus brachypterus benguellensis Bannerman, Bull. Brit. Orn. Cl., 47:147. HOLOTYPE, Adult male. Reg. no. 1928.10.25.37. Chicuma, Benguella Province, Angola, 9 Feb. 1927. Collected and presented by Admiral H. Lynes. benguellensis—Karrucincla schlegelii benguellensis W. L. Sclater, 1928 31 Bull. Br. Orn. Cl., 49:15. HOLOTYPE, Adult male. Reg. no. 1906.12.4.127. Huxe, Benguella, Angola, 5 Sept. 1905. Collected by and purchased of Dr W. J. Ansorge. benguellensis—Passer motitensis benguellensis Lynes, 1926 Ibis, (12) 2:380. HOLOTYPE, Adult male. Reg. no. 1906.12.4.299. Huxe, Benguella, Angola, 28 Aug. 1905. Collected by W. J. Ansorge. Ansorge Collection. Proc. zool. Soc. Lond., 1832: 90. SYNTYPE, Adult. Reg. no. 1880.1.1.4438. Deccan, India. Collected by W. H. Sykes. bennettii—Orthotomus bennettii Sykes, 1832 Presented by the Honourable East India Company Museum. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 217. There is another syntype in the collection not mentioned by Sharpe (op.cit.). bensoni—Anumomanes deserti bensoni R. Meinertzhagen, 1933. Bull. Br. Orn. Cl., 53:151. HOLOTYPE, Adult male. Reg. no. 1939.12.9.279. Tamsnigat, 6,800', Ahaggar Plateau, Sahara, 25 Feb. 1931. Collected by R. Meinertzhagen. Rothschild Bequest.

bellus—M viobius bellus P. L. Sclater, 1862 . . .

o i belisoni
bensoni—Apalis murina bensoni C. H. B. Grant and Mackworth-Praed, 1937
bensoni—Artisornis metopias bensoni Vincent, 1935 35 Bull. Brit. Orn. Cl., 55: 174. HOLOTYPE, Adult male. Reg. no. 1935.3.11.3. Chongoni Mountain, 6,000', 10 Feb. 1935. Collected and presented by J. Vincent.
bensoni—Cynnyris mediocris bensoni Williams, 1953 46 Bull. Brit. Orn. Cl., 73:10. HOLOTYPE, Adult male. Reg. no. 1952.48.1. Dedza Mountains, 7,000', Nyasaland, 19 July 1951. Collected and presented by C. W. Benson. The spelling of the generic name is a printer's error.
bensoni—Sheppardia bensoni Kinnear, 1938 31 Bull. Br. Orn. Cl., 58:138. HOLOTYPE, Adult male. Reg. no. 1938.6.17.1. Near Nkata Bay, 1,900', Chinteche District, West Nyasa District, Nyasaland, 15 April 1938. Collected by and purchased of C. W. Benson.
bensonorum—Acanthis flavirostris bensonorum R. Meinertzhagen, 1934 59 Ibis, (13) 4:55. HOLOTYPE, Adult male. Reg. no. 1965.M.51. Grogarry, South Uist, Outer Hebrides, 8 Nov. 1932. Collected by R. Meinertzhagen. Meinertzhagen Collection.
bentet—Lanius bentet Horsfield, 1821
berberae—Cisticola cisticola berberae R. Meinertzhagen, 1922 . 35 Ibis, (11) 4:12. HOLOTYPE, Adult male. Reg. no. 1923.8.7.2873. near Berbera, 3,000′, British Somaliland, 13 Jan. 1919. Collected by Sir G. F. Archer. Presented by Col. Stephenson Clarke.
berlepschi—Cyanolesbia berlepschi Hartert, 1898 25 Bull. Brit. Orn. Cl., 8: 16. Syntype, male, not fully grown. Reg. 1887.3.22.1364. Forest of Caripé, Venezuela, (June 1867, on label). Collected by A. Goering. Salvin-Godman Collection. Of the five remaining syntypes referred to by the author, Hartert, Novit. Zool., 27 (1920): 411, states that "male and female types" are in the American Museum of Natural History.

berlepschi—Phyllmyias berlepschi P. L. Sclater, 1887 9 Proc. zool. Soc. Lond., 1887.49. HOLOTYPE, Adult. Reg. no. 1888.1.13.435. Bahia, East Brazil. Sclater Collection; from Von Berlepsch.
berlepschi—Thryothorus coraya berlepschi Brabourne and Chubb, 1912
bermudiana—Carduelis carduelis bermudiana Kennedy, 1913 . 59 Bull. Brit. Orn. Cl., 33:33. HOLOTYPE, Adult male. Reg. no. 1913.10.29.38. Ireland Island, Bermuda, 23 Dec. 1912. Collected and presented by J. N. Kennedy.
bernieri—Malurus bernieri Ogilvie-Grant, 1909
bertrandi—Hyphantornis bertrandi Shelley, 1893 61 Ibis, (6) 5:23. SYNTYPE, Adult male. Reg. no. 1892.9.10.181. Zomba, Nyasaland, 7 Aug. 1891. Collected by A. Whyte. Presented by H. H. Johnston. There are two other syntypes in the collection.
bertrandi—Laniarius bertrandi Shelley, 1894
besti—Dicaeum besti Steere, 1890
besuki—Aethostoma pyrrogenys besuki Kloss, 1931 32B Treubia, 13:346. HOLOTYPE, Adult female (bill broken). Reg. no. 1947.60.51. Tamansari, near Banjoewangi, east Java, 20 Jan. 1920. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore.
bhamoensis—Stachyrhidopsis bhamoensis Harington, 1908 32B Ann. Mag. nat. Hist., (8) 2:245. SYNTYPE, Adult male. Reg. no. 1908.8.2.13. Sinlum-Kaba, Bhamo District, Upper Burma, 20 April 1908. Collected and presented by H. H. Harington. There are several other syntypes in the collection.

bichenovii—Fringilla bichenovii Vigors and Horsfield, 1827 60 Trans. Linn. Soc. Lond., 15:258. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.49. Shoal water Bay, Broad Sound, Queensland, Australia, Sept. 1802. Collected by R. Brown. Presented by the Linnean Society of London as the type.
bicolor—Amadina bicolor Fraser, 1843
bicolor—Chimarrhornis bicolor. See appendix, Corrections and Additions, vol. 2
bicolor—Crateropus bicolor Jardine, 1831
bicolor—Grallina bicolor Vigors and Horsfield, 1827 . 37 and 66 Trans. Linn. Soc. Lond., 15:233. Holotype, Adult male. Reg. no.1863.7.7.37. Prospect Hill, New South Wales. Collected by R. Brown. Presented by the Linnean Society as the type. The specimen is not a Grallina but an example of Petroica (Melanodryas) cucullata of the Muscicapidae. A specimen of Grallina melanoleuca from the Linnean Society was mistakenly referred to as the type of G. bicolor by Sharpe, Cat. Bds. Brit. Mus., 3 (1877): 273.
bicolor—Megabyas bicolor Elliott, 1859
bicolor—Merula bicolor Layard, 1876
bicolor—Saxicola bicolor Sykes, 1832
biddulphi—Podoces biddulphi Hume, 1874
bihe—Prinia flavicans bihé Boulton and Vincent, 1936 35 Bull. Brit. Orn. Cl., 57:7. HOLOTYPE, Adult male. Reg. no. 1931.12.21.94. Vouga, 5,800', Bihé, Central Angola, 11 Feb. 1931. Collected by Admiral H. Lynes and J. Vincent. Presented by Admiral H. Lynes.

bimaculata—Monarcha bimaculata G. R. Gray, 1860 38

Proc. zool. Soc. London., 1860: 352. SYNTYPE, Adult (male, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 420). Reg. no. 1860. 2.4.34. Batchian, Moluccas. Collected by A. R. Wallace. Purchased of Stevens.

There are several other syntypes in the collection.

bimaensis—Dicrurus bimaensis Wallace, 1863 64

Proc. zool. Soc. Lond., 1863: 492. SYNTYPE, Adult. Reg. no. 1873.5.12.1954. Flores, 1862. Collected by A. R. Wallace, Wallace Collection.

Wallace used a Temminck MS. name previously used by Bonaparte, Consp. Gen., 1 (1850): 352, as a nomen nudum. Wallace's description was based on both Lombok and Flores specimens, but Sumbawa was included in the localities. Rensch, Mitteil. zool. Mus. Berlin, 17 (1931): 589 restricted the type locality to Flores and Sumbawa. Vaurie, Bull. Amer. Mus. Nat. Hist., 93 (4) (1949): 297, considers that the name is valid for both Flores and Lombok birds but not for Sumbawa. In view of this, although one of the British Museum specimens Reg. no. 1873.5.12.1954., is marked on the label with the red stripe used by Wallace to denote key or type specimens, a specimen from Flores has been selected, although it appears a little small in size.

The second syntype cannot be traced in the collection.

This specimen was labelled as the type, and confirmed as the holotype by Swinhoe, *Proc. zool. Soc. Lond.*, 1863: 293.

- blanfordi—Cyrtostomus pectoralis blanfordi E. C. S. Baker, 1921 46
 Bull. Brit. Orn. Cl., 41:71. HOLOTYPE, Adult male. Reg. no.
 1886.12.1.970. Kondol Island, Nicobars, (13 March 1873, on label). Collected by W. Davidson. Hume Collection.
- There are two other syntypes in the collection. blanfordi—Laniarius blanfordi Sharpe, 1882 .

blanfordi—Montifringilla blanfordi Hume, 1876	
blanfordi—Psalidoprocne blanfordi Blundell and Lovat, 1899 Bull. Br. Orn. Cl., 10:20. HOLOTYPE, Adult male. Reg. no. 1900.1.3.110. Bilo, Abyssinia, 4 March 1899 (in text; 11 March on label). Collected by H. W. Blundell and Lord Lovat.	
blanfordi—Pycnonotus blanfordi Jerdon, 1862	
blanfordi—Sylvia blanfordi Seebohm, 1878	
blanfordi—Tephrocorys blanfordi Shelley, 1902 The Birds of Africa, 3:128, pl. 21. HOLOTYPE, Adult male. Reg. no. 1869.10.16.244. Senafé, 7,500', Tigré District, Abyssinia, 24 Feb. 1868. Collected by W. T. Blanford and presented by the Governor and Council of Bombay.	
blissetti—Diaphorophyia blissetti Sharpe, 1872	37
bloxami—Phytotoma bloxami Jardine and Selby, 1827. Illustration of Ornithology 1, pt. 1: pl. 4. HOLOTYPE, Adult male (relaxed mount). Reg. no. (recently assigned) 1968.46.1. Valparaiso, Chile. Collected by Lord Byron R.N. on the voyage of H.M.S. Blonde. Presented by the Admiralty.	
blundelli—Anaplectes blundelli Ogilvie-Grant, 1900 Ibis, (7) 6:132. Holotype, Adult male. Reg. no. 1900.1.3.118. Beni Schongul, Abyssinia, 3 May 1899. Collected by Lord Lovat. Presented by H. W. Blundell and Lord Lovat.	
	59
[blythi]—Sylvia curruca blythi Ticehurst and Whistler, 1933. Ibis, (13) 3: 556. HOLOTYPE?, Adult male. Reg. no. 1891.9.1. 101. Cawnpore, 17 Jan. 1876. Collected by W. E. Brook. Seebohm Bequest.	

The authors intended to designate a holotype but inadvertently made this a new name for <i>Curruca affinis</i> Blyth, 1845. The type would be invalid, <i>vide</i> , <i>International Code Zool. Nomencl.</i> (1961): 75, but application has been made for validation of the above holotype, <i>vide</i> Watson, <i>Bull. Zool. Nomencl.</i> 26 (1969): 39–41.
Catalogue of African birds: 63. SYNTYPE, Adult male. Reg. no. no. 1878.12.31.404. Huilla, Angola. Collected by J. Anchieta. Sharpe Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 227, but there is another syntype in the collection.
Proc. zool. Soc. Lond., 1892: 228, pl. 20. HOLOTYPE, Adult. Reg. no. 1892.5.31.3. San Miguel, St. Thomas Island, West Africa. Collected by F. Newton. Presented by R. B. Sharpe.
Monograph of the Nectariniidae, : 21, pl. 6. HOLOTYPE, Adult male. Reg. no. 1895.9.9.14. Caconda, Benguela (op. cit., p. xix). Collected by Anchieta. Shelley Collection, presented to him by Professor Barboza du Bocage of Lisbon.
Proc. zool. Soc. Lond., 1855: 109, pl. 101. HOLOTYPE, Adult. Reg. no. 1844.5.15.114. Bogota. Sclater Collection.
J. Linn. Soc. Lond., 17: 422. HOLOTYPE, Adult. Reg. no. 1884. 5.1.26. Sassa, Nyam-Nyam country, Central Africa, Nov. 1882. Collected by and purchased of F. Bohndorff.
Coliviana—Chrysomitris boliviana Sharpe, 1888
Proc. zool. Soc. Lond., 1852 (publ. Nov. 1853): 34, pl. 47. SYNTYPE, Adult male. Reg. no. 1884.5.15.211. Bolivia. Collected by Bridges. Sclater Collection. Labelled as the type, and listed as such by Sclater, Catalogue of a collection of American birds, (1862): 12.
Coliviana—Poospiza boliviana Sharpe, 1888
Cat. Bds. Brit. Mus., 10:402. HOLOTYPE, Adult. Reg. no. 1885.3.8.683. Yuyo, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.

bolivianus—Saltator cayanus bolivianus Chubb, 1921 56 Ann. Mag. Nat. Hist. (9) 8: 445. HOLOTYPE, Adult male. Reg. no. 1902.3.13.359. Chulumani, 2,000 m., Bolivia (57° W., 16° E., on label). 12 Jan. 1901. Collected by P. O. Simons. Purchased of Rosenberg.	0
bonapartii—Poospiza bonapartii P. L. Sclater, 1867 4 Proc. zool. Soc. Lond., 1867 : 341, pl. 20. SYNTYPE, Adult male. Reg. no. 1885.2.10.329. Lima, Peru, Jan. 1867. Collected by W. Nation. Sclater Collection. The female syntype is also in the collection.	9
bonapartii—Todopsis bonapartii G. R. Gray, 1859 3 Proc. zool. Soc. Lond., 1859: 156. SYNTYPE, Adult female. Reg. no. 1858.3.10.54. Aru Islands, 1857. Collected by A. R. Wallace. Wallace Collection. Gray had previously, Proc. zool. Soc. Lond., 1858: 176, pl. 134. described the male syntype, which is also in the collection, under Quoy and Gaimard's name of Todus cyanocephalus, but subsequently redescribed it and included the description of the female. See also Gould, Birds of New Guinea 2, pt. 8 (1878): pl. 22.	6
boranensis—Argya aylmeri boranensis Benson, 1947	В
borbae—Thamnophilus borbae Pelzeln, 1869. Zur Ornithologie Brasiliens, : 75, 140. SYNTYPE, Adult male. Reg. no. 1889.9.20.27. Borba, Brazil, July 1830. Collected by J. Natterer. Sclater Collection. Listed as "authentic specimen" by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 186.	4
borealis— Cettia minuta borealis Campbell, 1892	5
borealis—Cryptospiza borealis Percival, 1912 6 Bull. Brit. Orn. Cl., 27: 76. SYNTYPE, Adult male (immature, on label). Reg. no. 1912.2.6.1. Mt. Urguess, 7,000', north of Guasso Nyiro, Kenya, 25 Aug. 1911. Collected for and presented by A. B. Percival. This is the authors selected male syntype. The selected female syntype is also in the collection.	0
borealis—Troglodytes borealis Fischer, 1861	8

borneana—Chalcoparia singalensis borneana Kloss, 1921 J. Fed. Malay States Mus., 10: 209. Adult male. Reg. no. 1927. 4.18.1006. Bukar, Samarahan, Sarawak, Borneo, 26 Oct. 1919. Collected for the Federated Malay States Museum. Presented by H. C. Robinson. The described female syntype is also in the collection. borneanum—Dicaeum minullum borneanum Lönnberg, 1925. Sarawak Mus. J., 3:2. HOLOTYPE, Adult male. Reg. no. 1925. 9.10.3. Mount Murud, Rumah Tamabo Perak, North Sarawak. Borneo, Dec. 1923. Collected by E. Mjöberg. Presented by the Sarawak Museum. borneensis—Chlorura borneensis Sharpe, 1889 60 Ann. Mag. nat. Hist., (6) 3:424. SYNTYPE, Adult male. Reg. no. 1888.12.27.1. Kina Balu, 8,000', North-west Borneo, 12 Feb. 1888. Collected by and purchased of J. Whitehead. The two other syntypes are also in the collection. borneensis—Eupetes macrocercus borneensis Robinson and Kloss, 4.18.987. Samarahan, south Sarawak, 25 Nov. 1919. Collected by a Federated Malay States Museum collector and presented by H. C. Robinson. borneensis—Mirafra borneensis Swinhoe, 1871 . 17 Ann. Mag. nat. Hist., (4) 7:257. HOLOTYPE, Adult. Reg. no. 1873.5.12.800. Banjermassing, south Borneo. Collected by J. Motley (not Wallace as stated in text). Wallace Collection. Reg. no. 1855.2.10.486. Orizaba, southern Mexico. Collected by M. Botteri. Sclater Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 711. There are several other syntypes in the collection. boucardi—Formicivora boucardi P. L. Scalter, 1858 Proc. zool. Soc. Lond., 1858: 300. SYNTYPE, Adult male. Reg. no. 1858.9.27.12. Acatepec, 10,000', Oaxaco State, Mexico, April 1857. Collected by A. Boucard and purchased of A. Sallé. There are two other syntypes in the collection; only these two were listed as types by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 255. boucardi—Zonotrichia boucardi P. L. Sclater, 1867 49 Proc. zool. Soc. Lond., 1867: 1, pl. 1. SYNTYPE, Adult. Reg. no. 1885.2.10.488. La Peubla, Mexico. Collected by A Boucard. Sclater Collection.

This is the figured specimen, and was listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 714. There are several other syntypes in the collection.

- bourdilloni—Alcippe bourdilloni Hume, 1876 . 32B Stray Feathers, 4:485. HOLOTYPE, Adult male. Reg. no. 1886. 10.1.6531. Mynall, Travancore, India, 9 Feb. 1875. Collected by F. Bourdillon. Hume Collection.
- bourdilloni—Merula bourdilloni Seebohm, 1881 . Cat. Bds. Brit. Mus., 5:251, pl. 15. HOLOTYPE, Adult male. Reg. no. 1880.8.19.202. Travancore (Colathorpolay, Patnas, on label), 17 April 1880. Collected by and purchased of F. Bourdillon.
- bouruensis—Rhipidura bouruensis Wallace, 1863 . 37A Proc. zool. Soc. Lond., 1863: 29. SYNTYPE, Adult female. Reg. no. 1873.5.12.461. Bouru Island, Molucca Group, 1861. Collected by A. R. Wallace. Wallace Collection.

The author marked the label of this specimen in red, but there are other syntypes in the collection.

bouruensis—Tropidorhynchus bouruensis Wallace, 1863 Proc. zool. Soc. Lond., 1863: 19, 31. SYNTYPE, Adult (female, vide Gadow, Cat. Bds. Brit. Mus., 9 (1884): 276). Reg. no. 1873. 5.12.1021. Bouru Island, Molucca Group, 1861. Collected by A. R. Wallace. Wallace Collection.

Listed as the type by Gadow, op. cit. There is another syntype in the collection.

46 lected by Petit. Shelley Collection.

The female syntype cannot be traced, although it was received and registered with the male.

brachyptera—Drymoeca brachyptera Sharpe, 1870 Volta River, Gold Coast Colony. Collected by Governor Ussher. Sharpe Collection.

There is another syntype in the collection.

brachypterus—Turdus brachypterus Latham, 1801 Index Ornithologicus, Suppl.: xliii. ?HOLOTYPE, Water-colour drawing. New South Wales. No. 154 in Watling, Original water-colour drawings . . . Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus.; 2 (1906): 137.

Vol. 2, no. 32 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also

Mathews, Birds of Australia 10 (1923): 151.

brachypus—Fregilus graculus var. brachypus Swinhoe, 1871 . Proc. zool. Soc. Lond., 1871: 383. SYNTYPE, Adult male. Reg. no. 1897.6.1.387. China (Shaching, north of Pekin, 21 Sept. 1868, on label). Seebohm Collection.

There are other syntypes in the collection.

brachytarsus—Empidonax brachytarsus P. L. Sclater, 1859 Ibis, (1) 1:441. SYNTYPE, Adult (female, vide Sclater, Cat. Bds, Brit. Mus., 14 (1888):241), Reg. no. 1888.1.13.1030. Cordova, Mexico. Collected by A. Sallé. Sclater Collection. The second syntype is also in the collection.
brachyurus—Tyranneutes brachyurus P. L. Sclater and Salvin, 1881. Ibis, (4) 5: 269. Syntype, Adult male. Reg. no. 1888.1.20.196. Bartica Grove, British Guiana, 31 Dec. 1880. Collected by H. Whitely. Salvin-Godman Collection. This is the authors' selected male. The selected female and another female syntype are also in the collection.
bractiatus—Cinnyris chalybeus bractiatus Vincent, 1933 46 Bull. Brit. Orn. Cl., 53:146. HOLOTYPE, Adult female. Reg. no. 1898.5.2.32. Fort Chiquaqua, Mashonaland, Portuguese East Africa, 30 Aug. 1897. Collected by and purchased of J. L. Sowerby.
bradshawi—Alaemon bradshawi Sharpe, 1904
bradshawi—Cinnyris bradshawi Sharpe, 1898
brauni—Laniarius luehderi brauni Bannerman, 1939 23 Ibis, (14) 3:748, pl. 16. HOLOTYPE, Adult male. Reg. no. 1939. 8.4.1. Quicolungo, Angola, 11 April 1939 (8°27' S., 15°17' E., fide B. P. Hall, MS). Collected by R. H. Braun on behalf of the British Museum (Natural History).
brauniana—Chaptia brauniana Swinhoe, 1863
brenchleyi—Rhipidura brenchleyi Sharpe, 1879
brevicauda—Cittocincla brevicauda Ogilvie-Grant, 1899

- brevicauda—Muscicapa brevicauda Ogilvie-Grant, 1907 . . . 37 Bull. Brit. Orn. Cl., 19:107. HOLOTYPE, Adult female. Reg. no. 1907.6.26.206. (Pontheirville, on label), 2,000′, Upper Congo, 21 Feb. 1907. Collected by and purchased of D. Carruthers.
- brevicaudata—Hirundo brevicaudata McClelland, 1840 18
 in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840): 156. HoloTYPE, Adult. Reg. no. 1879.11.28.447. Assam. Collected by J.
 McClelland and presented by the Honourable East India Company
 Museum.
- brevicaudus—Lamprotornis brevicaudus Sharpe, 1897 62

 Bull. Brit. orn. Cl., 6:48. HOLOTYPE, Adult male. Reg. no. 1897.10.31.3. Elgeyu (3,700', Rift Valley, Kenya, on label), 15

 Aug. 1896. Collected and presented by F. J. Jackson.
- brevipes—Petrocincla brevipes Waterhouse, 1838 31 in J. E. Alexander, An Expedition of Discovery into the Interior of Africa, 2:263. SYNTYPE, Adult. Reg. no. 1838.4.16.134. South of Tans Mountain (now Gamsberg), Damaraland, 23°30′ S., 16°20′ E., 19 May 1837. Collected by J. E. Alexander and purchased through S. Stevens.
- brevirostris—Bathyrhynchus brevirostris McClelland, 1838 . . . 33
 India Review, 2:513. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4383. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.

H. J. Ross but this cannot be confirmed.) Gould Collection.

Bonaparte used a Gould MS name. An early label is marked "for descr. of this bird see Bonap. Review of Gray's Gen. and sub. gen. in B. M. 4 to paper p. 9". (The latter is a reprint of the above C. R. Acad. Sci. Paris.)

brevirostris—Nectarinia 1873			Blanford,
Ibis, (3) 3:86. SYNT Baluchistan, (Bampur, by W. T. Blanford. Re Calcutta.	YPE, Adult male. 2000', 5 April 18	72, on label).	1.11.23.50. Collected
The first British Muslater deleted. A female			
brevirostris—Pedilorhynch Bull. Brit. Orn. Cl., 25 1911.4.27.7. Assobam 1908. Collected and p	: 28. HOLOTYPE, , 2,000', Bumba R	Adult female. Liver, Cameroc	Reg. no.
brevirostris—Pomatorhinus 1919 . Ibis, (11) 1:578. Syn 277. Trang Bom, Co presented by C. B. Klos This is the authors' so is also in the collection		e. Reg. no. 1 ne 1918. Coll	32B 919.12.20. ected and
brevivexilla—Urocissa bre Proc. zool. Soc. Lond., 1897.6.1.221. Western Seebohm Collection.	, 1873 : 688. Hou	LOTYPE, Adult.	Reg. no. y Carles.
bridgesi—Thamnophilus ba Proc. zool. Soc. Lond., mount). Reg. no. 185 lected by T. Bridges an	1856 : 141. Holo 56.9.4.11. David,	TYPE, Adult ma Chiriqui, Verag	4 le (relaxed gua. Col-
bridgesi—Upucerthia bridge Proc. zool. Soc. Lond., Reg. no. 1846.9.9.63. Bridges. There is another synt	1889 : 32. SYNTY Bolivia. Collecte	PE, Adult (relaxed by and purch	
bridgesii—Nasica bridgesi in Jardine, Contribut SYNTYPE, Adult. Res Argentine vide Hellma (1925): 349). Collecte chased of E. Gerrard. Listed as type by P. : 157.	ions to Ornithologe, no. 1881.2.18. hyr, Catalogue of ed by T. Bridges.	.202. Bolivia Birds of the A Eyton Collec	(incorrect, mericas, 4 tion; pur-
britannica—Sitta europaed Novit. Zool., 7:526. 10.15.6. Tring Park, Collected by E. Harter Natural History, from	HOLOTYPE, Adult Hertfordshire, l rt. Presented by	: male. Reg. England, 13 C the American M	Oct. 1900.

britannicus—Acanthis carduelis britannicus Hartert, 1903 59 Vögel der Paläarktishen Fauna, 1:68. HOLOTYPE, Adult male. Reg. no. 1936.10.15.14. Rottingdean, Sussex, England, April 1902. Collected for Brazenor, Naturalists, of Brighton. Presented by the American Museum of Natural History, New York; from the Rothschild Collection.
brookei—Eurylaimus javanicus brookei Robinson and Kloss, 1919 Bull. Br. Orn. Cl., 40:15. Syntype, Adult male. Reg. no. 1919. 11.5.15. Ulu Paku, Saribas, southwest Sarawak, 25 Nov. 1916. Collected by a native for H. C. Robinson and C. B. Kloss and presented by the Federated Malay States Museum. Kuala Lumpur. The female syntype is in the collection.
brooki—Automolus brooki Chubb, 1916
brooksi—Phylloscopus brooksi Hume, 1874
brownii—Malurus brownii Vigors and Horsfield, 1827 36 Trans. Linn. Soc. Lond., 15:223. Holotype, Adult male (relaxed mount). Reg. no. 1863.7.7.23. Thirsty Sound, South Queensland, Australia, Sept. 1802. Collected by J. Brown. Presented as the type by the Linnean Society of London. The specimen has now lost the tail.
brucei—Alcippe brucei Hume, 1870
brunnea—Alcippe brunnea Gould, 1862
brunnea—Larvivora brunnea Hodgson, 1837

The specimen is labelled as the type, and listed as the type by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 213, who has confused the locality with that of the second syntype, a female from Salwatty,

which is also in the collection.

A misprint stated the tail to be only $\frac{5}{8}$ " long.

- brunneiceps—Brachypteryx brunneiceps Ogilvie-Grant, 1896 . . . 31

 Ibis, (7) 2:547. Syntype, Adult female. Reg. no. 1897.5.13.430.

 Canloon Volcano, Negros, Philippines, 14 April 1896. Collected by J. Whitehead and presented by the subscribers to his expedition.

A male syntype marked "type of male" by the author, is in the American Museum of Natural History, vide Hartert, Novit Zool.,

27 (1920): 479.

brunneiceps—Hylophilus brunneiceps P. L. Sclater, 1866 . . . 56

Proc. zool. Soc. Lond., 1866: 322. HOLOTYPE, Adult male. Reg.
no. 1886.9.15.311. (Rio Vaupé, Upper Rio Negro, north-west
Brazil, vide Sclater, Ibis, (4) 5 (1881): 306), (20 July 1831, on label).
Collected by J. Natterer. Sclater Collection; received by him in
exchange with the Vienna Museum.

brunneiceps—Lagonosticta brunneiceps Sharpe, 1890 60
Cat. Bds. Brit. Mus., 13:277. Syntype, Adult male. Reg. no. 1877.7.11.350. Maragaz, Eritrea, 27 July. Collected by W. Jesse on the Zoological Society Abyssinian Expedition. Sharpe Collection.

This specimen was labelled as the type by the author and listed as such by Shelley, *Birds of Africa* 4 (1905): 258, but there are several other syntypes in the collection.

- brunneiceps—Munia brunneiceps Walden, 1872 . . 60 Trans. zool. Soc. Lond., 8:73, pl. 9. HOLOTYPE, Adult male. Reg. no. 1873.5.12.114. Makassar, Celebes, (1856, on label). Collected by A. R. Wallace. Wallace Collection.
- brunneiceps—Yuhina brunneiceps Ogilvie-Grant, 1906 32

 Bull. Br. Orn. Cl., 16:121. SYNTYPE, Adult male. Reg. no. 1907.12.12.73. Racu Racu, 6,000', Mount Morrison, central Formosa, Feb. 1906. Collected by and purchased of W. Goodfellow.

There are two other syntypes in the collection.

- brunneicollis—Troglodytes brunneicollis P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858: 297. SYNTYPE, Adult male. Reg. no. 1858.9.27.2. Parada, 10,000', Oaxaca, Mexico, Dec. 1857. Collected by A. Boucard for A. Sallé and purchased of the latter. There is another syntype in the collection.
- 1878.11.12.7. Aru Islands, (1866, on label). Collected for Von Rosenberg. Presented by Leiden Museum. Although only one specimen came to the collection the author

evidently saw others.

- brunneiventris—Chlorura brunneiventris Ogilvie-Grant, 1894 . Bull. Brit. Orn. Cl., 3:50. SYNTYPE, Adult (male, on label). Reg. no. 1897.5.13.52. North-west Luzon, (La Trinadad, Benguet district, 18 Feb. 1894, on label). Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. A female syntype is also in the collection.
- brunnescens—Aplonis brunnescens Sharpe, 1890 . Cat. Bds. Brit. Mus., 13:132. SYNTYPE, Adult. Reg. no. 1878. 1.19.36. Savage Island, central Polynesia. Collected by the Rev. S. J. Whitmee.

There are two other syntypes in the collection.

- brunnescens—Cyphorhinus brunnescens Sharpe, 1881 Cat. Bds. Brit. Mus., 6:293. HOLOTYPE, Adult male (immature, vide Hellmayr, Catalogue of Birds of the Americas, 7 (1934): 292). Reg. no. 1885.3.6.166. Remedios, Cauca Valley, Antioquia. Collected by T. K. Salmon. Salvin-Godman Collection.
- brunnescens—Herpornis brunnescens Sharpe, 1876 Ibis, (3) 6:41. HOLOTYPE, Adult male. Reg. no. 1873.5.12.955. Sarawak, Borneo, 1854. Collected by A. R. Wallace. Wallace Collection.
- brunnescens—Horeites brunnescens Hume, 1872 lbis, (3) 2:109. HOLOTYPE, Adult. Reg. no. 1886.7.8.2185. Darjeeling, India. Collected by Gammie. Hume Collection.

- brunnescens—Lalage nigra brunnescens E. C. S. Baker, 1923 Bull. Br. Orn. Cl., 44:13. HOLOTYPE, Adult female. Reg. no. 1886.4.1.941. Klang, Selangor, Malay Peninsula, 11 Oct. 1879. Collected by W. Davison. Hume Collection.
- brunnescens—Margarornis brunnescens P. L. Sclater, 1856 Proc. zool. Soc. Lond., 1856: 27, pl. 116. HOLOTYPE, Adult. Reg. no. 1889.5.20.450. Bogota, Colombia. Sclater Collection: obtained by Sclater from Verreaux.
- brunnescens—Pipilo torquatus brunnescens Van Rossem, 1938 Bull. Brit. Orn. Cl., 58:131. HOLOTYPE, Adult male. Reg. no. 1899,2,1,3845. Totontepec, Oaxaca, Mexico, Feb. 1889. Collected by M. Truiillo, Salvin-Godman Collection.
- brunneus—Dasyornis? brunneus Gould, 1837. Proc. zool. Soc. Lond., 1837: 150. HOLOTYPE, (Immature). Reg. no. 1854.12.20.26a. Australia (South America, vide P. L. Sclater, Cat. Bds. Brit. Mus., 15 (1890): 44). Purchased of J. Gould.
- brunneus—Dromaeocercus brunneus Sharpe, 1877 Proc. zool. Soc. Lond., 1877: 23, pl. 2. HOLOTYPE, Adult. Reg. no. 1876.11.1.1. Near Antananarivo, Madagascar. Collected by A. Kingdon. Presented by A. Peckover who received it from the collector.
- brunneus— Sclerurus brunneus P. L. Sclater, 1857. Proc. zool. Soc. Lond., 1857: 17. HOLOTYPE, Adult. Reg. no. 1889.5.20.426. Bogota, Colombia, 1856. Collected by Bell. Sclater Collection.
- brunnifrons—Orthotomus (Prinia) brunnifrons Hodgson, 1845 Proc. zool. Soc. Lond., 1845: 29. There is no evidence that Hodgson's specimens in the collection can be considered as types or syntypes although listed as such by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 145.
- buaensis—Ptilotis procerior buaensis Wiglesworth, 1891. Abh. Ber. k. zool. anthrop. - ethn. Mus. Dresden, 6:34. SYNTYPE. Adult (female, on label). Reg. no. 1878.1.19.33. Bua, Vanua Levu, Fijis. Collected by E. L. Layard. Presented, possibly by the Rev. J. Whitmore.

Wiglesworth used a Layard MS. name. This is the syntype labelled 'P. buenis' referred to in the description, but the measurements are smaller than those given.

buchanani—Prinia buchanani Blyth, 1844 J. Asiat. Soc., Bengal, 13:376. New name for P. rufifrons Jerdon, 1840 (q.v.), on the grounds of its preoccupation by P. rufifrons Ruppell, 1835. Sclater, *Ibis*, (6) 4 (1892): 79, stated that the type of P. buchanani is in Calcutta but since it is only a new name the original syntypical specimens are valid.

buchenorum—Cinnvris sovimanga buchenorum Williams, 1953. Ibis. 95: 502, pl. 5. HOLOTYPE, Adult male. Reg. no. 1952.50.1. Menai Island, Cosmoledo Islands, Aldabra Archipelago, Indian Ocean, 15 April 1952. Collected and presented by D. Vesey-Fitzgerald. bucklevi—Calandrella bucklevi Shellev, 1873 Ibis, (3) 3:142. SYNTYPE, Adult. Reg. no. 1889.9.13.139. Accra, Gold Coast. Collected by T. E. Buckley. Shelley Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 599. bucklevi—Ptilochloris bucklevi P. L. Sclater and Salvin, 1880 Proc. zool. Soc. Lond., 1880: 158, pl. 16. SYNTYPE, Adult female. Reg. no. 1888.1.20.336. Pindo, eastern Ecuador. Collected by C. Buckley. Salvin-Godman Collection. The immature syntype is also in the collection. buckleyi—Tityra buckleyi Salvin and Godman, 1890 Biol. Centr.-Amer., Aves, 2:120. HOLOTYPE, Adult male. Reg. no. 1888.1.20.403. Yanayacu, eastern Ecuador (1877, on label). Collected by C. Buckley. Salvin-Godman Collection. buddae—Turdus merula buddae R. and A. Meinertzhagen, 1926 . 31 Bull. Br. Orn. Cl., 46:98. HOLOTYPE, Adult male. Reg. no. 1927.1.10.171. Chu Nallah, 14,000', Gyangtse, southern Tibet, 15 Dec. 1925. Collected and presented by F. Ludlow. buensis—Myiolestes buensis Layard, 1876 Ibis, (3) 6:145. There are several specimens from Bua Island, off Vanua Levu, Fiji Islands, in the collection and Mayr, Amer. Mus. Novit., 628 (1933): 9 regarded all such specimens as syntypes. The above specimens were not collected until late April 1876 and since the description appeared in the April 1876 number of the *Ibis* it seems improbable that they were available at the time. buffoni-Polioptila buffoni P. L. Sclater, 1861 Proc. zool. Soc. Lond., 1861: 127. SYNTYPE, Adult male. Reg. no. 1884.5.15.213. Cavenne. Collected for Verreaux. Sclater Collection. Listed as the type by Hellmayr, Catalogue of birds of the Americas 7 (1934): 496. A second syntype is in the collection. bulgeri—Rhipidura bulgeri Layard, 1877 Ibis, (4) 1:361. HOLOTYPE, Adult male. Reg. no. 1893.7.8.5. Ansevata, New Caledonia, 2 Feb. 1877. Collected by E. L. Layard. Received from P. L. Sclater in exchange. bulliens—Cisticola bulliens Lynes, 1930 35 Ibis, (12) 6; supplement: 315, pl. 11. HOLOTYPE, Adult male. Reg. no. 1928.10.25.2. Lobito Bay, Angola, 4 March 1928. Collected by Admiral H. Lynes and B. B. Osmaston. Presented by Admiral

H. Lynes.

bulomachus—Suthora bulomachus Swinhoe, 1866
bulubulu—Cisticola textrix bulubulu Lynes, 1931
bungurense—Piprisoma agile bungurense Chasen, 1934
burmae—Tchitrea affinis burmae Salomonsen, 1933 38 Ibis, (13) 3:736. HOLOTYPE, Adult male. Reg. no. 1909.4.7.84. Kani, Lower Chindwin, Burma, 9 May 1905. Collected and purchased by A. Mears.
burmanica—Estrelda burmanica Hume, 1876
burmanica—Leucocerca burmanica Hume, 1881
burmanica—Prinia inornata burmanica Harington, 1913 35 Bull. Brit. Orn. Cl., 31:111. SYNTYPE, Adult male. Reg. no. 1913.6.14.2. Mandalay, Upper Burma, 27 June 1911. Collected and presented by H. H. Harington. The female syntype is also in the collection.
burmanica—Saxicola caprata burmanica E. C. S. Baker, 1923 31 Bull. Br. Orn. Cl., 43:19. HOLOTYPE, Adult male. Reg. no. 1886.4.1.1492. Pegu, Burma. Collected by E. W. Oates. Hume Collection.
burmanicus—Criniger burmanicus Oates, 1889
burmanicus—Phylloscopus burmanicus Brooks, 1880

9.1.879. Pegu, Burma, 15 Nov. 1877. Collected by E. W. Oates. Seebohm Collection. One wing of the specimen is missing. The second syntype is also in the collection.
cat. Bds. Brit. Mus., 6:125. SYNTYPE, Adult male. Reg. no. 1863.5.15.83. Pegu, 1862. Collected and presented by W. T. Blanford. There are several other syntypes in the collection.
Ibis, (1) 4:21. Holotype, Adult (female, vide Sharpe, Cat Bds. Brit. Mus., 13 (1890):46). Reg. no. 1862.1.17.16. Near Thayetmya, Upper Burma. Collected and presented by T. C. Jerdon.
Ann. Mag. nat. Hist., (3) 10:445. HOLOTYPE, Adult. Reg. no. 1862.12.1.17. Cameroon Mountains, 7,000', West Africa. Collected by R. Burton. Presented by Mrs Burton.
Ann. Mus. civ. Stor. nat. Giacomo Doria, 12:341. HOLOTYPE, Adult female. Reg. no. 1873.5.12.910 Bouru Island, South-west Pacific, 1861. Collected by A. R. Wallace. Wallace Collection.
Buryi—Parisoma buryi Ogilvie-Grant, 1913
Bull. Brit. Orn. Cl., 13: 22. HOLOTYPE, Adult female. Reg. no. 1903.8.12.20. Dthubiyat, Upper Haushabi, Southern Arabia, (26 Aug. on label). Collected by and purchased of G. W. Bury.
Outleri—Anomalospiza butleri W. L. Sclater and Praed, 1918 61 Ibis, (10) 6:460. HOLOTYPE, Adult male. Reg. no. 1915.12.24. 1781. Kajo Kaji, Lado Enclave, Sudan, 8 April 1915. Collected and presented by A. L. Butler.
outleri—Cinnyris osea butleri W. L. Sclater and Mackworth-Praed, 1918
Bull. Brit. Orn. Cl., 21:17. HOLOTYPE, Adult male? Reg. no. 1916.9.20.1157. Chak Chak, Bahr-el-Gazal Province, Sudan, (17 Feb. 1907, on label). Collected and presented by A. L. Butler.

- butleri—Lagonosticta butleri Ogilvie-Grant, 1907 60

 Bull. Brit. Orn. Cl., 21:16. SYNTYPE, Female (immature, vide,
 Sclater and Mackworth-Praed, Ibis, (10) 6 (1918):446). Reg.
 no. 1907.11.21.2. Chak-Chak, Bahr-El-Gazal, Sudan, (17 Feb.
 Feb. 1907, on label). Collected and presented by A. L. Butler.
 The second syntype is also in the collection.

Nectarinia byronensis Bloxham, 1826, is a nomen nudum. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 9. Although the text refers to "a specimen" there would appear to be several syntypes.

cabanisi—Pyrgisoma cabanisi P. L. Sclater and Salvin, 1868 . . . 49
Proc. zool. Soc. Lond., 1868 : 324. SYNTYPE, Adult. Reg. no. 1885.2.10.590. San José, Costa Rica. Collected by Hoffman. Sclater Collection.

Listed as the type by Hellmayr and Conover, Catalogue of birds of the Americas, 11 (1938): 472. The specimen collected by J. Carmiol and listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 735, is not a syntype.

Ibis, (12) 4:433. HOLOTYPE, Adult male. Reg. no. 1928,10.7.1. Cercedilla, 5,000', Sierra de Guadarrama, Spain, 14 Oct. 1922. Collected and presented by H. F. Witherby. cacabata—Muscicapa sibirica cacabata. See appendix, Corrections and Additions, vol. 2 . . . cacharensis—Accentor cacharensis Hodgson, 1845 30 Proc. zool. Soc. Lond., 1845: 34. ?SYNTYPE, Adult. Reg. no. 1845.1.9.696. Nepal. Collected and presented by B. H. Hodgson. There are several other specimens in the collection which may be syntypes. cacharensis—Neornis cacharensis Hodgson, 1845 in Blyth, J. Asiat. Soc. Bengal, 14:590. ?HOLOTYPE, Adult. Reg. no. 1845.1.9.473. Nepal. Collected and presented by B. H. Hodgson. The specimen was labelled as the type in an unknown hand. Listed as the type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 138. caerebicolor—Dacnis caerebicolor P. L. Sclater, 1851 in Jardine, Contributions to Ornithology: 106. HOLOTYPE, Adult male. Reg. no. 1884.7.31.85. (Bogota, on label), Colombia, 1851. Collected for Argent. Sclater Collection. caerulatus—Cinclosoma caerulatus Hodgson, 1836 Asiat. Res., 19: 147. Syntype, Adult (relaxed mount). Reg. no. 1843.1.13.570. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection. caerulea—Myiothera caerulea Raffles, 1822. Trans. Linn. Soc. Lond., 13:301. SYNTYPE, Adult male (relaxed mount). Reg. no. 1880.1.1.4511. Sumatra. Collected by Sir T. S. Raffles. Presented by the Honourable East India Company Museum. A female syntype from the Museum of the Zoological Society of London is also in the collection. caerulea—Urocissa caerulea Gould, 1862. . Proc. zool. Soc. Lond., 1862: 282. SYNTYPE, Adult. Reg. no. 1863.2.16.3. Northern Formosa, April 1862. Collected by R. Swinhoe. Purchased of S. Stevens. Swinhoe Collection. There is a second syntype in the collection. caeruleocephala—Muscicapa caeruleocephala Sykes, 1832 Proc. zool. Soc. Lond., 1832:85. SYNTYPE, Adult. (female, vide, Horsfield and Moore, Catalogue of birds in the museum of the East India Company (1854): 138) (relaxed mount). Reg. no. 1880.1.1. 2079. Deccan, India. Collected by W. H. Sykes. Presented by the Honourable East India Company Museum.

cabrerae—Parus ater cabrarae Witherby, 1928

Renamed Hypothymis azurea sykesi E. C. S. Baker, 1920, on the grounds of preoccupation by M. caeruleocephala Scopoli, 1786. There is another syntype in the collection.

- caeruleogrisea—Campephaga caeruleogrisea G. R. Gray, 1858 . 20

 Proc. zool. Soc. Lond., 1858: 179. HOLOTYPE, Adult female.

 Reg. no. 1873.5.12.1821. Aru Islands, 1857. Collected by A. R. Wallace. Wallace Collection.
- caerulescens—Butalis caerulescens Hartlaub, 1865 37

 Ibis, (2) 1:267. SYNTYPE, Adult male. Reg. no. 1868.2.21.1.

 Natal. Collected by T. Ayres. Presented by J. H. Gurney.

 Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 4 (1879):155, but a female, in the possession of Gurney, was also referred to in the description.
- caerulescens—Diglossopis caerulescens P. L. Sclater, 1856 51 Ann. Mag. Nat. Hist., (2) 17:467. The author indicates that the description is based on specimens seen in Bremen and Paris. The Sallé specimen listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 12, has no type status.

A specimen in the collection was listed as type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 354, but it has no status.

caerulirostris—Chrysomitridops caerulirostris S. B. Wilson, 1890 . 55 Proc. zool. Soc. Lond., 1889 (publ. 1890) : 445. Syntype, Adult male. Reg. no. 1898.10.20.22. Kauai, (Waimea, 3,000 ft., vide Wilson and Evans, Aves Hawaiiensis, (1890) : 59), Sandwich Islands, Sept. (error for Oct. ?) 1888. Collected and presented by S. B. Wilson.

Wilson states that his first specimens were collected in October 1888, but this and a female specimen are labelled "sp. n.", and together with three specimens in the University Museum of Zoology, Cambridge, were collected in September 1888 according to the specimen labels. They would appear to be syntypes. Wilson also states that some specimens preserved in spirit were lost during his journey to England.

caesar—Poospiza caesar P. L. Sclater and Salvin., 1869 . . . 49
Proc. zool. Soc. Lond., 1869: 152, pl. 13. SYNTYPE, Adult. Reg.
no. 1885.2.10.336. Tinta, 11,000′, South Peru, 20 May 1868.
Collected by H. Whitely. Sclater Collection.

The second syntype is also in the collection. The specimen listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 736, was collected later and has no type status.

cajabambae—Leptasthenura striata cajabambae Chapman, 1921 . 3

Amer. Mus. Novit., : 18:9. HOLOTYPE, Adult female. Reg. no. 1899.6.1.81. 9,500′, Cajabamba, Peru. 28 Mar. 1894. Collected by O. T. Baron. Salvin-Godman Collection.

aledonicus—Corvus caledonicus Gmelin, 1787
Systema Naturae, 1, pt. 1:367. HOLOTYPE, Unsigned painting.
North Caledonia, 7 Sept. 1774. No. 53 in Forster, Unpublished
water-colour drawings made during Cook's second voyage.
Library of the Zoology Department, British Museum (Natural
History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser.,
1, (1959): 285.

- caledoniensis—Garrulus glandarius caledoniensis Hazelwood and Gorton, 1953 1953.71.1. Pitlochry, Perthshire, 25 Sept. 1951. Presented by the Bolton Public Museum.
- callainus—Malurus callainus Gould, 1867 . Proc. zool. Soc. Lond., 1867: 302. HOLOTYPE, Adult male. Reg. no. 1881.5.1.674. South Australia (Salt Bush Scrub, 300-400 miles north-west of Adelaide, vide Gould, Birds of Australia, Supplement. (1867): pl. 23). Collected by S. White. Gould Collection.
- callensis—Emberiza cia callensis Ticehurst and Whistler, 1938 Ibis, (14) 2:727. HOLOTYPE, Adult male. Reg. no. 1931.1.1.5865. Bom Jesus, Braga, Portugal, 31 May 1920. Collected by H. F. Witherby. H. F. Witherby Collection.
- callinota—Formicivora callinota P. L. Sclater, 1855 Proc. zool. Soc. Lond., 1855: 89, pl. 96. HOLOTYPE, Adult male. Reg. no. 1854.4.6.12. Bogota, Colombia. Purchased of Argent.
- calophrys—Chlorospingus calophrys P. L. Sclater and Salvin, 1876 51 Proc. zool. Soc. Lond., 1876: 354. HOLOTYPE, Adult. Reg. no. 1885.6.8.43. Tilotilo, Yungas Province, Bolivia, 1876. Collected by C. Buckley. Salvin-Godman Collection.
- caloptera—Formicivora caloptera P. L. Sclater, 1859. Proc. zool. Soc. Lond., 1859: 142. SYNTYPE, Adult. Reg. no. 1888.1.13.98. Pallatanga, Ecuador, Sept. or Oct. 1858. Collected by L. Fraser. Sclater Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 29. There are two other syntypes in the collection.
- calopterum—Todirostrum calopterum P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1857: 82, pl. 125. HOLOTYPE, Adult. Reg. no. 1886.6.24.360. Upper Napo River, Ecuador. Collected by Don Villavicencio for Prof. W. Jameson. Jardine Collection.
- camaronensis—Geocichla camaronensis Sharpe, 1905 Ibis, (8) 5:472. HOLOTYPE, Adult male. Reg. no. 1905.7.30.5. Efulen, Cameroon, West Africa, 5 Nov. 1903. Collected by G. L. Bates. Bates Collection.

- camaroonensis—Anthus rufulus camaroonensis Shelley, 1900 . . . 19 The Birds of Africa, 2:320. HOLOTYPE, Adult male. Reg. no. 1887.3.7.16. Camaroons Mt., 10,000', 8 Oct. 1886. Collected by H. H. Johnston and presented by the Cameroons Committee of the British Association.
- camarunensis—Smithornis camarunensis Sharpe, 1905 Ibis, (8) 5:469. SYNTYPE, Adult female. Reg. no. 1905.1.24.99. River Ja, Camaroon, West Africa, 17 May 1904. Collected by G. L. Bates and purchased of E. Gerrard.
- cambodiana—Aegithina tiphia cambodiana Hall, 1957 Ibis, 99:150. HOLOTYPE, Adult male. Reg. no. 1928.6.26.1485. Siem-Reap, Cambodia, 16 Dec. 1927. Collected by J. Delacour on the Fourth Franco-British Expedition to Indo-China.
- cambodiana—Aethovyga saturata cambodiana Delacour, 1948 46 J. Washington Acad. Sci., 38:55. HOLOTYPE, Adult male. Reg. no. 1928.6.26.1578. Bokor, Cambodia, 12 Dec. 1927. Collected by J. Delacour and W. P. Lowe, from the 4th Franco-British Expedition to Indo-China.
- cambodiana—Notodela cambodiana Delacour and Jabouille, 1928 . 31 Bull. Br. Orn. Cl., 48:132. HOLOTYPE, Adult male (tip of bill broken off). Reg. no. 1928.6.26.10. Bokor, 1,000 metres, south Cambodia, 13 Dec. 1927. Collected by J. Delacour and M. P. Jabouille on the Fourth Franco-British Expedition to French Indo-China and presented by J. Spedan Lewis.
- cambodianum—Dicaeum beccarii cambodianum Delacour and Jabouille, 1928. 45 Dec. 1927. Collected by J. Delacour and M. P. Jabouille, from the Fourth Expedition to Indo-China.
- cambodianus—Poliopsar cambodianus Sharpe, 1888 Ibis, (5) 6:477. HOLOTYPE, Adult. Reg. no. 1860.11.9.18. Cambodia. Collected by M. Mouhot. Purchased of Stevens.
- camburni—Hyphantornis camburni Sharpe, 1890 . . 61 Bull. Br. Orn. Cl., 10:35. HOLOTYPE, Adult male (female in text). Reg. no. 1900.1.19.17. Mount Kenya, 5,000', Nairobi Forest (17 July 1899, on label). Collected by C. Camburn (vide Shelley, The Birds of Africa, 4 (1905): 486). H. J. Mackinder Expedition Collection.
- cameroonensis—Anthreptes fraseri cameroonensis Bannerman, 1921 46 Bull. Brit. Orn. Cl., 41:137. HOLOTYPE, Adult male. Reg. no. 1908.5.25.91. Ja River, Cameroon, 18 Feb. 1906. Collected by and purchased of G. L. Bates.

cameroonensis—Dryoscopus angolensis cameroonensis Bannerman, queathed by Boyd Alexander. camerunensis—Bradypterus camerunensis Boyd Alexander, 1909 . 35 Bull. Brit. Orn. Cl., 25:19. HOLOTYPE, Adult male. Reg. no. 1911.12.23.4184. Peak of Cameroon Mountain, 7,000', 5 April 1909. Collected by W. Boyd Alexander. Boyd Alexander Bequest. camerunensis—Cryptolopha camerunensis Boyd Alexander, 1909 . 35 Bull. Brit. Orn. Cl., 25:13. HOLOTYPE, Adult male. Reg. no. 1911.12.23.4118. Peak of Cameroon, West Africa, 23 April, 1909. Collected by W. Boyd Alexander. Boyd Alexander Bequest. campbelli—Turdus campbelli Swinhoe, 1873 31 Ann. Mag. nat. Hist., (4) 12:374. HOLOTYPE, Adult male. Reg. no. 1896.6.1.837. Lighthouse Island, Chefoo, north China, 15 May 1873. Collected by Campbell, the lighthouse keeper, for Swinhoe. Seebohm Collection. The date of collection on the label, 1874, is incorrect: Swinhoe was in Chefoo in 1873 (vide Ibis, (4) 2 (1878): 127). He renamed this type in 1874 as Turdus chyropleurus (q.v.). cancrominus—Platyrhynchus cancrominus P. L. Sclater and Salvin, Collected by O. Salvin. Salvin-Godman Collection.
There are several other syntypes in the collection. candida—Nigrita canicapilla candida Moreau, 1942 60 Bull. Brit. Orn. Cl., 62:43. HOLOTYPE, Adult male. Reg. no. 1944.8.1.5. Ujamba Forest, 6,700', Kungwe Mountains, Kigoma District, Tanganyika, 16 Aug. 1941. Collected by Salemin Asumani for R. E. Moreau. Presented by R. E. Moreau. . 49 Godman Collection. canescens—Emberiza canescens Swinhoe, 1860 . Ibis, (1) 2:62. SYNTYPE, Adult male. Reg. no. 1898.11.1.1109. Amoy, China, 6 Feb. 1857. Collected by R. Swinhoe. Seebohm

The described female syntype is also in the collection.

Collection.

canescens—Empidonax canescens Salvin and Godman, 1889 . 9 Biol. CentrAmer., Aves 2:79. SYNTYPE, Adult male. Reg. no. 1899.4.20.2404. Mexicalcingo, near City of Mexico, 6 Feb. 1888. Collected by F. Ferrari-Perez. Salvin-Godman Collection. There are several other syntypes in the collection.
canescens—Erpornis xantholeuca canescens Delacour and Jabouille, 1928
canicapilla—Zonotrichia canicapilla Gould, 1839
canicapillus—Aethiops canicapillus Strickland, 1841 60 Proc. zool. Soc. Lond., 1841 : 30. Sharpe, Cat. Bds. Brit. Mus., 12 (1890) : 316. lists specimens as the types but, Salvin, Catalogue of a collection of birds of Strickland (1882) : 243, states that the type is in the Cambridge University Zoological Museum, and it would appear that the British Museum specimens from L. Fraser were collected a year later, vide Fraser, Proc. zool. Soc. Lond., 1842 (publ. Feb. 1843) : 145.
caniceps—Pastor caniceps Blyth, 1843
caniceps—Synallaxis caniceps Eyton, 1856
canipennis—Hemixus canipennis Seebohm, 1890
cantillans—Mirafra cantillans Blyth, 1845

- in the Calcutta Museum by P. L. Sclater, *Ibis*, (6) 4 (1892): 83. Two Jerdon specimens from Madras in the B.M. (N.H.) collection were listed as types by Sharpe, *Cat. Bds. Brit. Mus.*, 13 (1890): 606. *Mirafra cantillans* Blyth, *J. Asiat. Soc. Bengal*, 12 (1843): 181 is a nomen nudum; Blyth used Jerdon's MS name.

The male syntype is also in the collection.

- cantori—Criniger cantori Moore, 1854

 Proc. zoll. Soc. Lond., (May 1854): 279. Also in Horsfield and Moore, Catalogue of the Birds in the Museum of the Honourable East India Company, (Nov. 1854) 1: 410. SYNTYPE, Adult. Reg. no. 1880.1.1.1900. Malacca. Collected by T. Cantor and presented by the Honourable East India Company Museum.

There is what may be a second syntype in the collection.

cantori—Microtarsus cantori Horsfield and Moore, 1854 . . . 2.

Catalogue of the Birds in the Museum of the Honourable East India

Company, 1: 409. HOLOTYPE.

This is also the holotype of Brachypodius tristis Blyth, 1845 (a,v,)

cantoroides—Calornis cantoroides G. R. Gray, 1861 62

Proc. zool. Soc. Lond., 1861: 431. SYNTYPE, Adult. Reg. no. 1861.12.11.69. Mysol. Collected by A. R. Wallace. Wallace Collection.

There are several other syntypes in the collection.

capellanus—Corvus capellanus P. L. Sclater, 1876 7

Proc. zool. Soc. Lond., 1876: 694. pl. 66. SYNTYPE, Adult male.

Reg. no. 1878.12.10.2. Fao, Persian Gulf, Summer 1876. Collected by J. Huntley. Presented to the London Zoo by J. Huntley on 16 June 1876. Died 29 July 1877. Purchased of the Zoological Society.

A second syntype is in the collection.

capensis—Calandrella sclateri capensis Ogilvie-Grant, 1913 . . . 17

Ann. S. Afr. Mus., 13:41. Holotype, Adult male. Reg. no. 1912.11.27.1. Near Philipstown, Cape Province, 18 July 1912. Collected by H. L. Hare and presented by the South African Museum.

capensis—Drymoica capensis Smith, 1843
capitale—Todirostrum capitale P. L. Sclater
capitalis—Mixornis? capitalis Tweeddale, 1877 32B Ann. Mag. nat. Hist., (4) 20:535. Holotype, Adult female. Reg. no. 1888.4.20.1109. Dinagat Island, Philippines, June 1877. Collected by A. H. Everett. Tweeddale Collection.
capitalis—Muscicapa? capitalis McClelland, 1840 20 in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840): 157. Holotype, Adult (imperfect specimen). Reg. no. 1880.1.1.2938. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.
capitalis—Myiobius capitalis Salvin, 1865
capitalis—Thamnophilus capitalis P. L. Sclater, 1858 4 Proc. zool. Soc. Lond., 1858: 65. SYNTYPE, Sub-adult male. Reg. no. 1889.9.20.73. Rio Napo, Ecuador. Probably collected by Villavicencio. Sclater Collection; obtained by him from Verreaux. The female syntype is in the collection.
capito—Eopsaltria capito Gould, 1854
cara—Aethopyga cara Hume 1874
carbonaria—Cercomacra carbonaria P. L. Sclater and Salvin, 1873. Nomencl Av Neotron, : 161. Syntype, Adult male, Reg. no.

Natterer's MS name. carbonarium—Serrirostrum carbonarium Lafresnaye and D'Orbigny, of Verreaux by Sclater was listed by him, Cat. Bds. Brit. Mus. 11 (1886): 7, as "One of the types". It has no original label to verify this. According to Hellmayr, Catalogue of birds of the Americas 8 (1935): 233, the type is in the Paris museum. carneipes—Caereba carneipes P. L. Sclater, 1859. 51 Proc. zool. Soc. Lond., 1859: 376. SYNTYPE, Adult male. Reg. no. 1884.7.31.138. Playa Vicente, Oaxaca, Mexico. (April 1859. Collected by A. Boucard, vide Sclater, Catalogue of a collection of American birds (1862): 54). Sclater Collection. The other syntype is in the collection. carnipes—Coccothraustes? carnipes Hodgson, 1836 Asiat. Res., 19:151. SYNTYPE, Adult female. Reg. no. 1845,1.9. 301. Nepal. Collected by and purchased of B. H. Hodgson. Sharpe Cat. Bds. Brit. Mus., 12 (1888): 49, refers to this specimen, together with another female and an adult male, as the "types" of the species, although it is obvious from the description that Hodgson had not seen an adult male at that time. caroli—Aeithalus caroli Sharpe, 1871. Ibis, (3) 1:415. Syntype, Adult male. Reg. no. 1876.5.23.537. Ovaquenyama, Damaraland, South West Africa, 26 May 1867. Collected by C. J. Anderson. Sharpe Collection. There are two other syntypes in the collection. carruthersi—Alethe carruthersi Ogilvie-Grant, 1906 Bull. Br. Orn. Cl., 19:25. SYNTYPE, Adult male. Reg. no. 1906.12.23.1238. 150 miles west of Entebbe, 5,000', Uganda, 12 Dec. 1905. Collected by D. Carruthers and presented by the subscribers to the Ruwenzori Expedition. There is also a female syntype in the collection. carruthersi—Anthothreptes carruthersi Ogilvie-Grant, 1907 . 46 Bull. Brit. Orn. Cl., 19:106. SYNTYPE, Adult male. Reg. no. 1907.6.26.102. West shore of Lake Tanganyika, 3 Jan. 1907. Collected by and purchased of D. Carruthers. This is the author's selected male, his selected female syntype is also in the collection. carruthersi—Cisticola carruthersi Ogilvie-Grant, 1909 .

Bull. Brit. Orn. Cl., 23:94. HOLOTYPE, Adult female. Reg. no. 1905.12.23.902. Mokia, 3,400', South-east Ruwenzori, Uganda, 17 June 1906. Collected by D. Carruthers. Presented by the sub-

scribers to the British Museum Ruwenzori Expedition.

1889,9.20.431. Rio Brancho, Brazil, 22 Dec. 1831. Collected by

There is a female syntype in the collection. The authors used

J. Natterer. Sclater Collection

our united
carruthersi—Crateropus carruthersi Ogilvie-Grant, 1907 32E Bull. Brit. Orn. Cl., 19: 106. HOLOTYPE, Adult male. Reg. no. 1907.6.26.167. Near Niembe, 2,500′, Upper Congo, 8 Jan. 1907. Collected by and purchased of D. Carruthers.
carruthersi—Parus flavipectus carruthersi Hartert, 1917 42 Bull. Brit. Orn. Cl., 38: 19. HOLOTYPE, Adult male. Reg. no. 1909.10.26.199. Samarkand, 2,000', Turkestan, 25 Jan. 1908. Collected by and purchased of D. Carruthers.
cashmeriensis—Chelidon cashmeriensis Gould, 1858
cashmeriensis—Cinclus cashmeriensis Gould, 1859 27 Proc. zool. Soc. Lond., 1859 : 494. HOLOTYPE, Adult. Reg. no. 1881.5.1.6289. Cashmere (in text; Chimourarie Lake, Ladak, 26 July 1852, on label). Collected by Dr. A. L. Adams. Gould Collection.
cashmirensis—Motacilla cashmirensis Brooks, 1871 19 Proc. Asiat. Soc. Bengal, 1871 : 210. SYNTYPE, Adult. Reg. no. 1887.2.1.1468. Cashmere (Near Ganderbal, on label), 22 May 1871. Collected by W. E. Brooks. Hume Collection.
cashmirensis—Sitta cashmirensis Brooks, 1871
Casius—Planesticus casius Bonaparte, 1855
caspica—Pyrrhula pyrrhula caspica Witherby, 1908 59 Bull. Brit. Orn. Cl., 23: 48. HOLOTYPE, Adult male. Reg. no. 1907.12.21.569. South coast of Caspian Sea. 25 Feb. 1907. Collected by R. B. Woosnam. Presented by A. C. Bailward.
caspius—Garrulus atricapillus caspius Seebohm, 1883
Cassidix—Ptilotis cassidix Gould, 1867

- cassini—Sycobius cassini Elliot, 1859 61

 Ibis, (1) 1: 392. HOLOTYPE, Adult male. Reg. no. 1859.6.28.12.

 Gaboon, French West Africa. Received in May, 1859. Collected for and purchased of Verreaux.

- castaneiceps—Basileuterus castaneiceps P. L. Sclater and Salvin, 1877 54 Proc. zool. Soc. Lond., 1877: 521. HOLOTYPE, Adult. Reg. no. 1885.3.8.659. Jina (or Jima), Ecuador. Collected by C. Buckley. Salvin-Godman Collection.
- castaneiceps—Buarremon castaneiceps P. L. Sclater, 1859 . . . 49
 Proc. zool. Soc. Lond., 1859: 441. HOLOTYPE, Adult. Reg. no.
 1855.6.12.914. Rio Napo, Ecuador. Collected for Verreaux.
 Sclater Collection.
- castaneiceps—Conopophaga castaneiceps P. L. Sclater, 1857 . . . 5

 Proc. zool. Soc. Lond., 1857 : 47. SYNTYPE, Adult male. Reg. no. 1854.1.25.83. Bogota, Colombia. Purchased of Stevens.

The other syntype is in the collection. The specimen listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 332, is not accepted as typical material as it was not listed by the author in his Catalogue of American Birds, 1862: 193, and therefore must have been received by him well after the description was published.

Moore used a Gould MS. name. The specimen was listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 544.

- castaneiceps—Laniarius castaneiceps Sharpe, 1891 . . . 23

 Ibis, (6) 3: 445. HOLOTYPE, Adult female. Reg. no. 1893.12.1.19.

 Mount Elgon, central East Africa, 25 Feb. 1890. Collected and presented by F. J. Jackson.
- castaneiceps—Orthotomus castaneiceps Walden, 1872 35

 Ann. Mag. nat. Hist., (4) 10: 252. Holotype, Adult male. Reg.
 no. 1888.4.20.386. Guimaras Island, Philippines, March 1872.
 Collected by A. B. Meyer. Tweeddale Collection.
- castaneicollis—Chlorospingus castaneicollis P. L. Sclater, 1858 . 51

 Proc. zool. Soc. Lond., 1858: 293. HOLOTYPE, Adult. Reg. no. 1885.6.12.883. Interior of Peru . . . bordering on Bolivia. Collected for Verreaux. Sclater Collection.
- castaneiventris—Monarcha castaneiventris J. Verreaux, 1858. . . 38

 Rev. Mag. Zool., 10:304. HOLOTYPE, Adult male. Reg. no. 1858.5.8.13. Samoa, Navigator's Islands.

 Purchased of Cuming, the collection originally stated to be from Verreaux, but later ascribed to Cuming. (See also appendix, Corrections and Additions, vol. 2.)
- castaneothorax—Cinclosoma castaneothorax Gould, 1849 . . . 32A Proc. zool. Soc. Lond., 1848 (publ. 1849): 139, pl. 6. HOLOTYPE, Adult male (in very poor condition). Reg. no. 1855.11.5.31. Scrubby belts of trees growing on the tableland to the northward of the Darling Downs, New South Wales, vide Gould, The Birds of Australia, 1 (1865): 438. Collected by J. Coxen and purchased of J. Gould.
- castaneum—Philentoma castaneum Eyton, 1845 3'
 Ann. Mag. nat. Hist., (1) 16: 229. Syntype, Adult male (relaxed mount). Reg. no. 1846.3.4.31. Malacca, Malaya. Collected and presented by A. Charlton.

This and a female syntype were relaxed from mounting in 1884
and were omitted from Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 367.
Two further syntypes were received by Eyton and are also in the
collection.

castaneus—Mviophoneus castaneus R. G. W. Wardlaw Ramsay, 1880 31 Proc. zool. Soc. Lond., 1880: 16, pl. 1. SYNTYPE, Adult. Reg. no. 1888.4.20.45. Mount Sago, Pandang, Sumatra, 3 Sept. 1878. Collected by C. Bock. Tweeddale Collection.

This is the author's selected syntype. There is a second syntype

in the collection.

castanicauda—Siva castanicauda Hume, 1877 Stray Feathers, 5:100. HOLOTYPE, Adult. Reg. no. 1886.10.1. 6790. Moolyit, Tenasserim, 17 Feb. 1877 (on label; 7 Feb. in text). Collected by W. Davison. Hume Collection.

castaniceps—Abrornis castaniceps Hodgson, 1845. in Blyth, J. Asiat. Soc. Bengal, 14:593. SYNTYPE, Adult. Reg. no. 1843.1.13.386. Nepal. Presented by B. H. Hodgson.

castanonota—Alethe castanonota Sharpe, 1871 Catalogue of African Birds: 20. HOLOTYPE, Adult. Reg. no. 1876.5.23.523. Fantee. Gold Coast. Collected by Ussher. Sharpe Collection.

castanonota—Lochmias nematura castanonota Chubb, 1919 Bull. Br. Orn. Cl., 38:87. HOLOTYPE, Adult female. Reg. no. 1889.5.14.77. Aruparu River (Arranparu on label), 5,000', Kukenam Mountains, British Guiana, 29 Aug. 1883. Collected by H. Whitely. Salvin-Godman Collection.

castanonotus—Hemixus castanonotus Swinhoe, 1870 Ibis, (2) 6:251, pl. 9. HOLOTYPE, Adult. Reg. no. 1898.10.2.83. Tai-ping-sze, central Hainan, China, Feb. 1868. Collected by R. Swinhoe. Seebohm Bequest.

1887.9.28.62. Lado, Wadelai District, Equatorial Africa (vide, op. cit.: 18-19), 12 Aug. 1885. Collected for and presented by Emin Pasha.

There are four other syntypes in the collection.

castanopsis—Heliolais castanopsis Vincent, 1933. Bull. Brit. Orn. Cl., 53: 140. HOLOTYPE, Adult male. Reg. no. 1933.3.1.9. Mouth of Lurio River, Portuguese East Africa, 13°30′ S. 40°30′ E., 29 June 1932. Collected by J. Vincent. Presented by the Portuguese East Africa Expedition.

castanopterus—Pyriglena castanopterus Chubb, 1916 Bull. Br. Orn. Cl., 36:47. SYNTYPE, Adult female. Reg. no. 1916.8.24.48. Baeza, 6,000', east Ecuador, April 1914. Collected by W. Goodfellow and presented by E. J. Brook. There is another syntype in the collection.

castus—Monarcha castus P. L. Sclater, 1883	
catasticta—Tanagra catastica Oberholser, 1918	
caterinae—Saxicola caterinae Whitaker, 1898	
cathkinensis—Bradypterus barratti cathkinensis Vincent, 1948 . 3 Bull. Brit. Orn. Cl., 69:18. HOLOTYPE, Adult male. Reg. no. 1948.39.8. Near Cathkin Peak, 7,000′, Giant's Castle Game Reserve, Natal, 18 Oct. 1947. Collected and presented by J. Vincent.	35
caucasicus—Turdus torquatus caucasicus. See appendix, Corrections and Additions, vol. 2	31
caudata—Capsiempis caudata Salvin, 1897. Bull. Brit. Orn. Cl., 7:16. HOLOTYPE, Adult female. Reg. no. 1895.11.27.136. Ourumee, British Guiana, 22 Oct. 1890. Collected by H. Whitely. Salvin-Godman Collection. (See also Hellmayr, Novit. Zool. 13 (1906): 323.)	9
caudata—Formicivora caudata P. L. Sclater, 1855 Proc. zool. Soc. Lond., 1854 (publ. 1855) : 254, pl. 74. SYNTYPE, Adult male. Reg. no. 1889.9.20.360. Bogota, Colombia. Collected for Verreaux. Sclater Collection. There is a female syntype in the collection.	4
caudatus—Pseudotharrhaleus caudatus Ogilvie-Grant, 1895 Bull. Brit. Orn. Cl., 4:40. SYNTYPE, Adult male. Reg. no. 1897.5.13.263. Mount Data, Lepanto District, Northern Luzon, Philippines, 29 Jan. 1895. Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. A male and female were collected, vide Ogilvie-Grant, Ibis, (7) 1 (1895):449. The latter is now in the American Museum of Natural History, vide Hartert, Novit. Zool., 27 (1930):484.	
cavei—Bradypterus cinnamomeus cavei Macdonald, 1939	

- cavei—Certhilaudi albescens cavei Macdonald, 1953 17

 Bull. Brit. Mus. nat. Hist. (Zool.), 1:344. HOLOTYPE, Adult male.

 Reg. no. 1950.50.922. Smiles, 4,000′, 5 miles south-west of Witputs,

 Great Namaqualand, 26 Jan. 1950. Collected and presented by
 the British Museum South-West Africa Expedition.
- cavei—Eremopterix leucopareia cavei C. H. B. Grant and Mackworth-Praed, 1941 Bull. Br. Orn. Cl., 61:62. HOLOTYPE, Adult female. Reg. no. 1940.4.6.5. Nataparapoth, 2,000', Ataporopos Hill, Eastern District, Equatoria, Sudan, 15 March 1940. Collected and presented by F. O. Cave.
- cavendishi—Estrilda cavendishi Sharpe, 1900 60

 Ibis, (7) 6:110. Holotype, Adult male. Reg. no. 1898.11.16.39.

 Mapicuti, Mozambique, 15 Sept. 1898. Collected and presented by H. S. H. Cavendish.
- cavernicola—Thamnolaea cinnamomeiventris cavernicola Bates, 1933 31 Bull. Br. Orn. Cl., 53:176. HOLOTYPE, Adult male. Reg. no. 1932.8.6.10 (not 1932.8.6.18 as printed). Fiko, 30 miles east of Mopti, French Sudan, 10 Dec. 1931. Collected by and purchased of G. L. Bates.
- cayennensis—Coereba chloropyga cayennensis P. R. Lowe, 1912 . 54

 Ibis, (9) 6:506. HOLOTYPE, Adult. Reg. no. 1885.4.1.341.

 Oyapoc, Cayenne, French Guiana. Collected by Mde. Verdey.

 Salvin-Godman Collection.

The female syntype is also in the collection.

celebensis—Anthreptes celebensis Shelley, 1878
celebensis—Basilornis celebensis G. R. Gray, 1861 62 Proc. zool. Soc. Lond., 1861: 184, fig. 2. SYNTYPE, Adult male. Reg. no. 1873.5.12.1867. Menado, North Celebes Island, 1859. Collected by A. R. Wallace. Wallace Collection. Gray used a Temminck MS. name. There are several other syntypes in the collection.
celebensis—Broderipus celebensis Walden, 1872
centralasiae—Emberiza pyrrhuloides centralisiae Hartert, 1903 . 49 Vögel der Palaarktischen Fauna, 1:199. Holotype, Adult male. Reg. no. 1897.12.10.589. Maralbashi, 4,600', East Turkestan, Jan. 1874. Collected and presented by J. Biddulph.
cephaëlis—Cinnyris (Cyanomitra) olivaceus cephaëlis Bates, 1930 . 46 Bull. Brit. Orn. Cl., 51:52. HOLOTYPE, Adult male. Reg. no. 1930.12.4.1. Bitye, 2,000', Ja River, Cameroon, 20 March 1924. Collected by and purchased of G. L. Bates.
ceramensis—Cryptolopha ceramensis Ogilvie-Grant, 1910 35 Bull. Brit. Orn. Cl., 25:90. Syntype, Adult male. Reg. no. 1910.12.28.278. Karobi Mountain, 2,500', Ceram Island, Moluccas, Oct. 1909. Collected by W. Stalker. Presented by the subscribers to the Expedition to Dutch New Guinea. The female syntype is also in the collection.
ceramensis—Erythromyias ceramensis Ogilvie-Grant, 1910 37 Bull. Brit. Orn. Cl., 25: 90. HOLOTYPE, Adult male. Reg. no. 1910.22.28.199. Selemen, 3,000', Ceram Island, Moluccas, Nov. 1909. Collected by W. Stalker. Presented by the subscribers to the British Ornithologists' Union Expedition to Dutch New Guinea.
certus—Sclerurus mexicanus certus Chubb, 1919

- cervicalis—Philydor cervicalis P. L. Sclater, 1889 3

 Proc. zool. Soc. Lond., 1889 : 33. Syntype, Immature (vide Hellmayr, Novit. Zool., 13 (1906) : 335). Reg. no. 1889.5.20.372. Camacusa, British Guiana, 29 May 1882. Collected by H. Whitely. Sclater Collection.

Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 102. There are several other syntypes in the collection.

cervinicauda—Drepanornis albertisi cervinicauda P. L. Sclater, 1883 70 Proc. zool. Soc. Lond., 1883: 578. SYNTYPE, Adult male. Reg. no. 1883.12.17.3. Astrolabe Mountains, Port Moresby, South New Guinea. Collected by Goldie. Presented by Dr. G. Bennett, Sydney.

Three female syntypes are also in the collection.

The female syntype is also in the collection.

Listed as type by Sclater, Catalogue of American Birds, (1862): 158. There is another syntype in the collection.

- cerviniventris—Chlamydera cerviniventris Gould, 1850 69
 in Jardine, Contributions to Ornithology: 100 or 106 or 160 (page number in doubt). (The description in Proc. zool. Soc. Lond., 1850: 201 was not published until Feb. 1851.) Syntype, Adult. Reg. no. 1851.1.11.11. Cape York, northern Australia. Collected and presented by J. Macgillivray, from the voyage of H.M.S. Rattlesnake.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 393. A Gould specimen from the same locality is thought to be another syntype.

cerviniventris—Digenea cerviniventris Sharpe, 1879 37 Cat. Bds. Brit. Mus., 4:460. Holotype, Adult male. Reg. no. 1878.10.14.1. Manipur Hills, Burma. Collected and presented by H. H. Godwin-Austen.

cerviniventris—Phyllostrophus cerviniventris Shelley, 1894 . . . 27

Ibis, (6) 6:10, pl. 2. SYNTYPE, Adult. Reg. no. 1893.6.1.98.

Zomba, Nyasaland, Aug. 1892. Collected by A. Whyte and presented by Sir H. H. Johnston.

The six other syntypes are also in the collection.

collection, but are in bad condition.

cerviniventris—Rectes cerviniventris G. R. Gray, 1862 39

Proc. zool. Soc. Lond., 1861 (publ. 1862): 430. Syntype, Adult male. Reg. no. 1873.5.12.1300. Gagie Island (off western New Guinea). 1860. Collected by A. R. Wallace. Wallace Collection.

There are two other syntypes in the collection.

- cerviniventris—Turdinus cerviniventris Sharpe, 1901 32B Bull. Br. Orn. Cl., 12:3. Syntype, Adult male. Reg. no. 1895.5. 1.1377. Condé, Congo, 1876. Collected by L. Petit. Shelley Collection; from A. Bouvier's collection.

 There are at least two other syntypes in the collection.
- ceylonensis—Dicaeum erythrorhynchus ceylonensis Babault, 1920 . 45

 Mission Babault dans les Centrales Provinces de l'Inde: 293, pl. 5.

 Boista, South Ceylon, 15 May 1871. Collected by W. V. Legge.

 Purchased of E. Oates.
- ceylonensis—Dissemurus ceylonensis Sharpe, 1877 64
 Cat. Bds. Brit. Mus., 3:264. SYNTYPE, Immature female. Reg.
 no. 1876.6.6.3. Pan Colom, North-east Province, Ceylon, 7 Jan.
 1873. Collected and presented by W. V. Legge.

The specimen was loaned to Seebohm and received back with his collection, being given a second reg. no. 1897.6.1.591. There are several other syntypes in the collection.

- no. 1934.2.15.1. 8 miles north of Paletupani, east coast of Ceylon, 11 Jan. 1922. Collected by G. Brown and received in exchange from H. Whistler.
- ceylonensis—Hypothymis ceylonensis Sharpe, 1879 37

 Cat. Bds. Brit. Mus., 4:277. Syntype, Adult male. Reg. no. 1877.11.9.22. Kandy Hills, Ceylon. Collected by A. Whyte. Purchased of E. Gerrard.

 The female syntype is also in the collection.
- ceylonensis—Mirafra affinis ceylonensis Whistler, 1936. . . . 17

 J. Bombay nat. Hist. Soc., 38: 769. HOLOTYPE, Adult male. Reg.
 no. 1887.7.1.2667. Colombo, Ceylon, 7 Nov. 1877. Collected by
 the Hart brothers. Hume Collection
- ceylonensis—Pericrocotus peregrinus ceylonensis Whistler, 1933 . 20 J. Bombay nat. Hist. Soc., 36:344. HOLOTYPE, Adult male. Reg. no. 1916.9.20.621. Cocawatte, Ceylon, 7 Nov. 1895. Collected and presented by A. L. Butler.
- chadensis—Batis chadensis Boyd Alexander, 1908 38
 Bull. Brit. Orn. Cl., 21:105. SYNTYPE, Adult female. Reg. no.
 1911.12.23.1455. Arrigi, Lake Chad (30 Nov. 1904, on label).
 Collected by Boyd Alexander on the Alexander-Gosling Expedition.
 Boyd Alexander bequest.

The male syntype is also in the collection, but since this was stated to be similar to the male of *B. orientalis* the female has been selected here.

- chadensis—Bradypterus brachypterus chadensis Bannerman, 1936. 35
 Bull. Brit. Orn. Cl., 57: 43. HOLOTYPE, Adult male. Reg. no. 1936.2.21.1. Lake Chad, Central Africa, 1 Dec. 1904 (Nov., on label). Collected by Boyd Alexander. Presented by the Alexander-Gosling Expedition.
- chadensis—Calamocichla chadensis Boyd Alexander, 1907 . . . 35

 Bull. Brit. Orn. Cl., 19:63. SYNTYPE, Adult male. Reg. no.
 1911.12.23.2837. Lake Chad, French Equatorial Africa, (Wurmda,
 19 May 1905, on label). Collected by W. Boyd Alexander on the
 Alexander-Gosling Expedition. Boyd Alexander Bequest.

 This specimen was labelled as the type but there are several other

syntypes in the collection.

12.23.2997. Kowa Baga, Lake Chad, West Africa, 19 April Collected and bequeathed by Boyd Alexander. This is the author's selected male. His selected female synalso in the collection.	
chadensis—Tchagra senegala chadensis Bannerman, 1922. Rev. Zool. Afr., 9 (1921, publ. 1922): 355. HOLOTYPE, Adult Reg. no. 1911.12.23.2029. Yo, Lake Chad, Northern N 5 Jan. 1905. Collected by Boyd Alexander on the Alex Gosling Expedition. Boyd Alexander Bequest.	ligeria,
Chalcea —Nectarinia chalcea Hartlaub, 1862	
chalybeus—Turdus chalybeus Horsfield, 1821	1816.
Chapini —Cossypha bocagei chapini Benson, 1955. Bull. Br. Orn. Cl., 75: 104. HOLOTYPE, Adult male. Re 1955.41.2. Mporokoso, Northern Rhodesia, 30 July 1953. lected by C. W. Benson and presented by the National Muse Southern Rhodesia.	Col-
Chapini—Mirafra africana chapini C. H. B. Grant and Mack Praed, 1939 Bull. Br. Orn. Cl., 59: 140. HOLOTYPE, Adult female. Re 1931.12.21.21. Nasondoye, 3,350', south-east Belgian C 3 Jan. 1931. Collected and presented by Admiral H. Lynes.	. 17 eg. no. Congo,
chapmani—Pachyrhamphus chapmani Chubb, 1920 Bull. Brit. Orn. Cl., 40: 74. HOLOTYPE, Adult male. Re 1888.1.20.522. Santa Elena, Antioquia, Colombia, Sept. Collected by T. K. Salmon. Salvin-Godman Collection.	. 7 eg. no. 1878.
Chapmani — <i>Pipromorpha oleaginea chapmani</i> Chubb, 1919 . <i>Ann. Mag. nat. Hist.</i> , (9) 4:302. HOLOTYPE, Adult. Re 1899.4.20.241. Villaviciencio to Medina, Llanos of Meta 800–1.500′. Colombia. Collected by T. H. Wheeler.	River,

charmensis—Carpodacus thura charmensis Kinnear, 1940 . . . 59
Bull. Brit. Orn. Cl., 60: 56. SYNTYPE, Adult female. Reg. no.
1937.1.17.837. Le La, 15,000', South-east Tibet, 8 May 1936.
Collected by and purchased of F. Ludlow and G. Sheriff.
The male syntype is also in the collection.

Godman Collection.

- chayulensis—Suthora fulvifrons chayulensis Kinnear, 1940 . . . 33

 Bull. Br. Orn. Cl., 60: 56. HOLOTYPE, Adult male. Reg. no. 1937.1.17.838. Lung, 11,000′, Chayul Valley, south-east Tibet, 29 April 1936. Collected by F. Ludlow and G. Sherriff, and presented by the Godman-Thomas Fund.
- cheesmani—Ammomanes deserti cheesmani R. Meinertzhagen, 1923 17 Bull. Br. Orn. Cl., 43:157. HOLOTYPE, Adult. Reg. no. 1923.6. 19.1. Shatt-al-Adhaim, left bank of Tigris, Mesopotamia, 16 Oct. 1917. Collected by C. R. S. Pitman and presented by the Bombay Natural History Society.

- chermesina—Myzomela chermesina G. R. Gray and Mitchell, 1846 48
 Genera of Birds, 1: pl. 38. Holotype, Adult male (relaxed mount).
 Reg. no. 1846.7.16.147. (No locality given.) Presented by E. Wilson.

 See also reference to type by Forbes, Proc. zool. Soc. Lond., 1879:
- chiapensis—Campylorhynchus chiapensis Salvin and Godman, 1891 28 Ibis, (6) 3:609. HOLOTYPE, Adult female. Reg. no. 1890.12.20. 727. Tonala, Chiapas, Mexico, 6 Jan. 1890. Collected by W. B. Richardson. Salvin-Godman Collection.

- chianensis—Melozone biarcuatum chiavensis van Rossem, 1938 . . Bull. Brit. Orn. Cl., 58: 126. HOLOTYPE, Adult female. Reg. no. 1899.2.1.3938. Tuxtla, Chiapas, Mexico, 18 Mar. 1897. Collected by W. B. Richardson. Salvin-Godman Collection. chiapensis—Pipilo maculatus chiapensis van Rossem. 1938 Bull. Brit. Orn. Cl., 58: 130. HOLOTYPE, Adult male. Reg. no. 1899.2.1.3480. San Cristobal, Chiapas, Mexico, 14 May, 1897. Collected by W. B. Richardson. Salvin-Godman Collection. chiguancoides—Turdus chiguancoides Seebohm, 1881 Cat. Bds. Brit. Mus., 5: 231. SYNTYPE, Adult. Reg. no. 1896.6.1. 405. Plains of the Gambia River. Seebohm Collection. There are three other syntypes in the collection. chiniana—Drymoica chiniana Smith, 1843 Illustrations of the Zoology of South Africa, Aves: pl. 79. A single specimen was described. A specimen registered and labelled as the type, Reg. no. 1843.2.23.108. resembles the description in some aspect but lacks the distinctive head colour and was re-identified at an early date as a specimen of D. natalensis Smith, the label being altered. chirindensis—Apalis chirindensis Shelley, 1906 Bull. Brit. Orn. Cl., 16: 126. HOLOTYPE, Adult. Reg. no. 1911.5. 30.33. Chirinda Forest, 4,000', Northern Gazaland, Southern Rhodesia, 14 June 1905. Collected and presented by C. F. M. Swynnerton. chiriquensis—Geothlypis chiriquensis Salvin, 1872 Ibis, (3) 2:148. HOLOTYPE, Adult male. Reg. no. 1885.3.8.512. Volcan de Chiriqui, Veragua, 1870. Collected by E. Arcé. Salvin-Godman Collection. chiroleuca—Gymnocichla chiroleuca P. L. Sclater and Salvin, 1869 Proc. zool. Soc. Lond., 1869:417. SYNTYPE, Adult male. Reg. no. 1889.7.10.603. Tucurriqui, Costa Rica, 1864. Collected by E. Arcé. Salvin-Godman Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 272. chlorocephalus—Oriolus chlorocephalus Shelley, 1896 . 63 Ibis, (7) 2:183. HOLOTYPE, Adult female. Reg. no. 1896.3.14.44. Mount Chiradzulu, Nyasaland. Collected by A. Whyte. Presented by H. H. Johnstone.
- chlorocephalus—Phyllornis chlorocephalus Walden, 1871 . . . 22

 Ann. Mag. nat. Hist., (4) 7: 241. Syntype, Adult male. Reg. no. 1888.4.1.200. Near Tonghoo, Burma. Collected by Lloyd. Tweeddale Collection.

 The two other males and one female syntype are also in the

The two other males and one female syntype are also in the collection.

The label indicates that this is a Verreaux specimen. W. L. Sclater (in MS., now destroyed) stated "this, though not so marked, appears to be the type of the species."

- chloronotus—Abrornis chloronotus J. E. Gray, 1846 35 in Hodgson, Catalogue of the birds of Nepal, 1846 : 66, 152. Syn-Type, Adult. Reg. no. 1845.1.9.815. Nepal. Collected and presented by B. H. Hodgson.

Gray used a Hodgson MS number. Ticehurst, Systematic revue of the genus Phylloscopus, 1938:116, refers to this specimen as the type.

Renamed Z. gouldi by Bonaparte, Consp. Av., 1 (1850): 398, on the grounds of preoccupation by Dicaeum chloronothos Vieillot, 1817.

chlorophrys—Diaphorophyia chlorophrys Boyd Alexander, 1903. 38

Bull. Brit. Orn. Cl., 13:34. HOLOTYPE, Adult male. Reg. no.
1911.12.23.1442. Bakaki, Fernando Po, 19 Nov. 1902. Collected
by Boyd Alexander. Boyd Alexander Bequest.

chloroptera—Melanocharis chloroptera Salvadori, 1875
chloroptera—Myzomela chloroptera Walden, 1872
chloropygia—Nectarinia chloropygia Jardine, 1842
chlorurus—Pachycephala chlorurus G. R. Gray, 1859 39 Catalogue of birds of the tropical islands of the Pacific : 20. Syntype, Adult male. Reg. no. 1859.6.14.16. (Erromango Island, on label), New Hebrides. Collected by J. Macgillivray. Purchased of S. Stevens. The second syntype is in the collection.
chocolatina—Pnoepyga chocolatina Godwin-Austen and Walden, 1875
choloensis—Alethe choloensis W. L. Sclater, 1927 31 Bull. Br. Orn. Cl., 47: 86. HOLOTYPE, Adult female ("not breeding" on label). Reg. no. 1927.2.2.7. Cholo Mountain, Nyasaland, 6 Oct. 1926. Collected and presented by C. F. Belcher.
christianae—Cinnyris christianae Tristram, 1889
christinae—Aethopyga christinae Swinhoe, 1869

- christophori—Pachycephala christophori Tristram, 1879 . . . 39 Ibis, (4) 3:441. A specimen stated to be the type is listed by Sharpe, Cat. Bds. Brit. Mus., 8 (1883):201, but has not been traced in the collection. Tristram, Catalogue of a collection of birds (1889):191, states that the type was in his collection. Since this collection is now in the Liverpool Museum the type should be there also.

See C. H. B. Grant and Mackworth-Praed, *Bull. Br. Orn. Cl.*, 62 (1942): 43. for a note on the date of collection.

Two specimens in the collection were listed as types by Sharpe, *Cat. Bds. Brit. Mus.*, 7 (1883): 602, but the material used by Blyth is presumably still at the Calcutta Museum.

Stachyris chrysaea J. E. Gray, Zoological Miscellany, 1844:82, is a nomen nudum. Both authors used Hodgsons' MS. name.

The tail measurement quoted is inaccurate.

- chrysocephalum—Heteropelma chrysocephalum Pelzeln, 1869. 8

 Ornithologie Brasilien: 25, 185. The Natterer specimen received by Sclater (Cat. Bds. Brit. Mus., 14 (1888): 322) in exchange from the Vienna Museum and listed by him as "authentic" is not a typical specimen and is now considered to belong to a different species. It has not therefore been segregated as a syntype.
- chrysogaster—Chloropsis chrysogaster McClelland, 1840 . . . 22 in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840): 167. SYNTYPE, Adult male (in poor condition). Reg. no. 1860.4.16.258. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.

chrysogaster—Gerygone chrysogaster G. R. Gray, 1858 . . . 36

Proc. zool. Soc. Lond., 1858: 174. SYNTYPE, Adult male. Reg.
no. 1858.3.10.65. Aru Islands, 1857. Collected by A. R. Wallace.
Wallace Collection.

The female syntype is also in the collection.

- chrysogaster—Megarhynchus chrysogaster P. L. Sclater, 1860 . 9
 Proc. zool. Soc. Lond., 1860: 281. HOLOTYPE, Adult male. Reg.
 no. 1888.1.13.794. Babahoyo, Ecuador, Aug. 1859. Collected by
 L. Fraser Sclater Collection.
- chrysomelas—Tachyphonus chrysomelas P. L. Sclater and Salvin, 1869 51 Proc. zool. Soc. Lond., 1869: 440. pl. 32. SYNTYPE, Adult male. Reg. no. 1885.6.7.654. Cordillera del Chucu, Veragua, 1869. Collected by E. Arcé. Salvin-Godman Collection.

 This is the specimen figured. The female syntype is also in the

This is the specimen figured. The female syntype is also in the

collection.

chrysonota—Calliste chrysonota P. L. Sclater, 1850 5 in Jardine, Contributions to Ornithology: 50, pl. 51. SYNTYPE, Adult male. Reg. no. 1885.6.12.279. Cayenne. Collected for Verreaux. Sclater Collection.

This species was also described under the name of *C. luteola* by Sclater but the name was changed when pages in the above were suppressed and replaced (*vide* de Schauensee, *Proc. Acad. Nat. Sci. Philadelphia*, 109 (1957): 242).

- chrysoparia—Dendroeca chrysoparia Sclater and Salvin, 1860 . 54

 Proc. zool. Soc. Lond., 1860: 298. HOLOTYPE, Adult male. Reg.
 no. 1885.3.8.261. (Tactic, on label), Vera Paz, Guatemala. (4 Nov.
 1859, on label.) Collected by O. Salvin. Salvin-Godman Collection.
- chrysopasta—Euphonia chrysopasta P. L. Sclater and Salvin, 1869.
 51 Proc. zool. Soc. Lond., 1869: 438, pl. 30. SYNTYPE, Adult male.
 Reg. no. 1885.6.12.173. Lower Ucayali River, Eastern Peru.
 Collected by E. Bartlett. Sclater Collection.
 Listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 83.

There are four other syntypes in the collection.

chrysopeplus—Coccothraustes chrysopeplus Vigors, 1832 . . . 50

Proc. zool. Soc. Lond., 1832 : 4. HOLOTYPE, Adult male. Reg. no. 1855.12.19.19. Mexico (restricted to San Blas, Nayarit, by van Rossem, Bull. Mus. Comp. Zool., Harv. 77 (1934) : 480). Collected by H. Cuming. Purchased of the Zoological Society of London.

chrysophrys—Calliste chrysophrys P. L. Sclater, 1851
chrysophrys—Chlorospingus chrysophrys P. L. Sclater and Salvin, 1875
chrysopis—Nemosia chrysopis P. L. Sclater and Salvin, 1880 51 Proc. zool. Soc. Lond., 1880: 155. Holotype, Immature (vide Hellmayr, Novit. Zool., 13 (1906): 310). Reg. no. 1885.6.7.706. Sarayacu, Ecuador. Collected by C. Buckley. Salvin-Godman Collection.
chrysopleurus—Turdus chrysopleurus Swinhoe, 1874 31 Ibis, (3) 4: 444. HOLOTYPE. This is also the holotype of Turdus campbelli Swinhoe, 1873 (q.v.).
chrysops—Setophaga chrysops Salvin, 1878
chrysops—Sycalis chrysops P. L. Sclater, 1862
chrysops—Sylvia chrysops Latham, 1801
chrysops—Tyrannulus chrysops P. L. Sclater, 1859 9 Proc. zool. Soc. Lond., 1858 (publ. 1859): 458. SYNTYPE, Adult (female, vide Sclater, Catalogue of a Collection of American Birds (1862): 216). Reg. no. 1888.1.13.504. Gualaquiza, Ecuador, Jan. 1858. Collected by L. Fraser. Sclater Collection. The second syntype is also in the collection.

cl	rysoptera—Diphyllodes speciosus var. chrysoptera Elliot, 1873 .	
	Monograph of the Paradiseidae: pl. 13 and text. Holotype, Adult	
	Reg. no. 1881.5.1.1733. (Jobi Island?, on label.) Gould Collect	; _
	tion.	

Gould lent Elliot the specimen for figuring.

Vol. 3, no. 6 in Lambert drawings, in library at Knowsley, Lancashire. (See introduction to this catalogue, vol. 2.) See also Mathews, Birds of Australia, 11 (1923): 55.

chrysopterus—Merops chrysopterus Latham, 1801. 4

Index Ornithologicus, Suppl.: xxxiii. ? Syntypes, Water-colour drawings. New South Wales. No. 89, 90 in Watling, Original water-colour drawings Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 125; Mathews, Birds of Australia 12 (1925): 82.

Vol. 1, no. 33 in Lambert Drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2). A signed drawing by Watling, no. 104, named *Certhia mellivora* by Latham, is also of this species.

chrysorrhoa—Eopsaltria chrysorrhoa Gould, 1869 . . . 3'
Ann. Mag. nat. Hist., (4) 4:109. SYNTYPE, Adult male. Reg. no.
1881.5.1.1104. (Moreton Bay, on label), east part of New South
Wales. Gould Collection.

This specimen was labelled as the type. The female syntype is also in the collection.

There is another syntype in the collection; the selected specimen corresponds better with the description. See Roberts, *Annals Transvaal Mus.*, 1935: 320 for a note on the validity of the name.

Kimangitchi, Mount Elgon, 27 Feb. 1890. Collected and presented by F. J. Jackson.

This specimen was presented as the type but a female syntype was also referred to in the description.

- chubbi—Lamprocolius chubbi Boyd Alexander, 1903 62

 Bull. Brit. Orn. Cl., 13:48. HOLOTYPE, Adult female. Reg. no. 1911.12.23.3737. Moka, Fernando Po, 16 Dec. 1902. Collected and bequeathed by Boyd Alexander.

This specimen was labelled as the type by the author but there are

several other syntypes in the collection.

- chyula—Alseonax minimus chyula V. G. L. van Someren, 1939 . 37 J. E. Afr. and Uganda nat. Hist. Soc., 14:71. HOLOTYPE, Adult male. Reg. no. 1948.21.8. Chyulu Range, 5,500–7,200′, Kenya, 14 June 1938. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.
- chyulu—Apalis griseiceps chyulu V. G. L. van Someren, 1939. . . 35 J. E. Afr. and Uganda nat. Hist. Soc., 14:97. HOLOTYPE, Adult male. Reg. no. 1948.11.16. (Camp 3, on label.) Chyulu Range, Kenya, 26 June 1938. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.
- chyulu—Arizelocichla milanjensis chyulu V. G. L. van Someren, 1939 21 J. E. Afr. and Uganda nat. Hist. Soc., 14:64. HOLOTYPE, Adult male. Reg. no. 1948.11.17. Camp 3, 7,000′, Chyulu Hills, Kenya, 6 July 1938. Collected by V. G. L. van Someren and presented by the Coryndon Museum, Nairobi.
- chyulu—Cyanomitra olivacea chyulu V. G. L. van Someren, 1939 . 46 J. E. Afr. and Uganda nat. Hist. Soc., 14:120. HOLOTYPE, Adult male. Reg. no. 1949.11.24. Chyulu Hills, 7,600', (Camp 3), Kenya, 14 July 1938. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.

- chyulu—Geokichla gurneyi chyulu V. G. L. van Someren, 1939 . 3

 J. E. Afr. and Uganda nat. Hist. Soc., 14:77. HOLOTYPE, Adult.

 Reg. no. 1948.11.12. Camp 3, 7,000′, Chyulu Range, Kenya,

 4 July 1938. Collected by V. G. L. van Someren and presented by
 the Coryndon Museum, Nairobi.
- chyulu—Laniarius ferrugineus chyulu V. G. L. van Someren, 1939 . 23 J. E. Afr. and Uganda nat. Hist. Soc., 14:107. HOLOTYPE, Adult male. Reg. no. 1948.11.22. Chyulu Hills, 5,600', Kenya, 18 June 1938. Collected by V. G. L. van Someren and presented by the Coryndon Museum, Nairobi.
- chyulu—Melocichla mentalis chyulu V. G. L. van Someren, 1939 . 35 J. E. Afr. and Uganda nat. Hist. Soc., 14:91. Holotype, Adult male. Reg. no. 1948.11.14. (Camp 3, on label), 5,600′, Chyulu Range, Kenya, 26 June 1938. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.
- chyulu—Pycnonotus tricolor chyulu V. G. L. van Someren, 1939. 21 J. E. Afr. and Uganda nat. Hist. Soc., 14:60. Holotype, Adult male. Reg. no. 1948.11.20. Chyulu Hills, 6,500', Kenya, 27 April 1938. Collected by V. G. L. van Someren and presented by the Coryndon Museum, Nairobi.
- chyulu—Schoenicola brevirostris chyulu V. G. L. van Someren, 1939 35 J. E. Afr. and Uganda nat. Hist. Soc., 14:95. HOLOTYPE, Adult female. Reg. no. 1948.11.15. Camp 2, 5,600′, Chyulu Range, Kenya, 10 June 1936. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.
- chyulu—Seicercus umbrovirens chyulu V. G. L. van Someren, 1939. 35

 J. E. Afr. and Uganda nat. Hist. Soc., 14:89. Holotype, Adult male. Reg. no. 1948.11.9. Chyulu Range, Kenya, 6,000–7,000′, 21 June 1938. (5,600′, 1 June 1938, on label.) Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.

 In spite of discrepancies between description and label, this specimen which is labelled as the type appears to be the one on which the description was based.
- chyulu—Spinus citrinelloides chyulu van Someren, 1939 . . . 59 J. E. Afr. and Uganda nat. Hist. Soc., 14:127. HOLOTYPE, Adult male. Reg. no. 1948.11.26. Chyulu Hills, 6,560', Kenya, 24 April 1938. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.

chyuluensis—Anthus nicholsoni chyuluensis V. G. L. van Someren,
J. E. Afr. and Uganda nat. Hist. Soc., 14:56. HOLOTYPE, Adult male. Reg. no. 1948.11.25. Chyulu Camp, 5,000', Kenya, 4 May 1938 (in text; 4 June on label). Collected by V. G. L. van Someren and presented by the Coryndon Museum, Nairobi.
chyuluensis—Bradornis pallidus chyuluensis V. G. L. van Someren, 1939
Chyuluensis—Bradypterus cinnamomeus chyuluensis V. G. L. van Someren, 1939
chyuluensis—Chlorocichla flaviventris chyuluensis V. G. L. van Someren, 1939
chyuluensis—Cryptospiza salvadorii chyuluensis V. G. L. van Someren, 1939
chyuluensis—Phyllastrephus fischeri chyuluensis V. G. L. van Someren, 1939
chyuluensis—Zosterops chyuluensis V. G. L. van Someren, 1939 . 47 J. E. Afr. and Uganda nat. Hist. Soc., 14:114. Holotype, Adult male. Reg. no. 1948.11.24. Chyulu Range, 6,800', Camp 3, Kenya, 26 June 1938. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.
cinctura—Melanocorypha cinctura Gould, 1839

cinderella—Urolais epichlora cinderella Bates, 1928 35 Bull. Brit. Orn. Cl., 49:31. HOLOTYPE, Adult male. Reg. no. 1926.8.1.418. Oku, 6,000′, Banso Mountains, west of Kumbo, Cameroons, 11 Feb. 1925. Collected by and purchased of G. L. Bates.
cineracea—Myzomela cineracea P. L. Sclater, 1879 48 Proc. zool. Soc. Lond., 1879 : 448, pl. 37. HOLOTYPE, Adult male. Reg. no. 1881.3.29.79. New Britain, May 1878. Collected by the Rev. G. Brown. Purchased of E. Gerrard.
cineraceum—Trochalopteron cineraceum Godwin-Austen, 1874. 32B Proc. zool. Soc. Lond., 1874: 45, pl. 11. HOLOTYPE, Adult male. Reg. no. 1895.7.14.2249. Thobal Valley, Naga Hills, Munipur, India, 1872–73. Collected and presented by H. H. Godwin-Austen.
cineraceus—Edolius cineraceus Horsfield, 1821
cinerascens—Eurycercus cinerascens Walden, 1874 35 Ann. Mag. nat. Hist., (4) 14:156. HOLOTYPE, Adult male. Reg. no. 1888.4.20.496. Dobri, Lower Bengal, 27 Nov. 1873. Collected by F. Day. Tweeddale Collection.
cinerascens—Formicivora cinerascens P. L. Sclater, 1857 4 Proc. zool. Soc. Lond., 1857: 131. SYNTYPE, Adult male. Reg. no. 1855.12.17.92. Rio Napo, Ecuador. Gould Collection. Listed as type by Hellmayr, Novit. Zool., 12 (1905): 287. There is a female syntype in the collection.
cinerascens—Gerygone cinerascens Sharpe, 1878 36 J. Linn. Soc. Lond., 13: 494. HOLOTYPE, Adult. Reg. no. 1877.2. 9.16. South-east New Guinea. Collected by and purchased of O. C. Stone.
cinerascens—Muscicapa cinerascens Sharpe, 1879
cinerascens—Trichophorus cinerascens Hartlaub, 1858 21 Proc. zool. Soc. Lond., 1858 : 293. HOLOTYPE, Adult. Reg. no. 1858.1.4.122. Ashantee, West Africa. Purchased of J. Gould.
cinerea—Colluricincla cinerea Vigors and Horsfield, 1827 39 Trans. Linn. Soc. Lond., 15: 214. SYNTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.18a. Australia (=New South Wales? vide Mathews, Birds of Australia, 8 (1920): 240). Collected by Caley. Presented by the Linnean Society. The second syntype is also in the collection.

A female syntype in the Academy of Natural Sciences of Philadelphia was referred to as the type by W. Stone and G. M. Mathews, Austral Avian Record, 1 (1913): 180, but not listed by De Schauensee,

Proc. Acad. nat. Sci. Phil., 109 (1957): 123-246.

- cinereiceps—Cyclorhynchus cinereiceps P. L. Sclater, 1859 . . . 9

 Ibis, (1) 1: 443. HOLOTYPE, Adult (male, vide Sclater, Catalogue of a collection of American Birds, 1862: 220). Reg. no. 1888.1.13.715.

 Playa Vicente, Oaxaca, Mexico. Collected by A. Sallé for A. Boucard. Sclater Collection.
- cincreiceps—Drymocataphus cinereiceps Tweeddale, 1878 . . . 32B Proc. zool. Soc. Lond., 1887: 617. Holotype, Adult female. Reg. no. 1888.4.20.1134. Puerto Princesa, Palawan, Philippines, Jan. 1878. Collected by A. Everett. Tweeddale Collection.
- cinereiceps—Hylophilus cinereiceps P. L. Sclater and Salvin, 1860. 56

 Proc. zool. Soc. Lond., 1860: 299. HOLOTYPE, Adult. Reg. no. 1885.3.10.151. Choctam, Vera Paz, Guatemala. (Jan. 1860, on label.) Collected by O. Salvin. Salvin-Godman Collection.

cinereiceps—Pseudogerygone cinereiceps Sharpe, 1886 36 Nature, 34: 340. Holotype, Adult (female, vide Gould, Birds of New Guinea, 2 (1886): pl. 14). Reg. no. 1888.3.24.44. South-east
New Guinea. Vide Gould, op. cit. (Sogeri District, 1,750', Astrolabe Mountains, 26 Oct. 1881, on label). Collected by and purchased of H. O. Forbes.
cinereiceps—Trochalopterum cinereiceps Styan, 1887 32B Ibis, (5) 5:167, pl. 6. SYNTYPE, Adult. Reg. no. 1887.3.8.23. "Bought alive in Hankow, said to be from Yunnan, south-west China." Collected and presented by F. W. Styan. The second syntype was still alive at the date of first description.
cinereiceps—Tyrannulus cinereiceps P. L. Sclater, 1860 9 Proc. zool. Soc. Lond., 1860 : 69. HOLOTYPE, Adult female. Reg. no. 1888.1.13.478. Pallatanga, Ecuador (Oct. 1858, on label). Collected by L. Fraser. Sclater Collection.
cinereicollis—Basileuterus cinereicollis P. L. Sclater, 1864 54 Proc. zool. Soc. Lond., 1864: 166. SYNTYPE, Adult. Reg. no. 1884.5.15.444. Interior of Colombia. Collected for Verreaux. Sclater Collection. There is another syntype in the collection.
cinereicollis—Phyllergates cinereicollis Sharpe, 1888 35 Ibis, (5) 6: 479. HOLOTYPE, Adult. Reg. no. 1889.7.1.60. Kina Balu Mountain, Borneo, 15 Feb. 1887. Collected and presented by J. Whitehead. The specimen lacks a tail.
cinereifrons—Brachypus cinereifrons Tweeddale, 1878
cinereigulare—Todirostrum cinereigulare P. L. Sclater 1857 9 Proc. zool. Soc. Lond., 1856 (publ. 1857): 295. The type was a specimen collected by A. Sallé in Cordova, Vera Cruz. The specimen collected by Boucard from Oaxaca and listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 77, cannot be accepted.
cinereigularis—Dicaeum cinereigularis Tweeddale, 1878 45 Proc. zool. Soc. Lond., 1877 (publ. 1878): 829. SYNTYPE, Adult male. Reg. no. 1888.7.12.78. Butuan, Mindanao, Philippines, May 1877. Collected by A. O. Everett. Tweeddale Collection. This specimen was labelled as the type by the author but not designated. There are several other syntypes in the collection.
cinereiventris—Myrmotherula cinereiventris P. L. Sclater and Salvin, 1867

по. 1889.9.20.315.	Cayenne, French Guiana.	Sclater Collection
Listed as type by	Sclater, Cat. Bds. Brit. M	us., 15 (1890): 244.
There are several ot	her syntypes in the collection	n.

- cinereiventris—Pachyrhamphus cinereiventris Sclater, 1862 Catalogue of a collection of American birds: 242. HOLOTYPE, Adult male. Reg. no. 1888.1.13.1497. Santa Marta, Colombia. Collected for Verreaux. Sclater Collection.
- cinereocapilla—Prinia cinereocapilla Moore, 1854 in Horsfield and Moore, Catalogue of birds in the Museum of the Honourable East India Company, 1: 322. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.3061. (Hodgson's no. 890.) Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company Museum.

The specimen has a broken bill. Moore used a Hodgson MS.

name.

cinereoniger—Thamnophilus cinereoniger Pelzeln, 1869. Zur Ornithologie Brasiliens, : 76. 143. SYNTYPE, Adult male. Reg. no. 1889.9.20.51. Rio Amajau, Brazil, 16 Sept. 1831. Collected by J. Natterer. P. L. Sclater Collection.

This specimen and two others in the collection were listed as "authentic specimens" by Sclater, Cat. Bds. Brit. Mus., 15 (1890):

193.

- cinereus—Erythropygia coryphaeus cinereus Macdonald, 1952 Bull. Br. Orn. Cl., 72:91. HOLOTYPE, Adult female. Reg. no. 1950.50.355. 16 miles north of Port Nolloth, 50', Little Namaqualand, 19 Dec. 1949. Collected and presented by the South-West Africa Expedition.
- cinereus—Euprinodes cinereus Sharpe, 1891 *Ibis*, (6) 3:120. SYNTYPE, Adult male. Reg. no. 1893,12.1.21. Mount Elgon, north-east of Lake Victoria Nyanza, (25 Feb. 1890. vide Sharpe, Ibis, (6) 4 (1892): 155) Collected and presented as the type by F. J. Jackson, but a second male specimen was also available.
- cinereus—Malacopteron cinereus Eyton, 1839 Proc. zool. Soc. Lond., 1839: 103. Specimen Brit. Mus. Reg. no. 1881.2.18.24. is marked "original spec." but does not agree with the type-description and was not accepted as typical material by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 566.
- cinnamomeoventris—Iole virescens cinnamomeoventris E. C. S. Baker, Bull. Br. Orn. Cl., 38: 16. SYNTYPE, Adult male. Reg. no. 1886.9. 1.1081. Tenasserim Town, 25 Nov. 1874. Collected by W. Davison.

This is the author's selected male. His selected female syntype is also in the collection

- cinnamomeum—Trochalopterum cinnamomeum Davison, 1886. . 32B

 Ibis, (5) 4:204. Syntype, Adult. Reg. no. 1886.7.31.2. Hills
 to east of Cannanore, Travancore, India. Collected for A.
 Macgregor and presented by the Trivandrum Museum, Travancore.
- cinnamomeus—Cinclosoma cinnamomeus Gould, 1846 32A Proc. zool. Soc. Lond., 1846 : 68. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1846.5.22.87. The Depot, lat. 29° 40′ S., central Australia, 9 June 1845 (i.e., near Milparinka, north-west New South Wales, vide Whittell, The Literature of Australian Birds, 1954 : 694). Collected and presented by Captain Charles Sturt.
- cinnamomeus—Drymocataphus cinnamomeus Rippon, 1900 . . . 32B Bull. Brit. Orn. Cl., 11:12. Holotype, Adult. Reg. no. 1900. 12.20.286. Loi Mai, 6,000′, South Shan States, 15 April 1899. Collected and presented by G. Rippon.

- citriniventris—Emberiza citriniventris P. L. Sclater, 1905 . . . 49

 Bull. Brit. Orn. Cl., 16: 39. HOLOTYPE, Adult male. Reg. no. 1905.10.11.2. Kuryatein, Syria, 8 April 1905. Collected by and purchased of D. Carruthers.
- citrinus—Megalurus citrinus G. R. Gray and Mitchell, 1848. . . 35

 The Genera of Birds, 1: pl. 48. Holotype, Adult. Reg. no. Old
 Vellum Cat. no. 412b (Vol. 14). Java. Collected by T. Horsfield.
 Presented by the Honourable East India Company Museum.
- clanceyi—Emberiza calandra clanceyir R. Meinertzhagen, 1947 . 49
 Bull. Brit. Orn. Cl., 67:91. HOLOTYPE, Adult female. Reg. no. 1965.M.38. Aran Island, County Galway, West Ireland, 7 Feb. 1947. Meinertzhagen Collection.

Clarkei—Coccopygia clarkei Shelley, 1903
clarkei—Turdoides leucopygia clarkei Macdonald, 1939 32B Bull. Br. Orn. Cl., 60:10. HOLOTYPE, Adult male. Reg. no. 1938.5.18.1. Gummaro, near Gore, west Abyssinia, 11 July 1917. Collected by O. Kovacs and presented by Johannes Effendi.
clarkei—Turdus philomelos clarkei Hartert, 1909 31 Bull. Br. Orn. Cl., 23:54. HOLOTYPE, Adult male. Reg. no. 1936.10.15.12. Tring Park, Hertfordshire, England, 16 May 1902. Collected by E. Hartert and presented by the American Museum of Natural History, New York, from the Rothschild Collection.
clarki—Ixulus clarki Oates, 1894
clarus—Turdinulus epilepidotus clarus Robinson and Kloss, 1919 . 32B Ibis, (11) 1:582. Syntype, Adult male. Reg. no. 1919.12.20.292. Dalat, 5,000', south Annam, 3 May 1918. Collected by C. B. Kloss and presented by H. C. Robinson and Kloss. This is the authors' selected male. Their selected female syntype is also in the collection.
claudei—Apalis claudei W. L. Sclater, 1910
claudei—Crateropus tenebrosus claudei Bannerman, 1919 32B Bull. Br. Orn. Cl., 39:99. HOLOTYPE, Adult male. Reg. no. 1919.10.12.150. Poko, Welle River, north Belgian Congo, 13 Aug. 1914. Collected by C. Christy and presented by the Musee Royale D'Afrique Centrale, Tervuren, Belgium.
claudei—Lioptilus claudei Boyd Alexander, 1902 32B Bull. Br. Orn. Cl., 13: 34. SYNTYPE, Adult male. Reg. no. 1911. 12.23.2345. Near Bakaki, Fernando Po (on label), 17 Nov. 1902. Collected and bequeathed by Boyd Alexander. The locality in the text, Mount St. Ysabel, is that of the female syntype, in the collection, collected on 26 Nov.; vide Boyd Alexander, Ibis, (8) 3 (1903): 382.
claudi—Thamnolaea claudi Boyd Alexander, 1906 31 Bull. Br. Orn. Cl., 16: 124. HOLOTYPE, Adult female. Reg. no. 1936.2.21.77. Pettia, Lake Chad, central Africa, 25 Aug. 1904. Collected by Boyd Alexander. Boyd Alexander Collection; purchased of his nephew.

Ibis, (10) 2, Jubilee Supplement: 16, and text figures 1 and 2. SYNTYPE, Adult male. Reg. no. 1916.5.30.1140. Parimau, Mimika River, Dutch New Guinea, 18 Nov. 1910. Collected by C. H. B. Grant. Presented by British Ornithologists' Union New Guinea Expedition. This is the author's selected male, his selected female and several other syntypes are also in the collection.
cleaveri—Drymocataphus cleaveri Shelley, 1874
cleghorniae—Franklinia cleghorniae Blyth, 1867 35 Ibis, (2) 3: 24. A single specimen was "made over to Mr Gould". This specimen, from the banks of the Ganges, was listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 247, but cannot now be traced.
clio—Pachycephala clio Wallace, 1862
cobanensis—Vireosylvia cobanensis P. L. Sclater, 1860
cobbi—Troglodytes cobbi Chubb, 1909
coburni—Turdus coburni Sharpe, 1901
cockburniae—Anthus cockburniae Oates, 1890
cockerelli—Philemon cockerelli P. L. Sclater, 1877 48 Proc. zool. Soc. Lond., 1877: 104. HOLOTYPE, Adult. Reg. no. 1878.3.14.5. New Britain, Feb. 1876. Collected by the Rev. G. Brown. Purchased of E. Gerrard.

cockerelli—Ptilotis cockerelli Gould, 1869
codea—Alauda codea Smith, 1843
codringtoni—Hypochaera codringtoni Neave, 1907 61 Mem. Proc. Manchester Lit. Phil. Soc., 51 (10): 94. HOLOTYPE, Adult male. Reg. no. 1907.12.29.140. Molilo's, near Petauke, (East Loangwa Valley, on label), Northern Rhodesia, 3 Feb. 1905. Collected by S. A. Neave. Presented by the British South Africa Company.
coelestis—Hypothymis coelestis Tweeddale, 1877
coelica—Erythrura prasina coelica E. C. S. Baker, 1925 60 Bull. Brit. Orn. Cl., 45:84. HOLOTYPE, Adult male. Reg. no. 1888.8.13.42. Marintan-an, Borneo, Feb. 1888. Collected by A. H. Everett. Purchased of E. Gerrard.
coelicolar—Grandala coelicolar Hodgson, 1843
coelicolor—Calliste coelicolor P. L. Sclater, 1851
coelivox—Alauda coelivox Swinhoe, 1859

- coeruleotincta—Lioptila pulchella coeruleotincta Rothschild, 1921. 32B Novit. Zool., 28:38. SYNTYPE, Adult male. Reg. no. 1921.7.15. 291. Tengyueh District, Shweli-Salwin Divide, 6,000-7,000', Yunnan, May 1919. Collected by G. Forrest and presented by Stephenson Clarke.

 The three other syntypes are also in the collection.
- coerulifrons—Cyornis magnirostris coerulifrons E. C. S. Baker, 1918 37 Bull. Brit. Orn. Cl., 39:8. HOLOTYPE, Adult male. Reg. no. 1916.12.27.912. Klong Bang Lai, Siam, 24 Jan. 1916. Collected by C. Chunggat for E. G. Herbert. Presented by E. G. Herbert.
- cognita—Cryptolopha burkii cognita La Touche, 1922 35

 Bull. Brit. Orn. Cl., 43: 42. SYNTYPE, Adult female. Reg. no. 1900.1.18.84. Kautun, North-west Fohkien, China, 6 April 1898. Collected and presented by C. B. Rickett.

 According to Bangs, Bull. Mus. Comp. Zool. Harvard, 70 (1930): 340. the male syntype is at Harvard.
- colimae—Buarremon virenticeps colimae van Rossem, 1938 . . . 49
 Bull. Brit. Orn. Cl., 58: 125. Holotype, Adult male. Reg. no. 1894.7.1.972. Sierra Nevada de Colima, 8,000-12,000 ft., Jalisco, Mexico, 8 April, 1889. Collected by W. B. Richardson. Salvin-Godman Collection
- colimae—Junco phaeonotus colimae van Rossem, 1938 49
 Bull. Brit. Orn. Cl., 58: 132. HOLOTYPE, Adult male. Reg. no.
 1899.2.1.2255. Sierra Nevada de Colima (Jalisco, on label), Mexico,
 9 Apr. 1889. Collected by W. Lloyd and W. B. Richardson.
 Salvin-Godman Collection.
- colimae—Troglodytes brunneicollis colimae Van Rossem, 1938 . 28
 Bull. Br. Orn. Cl., 59:14. HOLOTYPE, Adult male. Reg. no. 1890.12.20.568. Sierra Nevada de Colima, 12,000', south Jalisco, Mexico, 5 Dec. 1889. Collected by W. B. Richardson. Salvin-Godman Collection.

collaris—Ploecus collaris J. E. Gray, 1831
collaris—Ploceus collaris Fraser, 1842
colletti—Poliopsar colletti Sharpe, 1888
colliei—Pica colliei Vigors, 1829
collingwoodi—Rhynchocyclus flaviventris collingwoodi Chubb, 1920 9 Bull. Brit. Orn. Cl., 40: 109. Holotype, Adult. Reg. no. 1909. 8.7.168. Macqueripes Valley, Trinidad (Tuches Valley, on label)., 27 Dec. 1905. Collected and presented by H. Collingwood Ingram.
collsi—Erythropygia collsi Boyd Alexander, 1907 31 Bull. Br. Orn. Cl., 19:46. HOLOTYPE, Adult male. Reg. no. 1911.12.23.1880. Libokwa, Welle River (Joga River on label), north Belgian Congo, 8 March 1906. Collected and bequeathed by Boyd Alexander.
colombiana—Spermophila colombiana Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:99. HOLOTYPE, Adult male. Reg. no. 1885.12.14.520. Bogota, Colombia. Collected by G. Crowther. Salvin-Godman Collection.
commixtus—Parus commixtus Swinhoe, 1868

comorensis—Artamia comorensis Shelley, 1894
comorensis—Zosterops comorensis Shelley, 1900
compressirostris—Myiolestes compressirostris Layard, 1876 38 Ibis, (3) 6:153. HOLOTYPE, Adult. Reg. no. 1897.6.1.625. Kandavu, Fiji Islands. Collected by C. Pearce. Seebohm Bequest.
comptus—Buarremon comptus P. L. Sclater and Salvin, 1879 49 Ibis, (4) 3: 426. Syntype, Adult. Reg. no. 1885.5.8.434. Maravina, Ecuador. 1876. Collected by C. Buckley. Salvin-Godman Collection. A second syntype is also in the collection.
comrii—Manucodia comrii Sclater, 1876
concinens—Calamoherpe concinens Swinhoe, 1870 35 Proc. zool. Soc. Lond., 1870 : 432. HOLOTYPE, Adult female. Reg. no. 1898.9.1.1274. Pekin, 2 July 1868. Collected by R. Swinhoe. Seebohm Bequest.
concinna—Euphonia concinna P. L. Sclater, 1855
concinna—Pitta concinna Gould, 1857
concinna—Psaltria concinna Gould, 1855

There are three other syntypes in the collection. Gould, *Proc. zool. Soc. Lond.*, 1837: 22, used the same name for a different bird, apparently unaware of Sykes' description, but subsequently renamed his species *Progne modesta* Gould, 1839.

The name is preoccupied by *H. concolor* Sykes, 1832 (q.v.), and was renamed *Progne modesta* by Gould, *Zoology of the voyage of H.M.S. Beagle*, 3 *Birds* (1839): 39, although no reason was given.

- concolor—Horeites acanthizoides concolor Ogilvie-Grant, 1912 . 35

 Bull. Brit. Orn. Cl., 29: 107. SYNTYPE, Adult male. Reg. no.
 1913.1.29.40. Mount Arizan, 8,000', Formosa, April 1912. Collected by W. Goodfellow. Purchased of E. J. Johnstone.

 The female syntype is also in the collection.
- condorensis—Mixornis rubricapilla condorensis Robinson, 1921. 32B J. nat. Hist. Soc. Siam, 4:88. HOLOTYPE, Adult male. Reg. no. 1955.1.5989. Pulo Condore, coast of Cochin China, 8 Sept. 1919. Collected for and presented by Sir W. J. F. Williamson.

congica—Lagonisticta congica Sharpe, 1890	60 alt female.
Reg. no. 1887.8.12.11. Kassongo, Upper Congo, Ju lected by and purchased of F. Bohndorff.	ine. Col-
congicus—Dryoscopus congicus Sharpe, 1901 in Jackson, Ibis, (8) 1:39. SYNTYPE, Adult female. 1878.2.23.64. St. Antoine, Lower Congo River. Collect Lucan and purchased of A. Bouvier. The other syntype is also in the collection.	
congicus—Erythrocercus congicus Ogilvie-Grant, 1907. Bull. Brit. Orn. Cl., 19: 41. Adult male. Reg. no. 1906.1 (Irumu, Ituri Forest, vide Ogilvie-Grant, Trans. zool. St. 19 (1910): 403), 3,000′, Eastern Congo Forest, 16 Collected by R. E. Dent. Presented by the subscribe Ruwenzori Expedition.	oc. Lond., Oct. 1906.
congo—Cisticola pipiens congo Lynes, 1936 Bull. Brit. Orn. Cl., 56:110. HOLOTYPE, Adult male. 1936.4.13.2. Elizabethville, 4,000', South-east Congo, 1934. Collected by Admiral H. Lynes and J. Vincent. by Admiral H. Lynes.	, 10 Jan.
congrevei—Parus palustris congrevei Kinnear, 1928. Bull. Brit. Orn. Cl., 49:50. Holotype, Adult male. 1928.6.16.32. Reytezet, 3,400', (near Guraylata, in Transylvania, 14 April 1928. Collected and presented Congreve.	42 Reg. no. register), by W. M.
conirostris—Alauda conirostris Sundevall, 1850 K. svenska Vetensk-Akad. Handl. Stockholm, 7:99. Adult female. Reg. no. 1878.12.31.734. Megaliesber Transvaal, 28 Oct. 1844. Collected by J. Wahlberg. Stlection; received by Sharpe in exchange with the Sundevalum. Listed as "one of the typical specimens" by Sharpe, In Soc. Lond., 1874:631.	rg Range, narpe Col- Stockholm
connectens—Alophoixus phaeocephalus connectens Chasen a 1929	21 no. 1947. eo, 6 July

connectens—Vireo gilvus connectens van Rossem, 1940.

Salvin-Godman Collection.

Trans. San Diego Soc. Nat. Hist., 9:77. HOLOTYPE, Immature, Reg. no. 1891.11.3.244. Chilpancingo, 4,600', Sierra Madre del Sur, Guerrero, Mexico, Aug. 1888. Collected by D. W. Smith.

- consanguineus—Analcipus consanguineus Wardlaw Ramsay, 1881 . 63 Ibis, (4) 5:33, pl. 1. SYNTYPE, Adult male. Reg. no. 1887.11.20. 627. Paio, Sumatra. Collected by Bock. Tweeddale Collection. There are several other syntypes in the collection. consobrina—Formicivora consobrina P. L. Sclater, 1860 Proc. zool. Soc. Lond., 1860: 279. SYNTYPE, Adult male. Reg. no. 1889.9.20.372. Babahoya, Ecuador, Aug. 1859. Collected by L. Fraser. Sclater Collection. Listed as type by Sclater, Catalogue of American Birds, 1862: 183. The female syntype is in the collection. consobrinus—Aegithalus consobrinus Swinhoe, 1870 41 Proc. zool. Soc. Lond., 1870:133. HOLOTYPE, Adult male. Reg. no. 1898.9.20.456. Sha-she, on Ichang River, Hupeh, China, 4 April 1869. Obtained by Swinhoe as a captive specimen. Seebohm Bequest. consobrinus—Oriolus consobrinus Wardlaw Ramsay, 1879 Proc. zool. Soc. Lond., 1879: 907. HOLOTYPE, Adult female. Reg. no. 1887.11.20.526. Sandakan, North Borneo. Collected by Cutter. Tweeddale Collection. consobrinus—Philydor consobrinus P. L. Sclater, 1870 . Proc. zool. Soc. Lond., 1870: 328. SYNTYPE, Adult. Reg. no. 1889.5.20.358. Bogota, Colombia, 1869. Collected for Verreaux. Sclater Collection. There is another syntype in the collection. conspicillata—Microeca conspicillata G. R. Gray, 1859 Proc. zool. Soc. Lond., 1859: 156. SYNTYPE, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 222). Reg. no. 1859.418.32. Dorey, New Guinea. Collected by A. R. Wallace. Purchased of Stevens. A second syntype is also in the collection. conspicillatus—Basileuterus conspicillatus Salvin and Godman, 1880 54 Ibis, (4) 4:117. SYNTYPE, Adult female. Reg. no. 1885.3.8.661. San José, Santa Marta, Colombia. 8 June 1879. Collected by F. Simons. Salvin-Godman Collection. The second syntype is also in the collection. continentalis—Emberiza yessoënsis continentalis Witherby, 1913 . 49 Bull. Brit. Orn. Cl., 31:74. HOLOTYPE, Adult male. Reg. no. 1924.12.20.6. Yashwanen, near Nanking, China, 31 Jan. 1911. Collected by H. Lynes. Presented by H. F. Witherby. continentis—Calyptomena viridis continentis Robinson and Kloss,

30 cooki
Robinson and C. B. Kloss, and presented by the Raffles Museum, Singapore. There is a female syntype in the collection.
ooki—Cyanopica cooki Bonaparte, 1851
ooki—Suya crinigera cooki Harington, 1913
oquerellii—Nectarinia coquerellii Hartlaub, 1860
Proc. zool. Soc. Lond., 1856: 29. SYNTYPE, Adult male. Reg. no. 1888.1.13.1268. Bogota, New Colombia. Collected for M. Verreaux. Sclater Collection Sclater used a Verreaux MS. name. The immature and female syntypes are also in the collection.
Proc. zool. Soc. Lond., 1891: 133, pl. 13. SYNTYPE, Adult. Reg. no. 1896.12.28.1. 8 leagues from Sacaya, Tarapacá, Chile, 20 March 1890. Collected by A. A. Lane. Presented by H. Berkeley James. Listed as the type by Hellmayr and Conover, Catalogue of birds of the Americas, 11 (1938): 350. The second syntype is also in the collection.

corallina—Sitta corallina Hodgson, 1836 J. Asiat. Soc. Bengal, 5: 779. SYNTYPE, Adult (female, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883): 359). Reg. no. 1843.1.13.254. Nepal. Collected and presented by B. H. Hodgson. Listed as the type by Gadow (op. cit.).

cordofanica—Mirafra cordofanica Strickland, 1850 Proc. zool. Soc. Lond., 1850: 218, pl. 23. HOLOTYPE, Adult (female on label). Reg. no. 1851.4.16.4. Kordofan, east Sudan. Collected by J. Petherick and purchased of H. A. Strickland.

22D

Ibis, (8) 5: 330, pl. 7. HOLOTYPE, Adult (referred to in text as both male and female). Reg. no. 1916.9.20.1170. Jebel Melbis, Kordofan, Sudan, 16 April 1904. Collected and presented by A. L. Butler.
corea—Sitta corea Ogilvie-Grant, 1906
Cornix—Graucalus cornix Reichenow, 1900
coronata—Todopsis coronata Gould, 1878
coronatus—Dryoscopus coronatus Sharpe, 1874 23 Proc. zool. Soc. Lond., 1874 : 205, pl. 33. Holotype, Adult male. Reg. no. 1874.2.17.7. Danger River, Gaboon. Collected and presented by H. T. Ansell.
coronatus—Lanius coronatus Raffles, 1822
coronatus—Malimbus coronatus Sharpe, 1906 61 Bull. Br. Orn. Cl., 19:18. HOLOTYPE, Adult male. Reg. no. 1908.5.25.100. Ja River, Cameroons, 18 Feb. 1906. Collected by and purchased of G. L. Bates.
coronatus—Malurus coronatus Gould, 1857

of A. C. Gregory.

This is the author's described male, the described female syntype is also in the collection. See also, Chisholm, Queensland Naturalist, 17 (1964): 65.

102 Colonatas
coronatus—Orthotomus coronatus Blyth and Jerdon, 1861 35 Proc. zool. Soc. Lond., 1861: 200. SYNTYPE, Adult. Reg. no. 1862.1.17.27. Darjeeling, Sikkim. Collected and presented by T. C. Jerdon. There are two other syntypes in the collection.
coronatus—Platyrhynchus coronatus P. L. Sclater, 1858 9 Proc. zool. Soc. Lond., 1858:71. HOLOTYPE, Adult. Reg. no. 1888.1.13.230. Napo River, Ecuador. Collected for Verreaux. Sclater Collection.
coronoides—Corvus coronoides Vigors and Horsfield, 1827
coronulatus—Masius coronulatus P. L. Sclater, 1860 8 Proc. zool. Soc. Lond., 1860: 91. HOLOTYPE, Adult male. Reg. no. 1888.1.13.1241. Nanegal, 4,000', western shore of Pichincha, north of Quito, west Ecuador. Collected by L. Fraser. Sclater Collection.
coruscans—Neodrepanis coruscans Sharpe, 1875
corvina—Spermophila corvina P. L. Sclater, 1860
corvinus—Thamnophilus corvinus P. L. Sclater, 1855 4 Edinb. New philos. J., (n.s.) 1:234. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1855.11.5.7. Ucayali River, east Peru, June 1852. Collected by J. Hauxwell and purchased of J. Gould. Sclater used Gould's MS name.
costaricensis—Pachyramphus costaricensis Chubb, 1920 7 Bull. Brit. Orn. Cl., 40:74. HOLOTYPE, Adult male. Reg. no. 1899.5.1.428. Bebedero, Costa Rica, 15 Sept. 1893. Collected by C. F. Underwood. Salvin-Godman Collection.
costaricensis—Thryophilus costaricensis Sharpe, 1881

cotta—Elania cotta Gosse, 1849 Ann. Mag. nat. Hist., (2) 3:257. SYNTYPE, Adult (relaxed mount). Reg. no. 1847.6.8.10. Jamaica. Collected by and purchased (?) of P. H. Gosse (register states that this particular specimen was 'purchased of Golding"). This specimen was purchased as the type, and listed as the type by Sharpe, Hist, Coll. Brit. Mus., 2, Birds, (1906): 373. There are two other syntypes in the collection. couesi—Campylorhynchus couesi Sharpe, 1881 28 Cat. Bds. Brit. Mus., 6: 196. SYNTYPE, Adult female. Reg. no. 1880.11.24.6. Larido, Texas, 28 Feb. 1867. Collected by H. B. Butcher and presented by the Smithsonian Institution. This was the only specimen in the collection at the time of description. Other syntypes were presented later with the Salvin-Godman Collection. cowani—Cotile cowani Sharpe, 1882 . 18 J. Linn. Soc. Lond., 16: 322. HOLOTYPE, Adult female, Reg. no. 1882.2.27.70. Forest of Ankáfana, Betsileo, south-east Madagascar, 11 March 1881. Collected by and purchased of the Rev. D. Cowan. coxi—Ammomanes deserti coxi R. Meinertzhagen, 1923 17 Bull. Br. Orn. Cl., 44:15. HOLOTYPE, Adult male. Reg. no. 1905.10.11.46. Karyatein, 2,000', north Syrian desert, 24 Mar. 1905. Collected by D. Carruthers and J. H. Miller and purchased of the former. cozumelae—Xanthoura luxuosa cozumelae Van Rossem, 1924 Bull. Mus. comp. Zool., Harvard, 77:397. HOLOTYPE, Adult. Reg. no. 1886.9.9.862. Cozumel Island, Yucatan (off coast of Quintana Roo, Mexico, on label), Jan. 1866. Collected by G. F. Gaumer. Salvin-Godman Collection. craddocki—Suthora craddocki Bingham, 1903 Adult male. Reg. no. 1903.12.24.232. Loi-Pang-Nan, 8,500', east of Kengtung, valley of Mekong River, South Shan States, Upper Burma (2 April 1902, vide Bingham, Ibis, loc. cit.: 587). Collected by H. N. Thompson and W. H. Craddock and purchased through E. Gerrard. cranbrooki—Grammatoptila striata cranbrooki Kinnear, 1932 Bull. Br. Orn. Cl., 53:79. HOLOTYPE, Adult female. Reg. no. 1932.12.10.2. Adung Valley, 6,000', 28°10' N., 97°40' E., north-east Burma, 16 April 1931. Collected and presented by Lord Cranbrook and F. Kingdon Ward. cranbrooki—Mirafra cranbrooki Boyd Alexander, 1907 Bull. Br. Orn. Cl., 19:88. SYNTYPE, Adult male. Reg. no. 1911. 12.23.3003. Bwando, River Welle, Congo, 25 Dec. 1905. Col-

Bequest. The female syntype is also in the collection. Boyd Alexander Expedition. Boyd Alexander Expedition.
crassa—Calornis crassa P. L. Sclater, 1883 62 Proc. zool. Soc. Lond., 1833: 56, pl. 14. SYNTYPE, Adult male. Reg. no. 1883.5.30.79. Larat, Timor Laut, Tenimber Island Group, 1 Aug. 1882. Collected by W. O. Forbes. Presented by the Timor Laut Committee of the British Association. The female syntype is also in the collection.
crassirostris—Calornis crassirostris Walden, 1872 62 Trans. zool. Soc. Lond., 8:80. HOLOTYPE, Adult. Reg. no. 1888.9.20.422. Lombock. Collected by A. R. Wallace (vide specimen label). Tweeddale Collection.
crassirostris—Camarhynchus crassirostris Gould, 1837 53 Proc. zool. Soc. Lond., 1837: 6. The specimen listed by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 16, is not the type, vide Swarth, Occ. Pap. Calif. Acad. Sci., 18 (1931): 209.
crassirostris—Carpodacus crassirostris Blyth, 1847
crassirostris—Euphonia crassirostris P. L. Sclater, 1857 51 Proc. zool. Soc. Lond., 1856 (publ. 1857): 277. SYNTYPE, Adult. Reg. no. 1857.11.18.5. (recently assigned). Bogota, Colombia. Presented by P. L. Sclater. Sclater, Cat. Bds. Brit. Mus., 11 (1886): 77, lists a specimen from Santa Marta as the type, apparently in error.
crassirostris—Geositta crassirostris P. L. Sclater, 1866
crassirostris—Myiophoneus crassirostris Robinson, 1910 31 Bull. Br. Orn. Cl., 25:99. HOLOTYPE, Adult male. Reg. no. 1910.12.27.271. Kokhan, Trang, northern Malay Peninsula, 12 Jan. 1910. Presented by the Kuala Lumpur Museum.
crassirostris—Pitta crassirostris Wallace, 1862

crassirostris—Pyromelana crassirostris Ogilvie-Grant, 1907 61 Bull. Br. Orn. Cl., 21:14. Holotype, Adult male. Reg. no. 1906.12.23.241. North end of Ruwenzori, 3,500′, 19 Aug. 1906. Collected by R. E. Dent. Presented by the subscribers to the Ruwenzori Expedition.
craterum—Laniarius atroflavus craterum Bates, 1926 23 Bull. Br. Orn. Cl., 46:88. HOLOTYPE, Adult male. Reg. no. 1926.8.8.2. Oku, 6,000′, west of Kumbo, Cameroons, 11 Feb. 1925. Collected by and purchased of G. L. Bates.
crawfordi—Dendroeca crawfordi Nicoll, 1904 54 Bull. Brit. Orn. Cl., 14: 95. SYNTYPE, Adult ?female. Reg. no. 1904.8.17.116. Little Cayman Island, West Indies, (14 Mar. 1904 on label). Collected by M. J. Nicoll. Presented by the Earl of Crawford, from the voyage of R. Y. S. Valhalla. This specimen was marked "type" and "figured in Ibis", (8) 4 (1904), pl. 11. There are two other syntypes in the collection.
crawfurdi—Eremomela crawfurdi Clarke, 1911
creaghi—Amydrus creaghi Ogilvie-Grant and Forbes, 1903 62 The Natural History of Sokotra and Abd-el-Kuri: 58, HOLOTYPE, Adult male. Reg. no. 1899.8.11.143. Abd-el-Kuri Island, 22 Feb. 1899. Collected by W. R. Ogilvie-Grant and H. O. Forbes. Presented by the Royal Society.
crepitans—Muscicapa crepitans Latham, 1801
criniger—Suya criniger Hodgson, 1836
crissalis—Fringilla crissalis Vigors, 1839

crissalis—Helminthophilia crissalis Salvin and Godman, 1889 . 54 Ibis, (6) 1:380. HOLOTYPE, Adult female. Reg. no. 1892.4.1.195. Sierra Nevada de Colima, 8,000', Jalisco, Mexico, 6 April 1889. Collected by W. Lloyd and W. B. Richardson. Salvin-Godman Collection.
crissalis—Pheucticus crissalis P. L. Sclater, 1877
crissalis—Strepera crissalis Sharpe, 1877
crissalis—Zosterops crissalis Sharpe, 1844
cristatellus—Tachyphonus cristatellus P. L. Sclater, 1862 51 Catalogue of a collection of American birds: 86. HOLOTYPE, Adult male. Reg. no. 1885.6.12.756. Bogota, Colombia. (Collected by Parzudaki, on label.) Sclater Collection.
croceus—Pericrocotus croceus Sharpe, 1888
crocopygia—Poliospiza crocopygia Sharpe, 1871
crocotius—Paradoxornis poliotis crocotius Kinnear, 1954
crossensis—Chalcomitra rubescens crossensis Serle, 1963 46 Bull. Brit. Orn. Cl., 83:118. Holotype, Adult male. Reg. no. 1963.12.1. Mamfe, on Cross River, Federated Republic of Cameroon, 10 March 1953. Collected and presented by W. Serle.

crossleyi—Turdus crossleyi Sharpe, 1871
cruralis—Brachypteryx cruralis Hodgson, MS
cruralis—Megalurus cruralis Vigors and Horsfield, 1827 35 Trans. Linn. Soc. Lond., 15: 228. SYNTYPE, Adult. Reg. no. 1863.7.7.29b. Australia. Collected by G. Caley. Presented by the Linnean Society of London. Another syntype is in the collection.
crypterythrus—Myiobius crypterythrus P. L. Sclater, 1861 9 Proc. zool. Soc. Lond., 1860 (Publ. 1861): 464. SYNTYPE, Adult male. Reg. no. 1888.1.13.875. Pallatanga, Ecuador, Oct. 1858. Collected by L. Fraser. Sclater Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 210. There are several other syntypes in the collection.
cryptoleuca—Myrmecocichla cryptoleuca Sharpe, 1891 31 Ibis, (6) 3:445. HOLOTYPE, Adult. Reg. no. 1893.12.1.28. Kikuyu, central East Africa, 14 Sept. 1894. Collected and presented by F. J. Jackson.
cryptolopha—Lathria cryptolopha P. L. Sclater and Salvin, 1877 . 7 Proc. zool. Soc. Lond., 1877: 522. HOLOTYPE, Adult. Reg. no. 1888.1.20.599. Mongi, eastern Ecuador, Mar. 1877. Collected by C. Buckley. Salvin-Godman Collection.
cryptoxanthus—Myiobius cryptoxanthus P. L. Sclater, 1861 9 Proc. zool. Soc. Lond., 1860 (publ. 1861): 465. SYNTYPE, Adult male. Reg. no. 1888.1.13.878. Zamora, Ecuador, Jan. 1858. Collected by L. Fraser. Sclater Collection. The female syntype is also in the collection.
cucullata—Conopophaga cucullata P. L. Sclater, 1856 4 Proc. zool. Soc. Lond., 1856 : 29, pl. 119. HOLOTYPE, Adult. Reg. no. 1889.9.20.654. Bogota, Columbia. Collected for Verreaux.
cucullata—Crypsirhina (Temia) cucullata Jerdon, 1862 71 Ibis, (1) 4:20. Syntype, Adult. Reg. no. 1862.1.17.12. Theyetmo, Upper Burma. Collected and presented by T. C. Jerdon. There is another syntype in the collection.
cucullata—Eopsaltria cucullata G. R. Gray, 1859 39 Catalogue of birds of tropical islands of the Pacific : 21. HOLOTYPE, Adult (female, on label). Reg. no. 1859.6.14.14.

Anieteum, New Hebrides, (26 Oct. 1858, on label). Collected by J. Macgillivray on the voyage of H.M.S. Rattlesnake. Purchased of S. Stevens.

cucullata—Geothlypis cucullata Salvin and Godman, 1889 . Ibis, (6) 1:237. SYNTYPE, Adult male. Reg. no. 1892.4.20.1526. Cofre de Perote, Jalape, Mexico, July 1888. Collected by M. Trujillo. Salvin-Godman Collection.

There are two other syntypes in the collection. This species renamed G. nelsoni C. W. Richmond Auk 17 (1900): 179, on the

grounds of preoccupation by Sylvia cucullata Latham 1790.

cucullata—Urocissa cucullata Gould 1861 Birds of Asia 5, pt. 13 (May 1861): pl. 51. SYNTYPE, Adult. Reg. no. 1876.1.7.89. Kulu or Kooloo Valley, 50 miles north of Simla, India. Collected by Captain M. Tweedie. Purchased of J. Gould. Gould Collection.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 3 (1877): 74.

cucullatus—Tarsiger cucullatus Blyth, 1867. Ibis, (2) 3:16. SYNTYPE, Adult male. Reg. no. 1867.3.16.90. "Africa or India". Gould Collection.

The second syntype is also in the collection. Blyth used Gould's

MS name.

- culequita—Troglodytes brunneicollis culequita van Rossem, 1938 . 28 Bull. Br. Orn. Cl., 59:13. HOLOTYPE, Adult. Reg. no. 1890.12. 20.548. Coajimalpa (Tacubaya), Mexico, 7 March 1888. Collected by F. Ferrari-Perez. Salvin-Godman Collection.
- culminatus—Corvus culminatus Sykes, 1832. Proc. zool. Soc. Lond., 1832:96. SYNTYPE, Adult. Reg. no. 1860.4.16.543. Deccan, Collected by W. H. Sykes. Presented by the Honourable East India Company Museum.
- cumingi—Minodoria striaticeps cumingi Hachisuka, 1934 Tori, 8, no. 38:221. HOLOTYPE, Adult (relaxed mount). Reg. no. 1842.2.15.1. Philippines ("South of the Island of Manilla" in register). Collected by and purchased of H. Cuming.
- cummingi—Saxicola cummingi Whitaker, 1899 31 Bull. Br. Orn. Cl., 10:17. HOLOTYPE, Adult. Reg. no. 1885.3.7.81. Fao, Persian Gulf. Collected and presented by W. D. Cumming.
- cunhaci—Mesia argentauris cunhaci Robinson and Kloss, 1919 Ibis, (11) 1:591, pl. 16. HOLOTYPE, Adult male. Reg. no. 1919. 12.20.345. Dalat, 5.000', south Annam, 11 April 1919. Collected and presented by C. B. Kloss.
- cupreonitens—Nectarinia cupreoniteus Shelley, 1876 46 Monograph of the Nectariniidae, : 17, pl. 6. HOLOTYPE, Adult male. Reg. no. 1874.5.1.11. Abyssinia, Aug. 1850. Collected for J. Verreaux. Sharpe Collection.

curtata—Synallaxis curtata P. L. Sclater, 1869 Proc. zool. Soc. Lond., 1869: 636, pl. 49. SYNTYPE, Adult. Reg. no. 1889.5.20.220. Bogota, Colombia, Collected by Trübner, Sclater Collection. There is another syntype in the collection, but only the selected specimen agrees with the plate. curvirostris—Limnornis curvirostris Gould, 1839 Zoology of the Voyage of H.M.S. Beagle, 3 Birds: 81, pl. 25, SYNTYPE, Adult (relaxed mount). Reg. no. 1855.12.19.74. Maldonado, Uruguay, June. Collected by Charles Darwin and purchased of the Zoological Society of London. There is another syntype in the collection. cuzcoensis—Cvanolyca viridicvanea cuzcoensis W. L. Sclater, 1917. 71 *Ibis*, (10) 5: 463, pl. 8. HOLOTYPE, Adult. Reg. no. 1886.9.15.399. Huasampilla, 10,000′, N.E. of Cuzco, Andes, July 1871. Collected by H. Whitely. Sclater Collection. cvanea—Muscitrea cvanea Hume, 1877 Stray Feathers, 5:101. SYNTYPE, Adult male. Reg. no. 1886.4.1. 4264. Meetan, Tenasserim, (25 Feb. 1877, collected by W. Davison, on label). Hume Collection. The second syntype is also in the collection. cyaneifrons—Erythrura cyaneifrons Layard, 1878 60 Ann. Mag. nat. Hist., (5) 1:375. Although Sharpe, Hist. Coll. Brit. Mus., 2 (1906) 474, claimed that the types were in the collection. the original description was based on an old collection and no specimen in the museum collection would have been available at that period. cyanescens—Calliste cyanescens P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1856 (publ. 1857): 260. SYNTYPE, Adult male. Reg. no. 1885.6.12.337. Venezuela. Sclater Collection; bought of Leadbeater. The specimen is marked as the type on the label, and listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 123. cvaneus—Turdus cvaneus Horsfield, 1821 Trans. Linn. Soc. Lond., 13:149. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4521. Java. Presented by the Honourable East India Company Museum. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883):12. There is a second syntype in the collection. cyanilia-Tanagra cyanilia Bonaparte, 1854 C.R. Acad. Sci. Paris, 38:383. HOLOTYPE, Adult. Reg. no. 1854.1.17.14. Venezuela. Collected for and purchased of Verreaux fréres. The specimen is marked "Type" on the original label and Bonaparte states that he saw the new species in Verreaux's collection.

Listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 159.

cyaniventer—Tesia cyaniventer Hodgson, 1837
cyanochlamys—Malurus cyanochlamys Sharpe, 1881 36 Proc. zool. Soc. Lond., 1881 : 788. Holotype, Adult male. Reg. no. 1881.5.1.712. Moreton Bay, Queensland, Australia. Collected by F. Strange. Gould Collection.
cyanogenys—Psilorhinus cyanogenys Sharpe, 1877 71 Cat. Bds. Brit. Mus., 3:140, pl. 9. Holotype, Adult. Reg. no. 1867.7.24.1. Pearl-bay Lagoon, Mosquito, Nicaragua. Collected and presented by H. Wickham. Sharpe, used a nomen nudum from G. R. Gray, Handlist of Birds, 2 (1870):6.
cyanolaima—Calliste cyanolaima Bonaparte, 1851
cyanolaimus—Nectarinia cyanolaimus Jardine and Fraser, 1851 . 46 Contributions to Ornithology: 154. ? SYNTYPE, Adult male (relaxed mount). Reg. no. 1851.11.27.1. Fernando Po, (West Africa, on label). Collected by L. Fraser. Purchased through E. Gerrard. Specimens from the Jardine Collection were received and registered but cannot be traced. From the date of receipt it seems certain that this is a syntype. A female syntype is also in the collection.
cyanonota—Pitta cyanonota G. R. Gray, 1860
cyanopterus—Chloropsis cyanopterus Hodgson, 1844
cyanopygia—Calliste cyanopygia P. L. Sclater, 1883 51 Proc. zool. Soc. Lond., 1883 (Dec.): 653. Berlepsch and Taczanowski, using Sclater's MS name, published a description of the species based on a specimen at Warsaw, which appeared in the same volume of the Proc. zool. Soc. Lond. (p. 545) in September. The Warsaw specimen is therefore the type and Sclater's specimen has no type status.
cyanotis—Calliste cyanotis P. L. Sclater, 1858 51 Proc. zool. Soc. Lond., 1858: 294. Holotype, Adult. Reg. no. 1885.6.12.377. Interior of Peru bordering Bolivia. Collected for Verreaux. Sclater Collection.

- cyanotis—Turdus cyanotis Jardine and Selby, 1828 31 Illustrations of Ornithology, 1: pl. 46. HOLOTYPE, Adult. Reg. no. 1886.6.24.258. India. Jardine Collection; purchased through E. Gerrard.
- cyanouroptera—Siva cyanouroptera Hodgson, 1838 32B India Review, 2:88. Syntype, Adult (relaxed mount). Reg. no. 1843.1.13.446. Nepal. Collected and presented by B. H. Hodgson. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883):640. There are several other syntypes in the collection.
- cyornithopsis—Callene cyornithopsis Sharpe, 1901 . . . 31

 Bull. Br. Orn. Cl., 12:4. HOLOTYPE, Adult male. Reg. no.
 1901.10.29.28. Efulen, Cameroon, 20 June 1901. Collected by and purchased of G. L. Bates.
- cypriotis—Fringilla coelebs cypriotis J. M. Harrison, 1945 . . . 58

 Bull. Brit. Orn. Cl., 66: 6. HOLOTYPE, Adult male. Reg. no.
 1951.13.2737. Platres, Cyprus, 2 April 1938. Collected and presented by W. A. Payn.
- czarnikowi—Phyllanthus czarnikowi Ogilvie-Grant, 1907 . . . 32E Bull. Brit. Orn. Cl., 19: 40. HOLOTYPE, Adult male. Reg. no. 1906.12.23.1284. Mawambi, 3,000′, east Congo Forest, 18 Oct. 1906. Collected by R. B. Woosnam and presented by the Subscribers to the Ruwenzori Expedition.
- daaroodensis—Calandrella cinerea daaroodensis White, 1960 . . . 17
 Bull. Br. Orn. Cl., 80: 24. HOLOTYPE, Adult male. Reg. no. 1923.8.7.2739. Sheikh, British Somaliland, 1 Nov. 1917. Collected by G. F. Archer and presented by Col. Stephenson Clarke.
- dacotiae—Pratincola dacotiae Meade-Waldo, 1889 31

 Ibis, (6) 1:504, pl. 15. Syntype, Adult male. Reg. no. 1905.

 12.22.7. Gran Tanajal, Fuertaventura, Canary Islands. Collected and presented by E. G. Meade-Waldo.

 There are three other syntypes in the collection, and one in the

There are three other syntypes in the collection, and one in the University Museum of Zoology, Cambridge.

- daflaensis—Actinura daflaensis Godwin-Austen, 1875 32B Ann. Mag. nat. Hist., (4) 16:340. Holotype, Adult male. Reg. no. 1895.7.14.1422. Shengorh Peak, 7,000', Dafla Hills, Assam, Feb. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.
- daflaensis—Suthora daflaensis Godwin-Austen, 1876 33
 Ann. Mag. nat. Hist., (4) .17 : 32. HOLOTYPE, Adult. Reg. no.

- 1895.7.14.1516. Under Torúpútú Peak, 5,000', Dafla Hills, Assam, Jan. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.

- danjoui—Rimator danjoui Robinson and Kloss, 1919 32B Ibis, (11) 1:578, pl. 14. Syntype, Adult male. Reg. no. 1919. 12.20.279. Langbian Peaks, 6,000', south Annam, 27 April 1918. Collected by C. B. Kloss and presented by H. C. Robinson and Kloss.

This is the authors' selected male. Their selected female syntype is also in the collection.

- dankali—Fringillaria striolata dankali Thesiger and Meynell, 1934. 49

 Bull. Brit. Orn. Cl., 55: 79. HOLOTYPE, Adult male. Reg. no.
 1934.8.9.1. Adua Danakil, east Abyssinia, 9 March 1934. Collected by W. Thesiger. Thesiger Collection.
- dankali—Gynmoris pyrgita dankali Thesiger and Meynell, 1935. 61

 Bull. Br. Orn. Cl., 55: 122. HOLOTYPE, Adult male. Reg. no.
 1934.8.9.3. Adau, Danakil, eastern Abyssinia, 6 Mar. 1934.
 Collected and presented by W. Thesiger.

dartmouthi—Nectarinia dartmouthi Ogilvie-Grant, 1906 46 Bull. Brit. Orn. Cl., 16:117. SYNTYPE, Adult male. Reg. no. 1906.12.23.549. Eastern Ruwenzori, 13,000', West Uganda, 19 Feb. 1906. Collected by R. E. Dent. Presented by the subscribers to the Ruwenzori Expedition. Labelled as the male type by the author. There are several other syntypes in the collection.
darwini—Upucerthia darwini Scott, 1900
davidiana—Suthora davidiana Slater, 1897
davisoni—Acanthopneuste davisoni Oates, 1889 35 The Fauna of British India, Birds, 1:420. SYNTYPE, Adult male. Reg. no. 1886.7.8.643. Mooleyit, 6,000', Tenasserim, 19 Feb. 1877. Collected by W. Davison. Hume Collection. There are two other syntypes in the collection.
davisoni—Alcippe phaeocephala davisoni Harington, 1915 32B J. Bombay nat. Hist. Soc., 23:447, 453. SYNTYPE, Adult male. Reg. no. 1886.10.1.6494. Tavoy, Tenasserim, 20 April 1878. Collected by W. Davison. Hume Collection.
davisoni—Brachyurus davisoni Hume, 1875
davisoni—Cryptolopha davisoni Sharpe, 1888
davisoni—Hemixus davisoni Hume, 1877
davisoni—Ixus davisoni Hume, 1875
davisoni—Leioptila davisoni Hume, 1877

1.763. Hills north of Mooleyit, 6,000', Tenasserim, 1 Feb. 1877. Collected by W. Davison. Hume Collection. There are four other syntypes in the collection.
davisoni—Stachyris davisoni Sharpe, 1892
davisoni—Turdinulus davisoni Ogilvie-Grant, 1910
davisoni—Turdulus davisoni Hume, 1877
decipiens—Mimeta decipiens P. L. Sclater, 1883 6. Proc. zool. Soc. Lond., 1883: 199. SYNTYPE, Adult (female on label). Reg. no. 1883.5.30.98. Larat, Timor Laut, Tenimber Island Group, 15 Sept. 1882. Collected by H. O. Forbes. Presented by the Timor Laut Committee of the British Association. The second syntype is in the collection.
decora—Vidua decora Hartlaub, 1862 6 Ibis, (1) 4:340. HOLOTYPE, Adult male. Reg. no. 1873.12.10.6. Angola. Collected by and purchased of J. J. Monteiro.
decorata—Niltava grandis decorata Robinson and Kloss, 1919 . 3 Ibis, (11) 1: 444. SYNTYPE, Adult female. Reg. no. 1919.12.20.17. Langbian Peaks, 6,500', South Annam, 23 April 1918. Collected and presented by C. B. Kloss. This is the described female. The described male syntype (which differs less distinctly from N. g. decipiens) is also in the collection.
decoratus—Lamprocolius decoratus Hartlaub, 1862 6 Ibis, (1) 4:148. HOLOTYPE, Adult (relaxed mount). Reg. no. 1874.5.1.72. Cape Colony, South Africa. Collected by E. L. Layard. Sharpe Collection.
degeni—Mirafra degeni Ogilvie-Grant, 1902
dehrae—Sturnopastor capensis dehrae E. C. S. Baker, 1925 6 Bull. Br. Orn. Cl., 45: 103. HOLOTYPE, Adult female. Reg. no. 1887.7.1.146. Dehra Dun, India, 20 Dec. 1870. Collected by G. King. Hume Collection.

deignani—Aegithina tiphia deignani Hall, 1957
delacouri—Neomixis viridis delacouri Salomonsen, 1934 32B Ann. Mag. nat. Hist., (10) 14:63. HOLOTYPE, Adult male. Reg. no. 1931.8.18.2937. Maraontsetra, north-east Madagascar, 23 June 1930. Collected by the Archbold-Vernay Expedition.
delacouri—Smithornis capensis delacouri Bannerman, 1923 1 Bull. Br. Orn. Cl., 43:160. HOLOTYPE, Adult. Reg. no. 1923. 11.12.400. Bandama, Ivory Coast, Upper Guinea, 29 Dec. 1922. Collected by W. P. Lowe and H. R. Hardy.
delamerei—Coliuspasser delamerei Shelley, 1903 61 Bull. Br. Orn. Cl., 13:73. HOLOTYPE, Adult male. Reg. no. 1901.2.22.37. Ngari Mossor, north Kenya, 17 Mar. 1900. Collected and presented by Lord Delamere.
delamerei—Estrilda delamerei Sharpe, 1900
delamerei—Pseudalaemon delamerei Sharpe, 1900 17 Bull. Br. Orn. Cl., 10:102. HOLOTYPE, Adult male. Reg. no. 1901.2.22.244. Athi River, British East Africa, 17 Nov. 1899. Collected and presented by Lord Delamere.
delicatula—Zosterops delicatula Sharpe, 1882
deliciosa—Pipra deliciosa P. L. Sclater, 1860
deltarhyncha—Guiraca caerulea deltarhyncha van Rossem, 1938 . 50 Bull. Brit. Orn. Cl., 58: 133. Holotype, Adult female. Reg. no. 1899.2.1.257. Tepic, Nayarit, Mexico, 8 June 1889. Collected by W. B. Richardson. Salvin-Godman Collection.
demissa—Synallaxis demissa Salvin and Godman, 1884 3 Ibis, (5) 2:449. SYNTYPE, Adult male. Reg. no. 1889.5.14.119.

- 5,000', Roraima, British Guiana, 16 Aug. 1883. Collected by H. Whitely. Salvin-Godman Collection.

 There are two female syntypes in the collection.
- dennisi—Petroica multicolour dennisi Cain and Galbraith, 1955. 37

 Bull. Brit. Orn. Cl., 75: 93. HOLOTYPE, Adult female. Reg. no. 1959.21.957. Turipava, 4,100', Guadalcanar, Solomon Islands, 19 Aug. 1953. Collected by a native hunter. Presented by the Oxford University Expedition to the Solomon Islands.
- dentata—Malacothraupis dentata P. L. Sclater and Salvin, 1876.

 Proc. zool. Soc. Lond., 1876: 353, pl. 31. HOLOTYPE, Adult.
 no. 1885.6.7.522. Tilotilo, Yungas Province, Bolivia, 1876. Collected by C. Buckley. Salvin-Godman Collection.

desertae—Prinia superciliosa desertae Macdonald, 1941 35 Bull. Brit. Orn. Cl., 62:27. HOLOTYPE, Adult male. Reg. no. 1922.12.8.1221. Kulme, (3,300', on label), Darfur, Western Sudan, 9 June 1921. Collected by Admiral H. Lynes. Presented by the Lynes-Lowe Expedition.
desolata—Tchitrea desolata Salomonsen, 1933
deva—Alauda deva Sykes, 1832
diabolicus—Bradyornis diabolicus Sharpe, 1877
diademata—Yuhina diademata J. Verreaux, 1869 32B Nouv. Arch. Mus. Paris Bull., 5:35. HOLOTYPE, Adult male. Reg. no. 1898.10.3.836. Moupin, west China, 4 March 1869. Collected by Père David. Seebohm bequest. Received by Swinhoe from Paris Museum as the type (vide specimen label).
dichroa—Monarcha dichroa G. R. Gray, 1859
dichrous—Cyphorhinus dichrous P. L. Sclater and Salvin, 1879. 28 Proc. zool. Soc. Lond., 1879: 492, pl. 41. HOLOTYPE, Adult. Reg. no. 1886.9.15.72. Remedios, 2,360', far north-east of Medellin, Antioquia. Collected by T. K. Salmon. Sclater Collection.
dichrous—Parus dichrous Blyth, 1845

There is another syntype in the Australian Museum, Sydney (vide Hindwood, Records of the Australian Museum, 21 (1946): 393).

110 dienegae
Ann. Mag. Nat. Hist., 15 (1845): 326, was not apparently published until May of that year.
dickeyae—Icterus graduacauda dickeyae van Rossem, 1938 57 Bull. Brit. Orn. Cl., 58:137. HOLOTYPE, Adult male. Reg. no. 1898.12.14.1180 (not 1177 as printed, op.cit). Xautipa, Guerrero, Mexico, July 1888. Collected by D. W. Smith. Salvin-Godman Collection.
dicruriformis—Dissemuroides dicruriformis Hume, 1873 64
Stray Feathers, 1:408. SYNTYPE, Adult male. Reg. no. 1886.3.1. 2155. Great Coco Island, Indian Ocean, (24 March 1873. Collected by H. Iwaries, on label). Hume Collection. There are two other syntypes in the collection.
dictator—Timelia pileata dictator Kinnear, 1930 32E
Bull. Br. Orn. Cl., 50: 55. HOLOTYPE, Adult male. Reg. no. 1919.12.20.282. Dran, south Annam, 11 May 1918. Collected by C. B. Kloss and presented by H. C. Robinson and Kloss.
dieffenbachii—Miro dieffenbachii J. E. Gray, 1843
dieffenbachii—Miro dieffenbachii J. E. Gray, 1843 37 in Dieffenbach, Travels in New Zealand, 2:191. Holotype, Adult male (relaxed mount). Reg. no. 1842.10.30.15. Chatham Island, (New Zealand, on label; South Island, New Zealand, vide Fleming, Trans. Roy. Soc. N.Z., 78 (1950):29, 36). Collected by Dr Sinclair and presented through Sir John Richardson.
diemenensis—Acanthiza diemenensis Gould, 1838
Synopsis of the birds of Australia, pt. 4: pl. 59. (Also in Proc. zool. Soc. Lond., 1837, (publ. Dec. 1838): 146). SYNTYPE, Adult (relaxed mount). Reg. no. 1838.1.15 specimen number omitted from label and not certainly known. Tasmania. Collected and presented by R. Gunn. Mathews, Austral Avian Record 1 (1912): 163, refers to a type in the Philadelphia Museum.
diemenensis—Rhipidura diemenensis Sharpe, 1879
Ibis, (4) 3:368. New name for R. saturata Sharpe, 1879 (q.v.), on the grounds of its preoccupation by R. saturata Salvadori, 1878.
difficilis—Botha difficilis Shelley, 1902
The Birds of Africa 3: 105, pl. 22. HOLOTYPE, Immature (female). Reg. no. 1889.9.13.164. Orange River Colony, 1875. Collected by T. E. Buckley. Shelley Collection.
difficilis—Geospiza difficilis Sharpe, 1888
Cat Rds Rrit Mus 12:12 Syxtype Adult male Dea no

Cat. Bds. Brit. Mus., 12:12. SYNTYPE, Adult male. Reg. no. 1885.12.14.321. Galapagos Archipelago (restricted to Abingdon Island, vide Rothschild and Hartert, Novit Zool., 6 (1899):163), Dec. 1868. Collected by A. Habel. Salvin-Godman Collection. Of the two male syntypes this is the described specimen. There are six other syntypes in the collection.

unut 11	_
difficilis—Zosterops difficilis Robinson and Kloss, 1918 47 J. Fed. Malay States Mus., 8, pt. 2:250. PARATYPE, Adult female. Reg. no. 1883.2.19.2. Mount Dempo, 8,000', Sumatra, 1 May 1881. Collected by H. O. Forbes. Presented by F. Nicholson. This paratype has been included because the original description was based on a figure of a specimen in the Liverpool Museum and a description by Sharpe, Cat. Bds. Brit. Mus., 9 (1884):191. The Liverpool specimen was destroyed (vide Gibson-Hill, Bull. Raffles Mus., 19 (1949):163, footnote) and this paratypical specimen on which Sharpe's description was based is therefore the only remaining type material.	7
diffusa—Pyrgita diffusa Smith, 1836 6 Report of the expedition for exploring Central Africa of 1834. : 50. SYNTYPE, Adult (relaxed mount). Reg. no. 1838.6.9.132 and Old Vol. Cat. 21, 275b. Between the Orange River and the Tropic. (Restricted to Kuruman, northern Cape Province by Macdonald and Hall, Ann. Transvaal Mus., 23 (1) (1955) : 35). Collected by A. Smith. Purchased from the South African Museum, London. There is a second syntype in the collection.	1
diffusus—Oriolus diffusus Sharpe, 1877	3
dignissima—Grallaria dignissima P. L. Sclater and Salvin, 1880. Proc. zool. Soc. Lond., 1880: 160, pl. 17. SYNTYPE, Adult. Reg. no. 1889.7.10.836. Sarayacu, Ecuador. Collected by C. Buckley. Salvin-Godman Collection. There is another syntype in the collection.	4
digressa—Ortygospiza atricollis digressa Clancey, 1958 6 Durban Mus. Novit., 5:141. HOLOTYPE, Adult male. Reg. no. 1933.7.14.477. Whitson, Umhlongo Nek, Richmond, Natal, 13 Sept. 1927. Collected by and purchased of J. Vincent.	
diluta—Cotile diluta Sharpe & Wyatt, 1893	8
diluta—Otocorys diluta Sharpe, 1890	7
diluta—Rhipidura diluta Wallace, 1864	

- Reg. no. 1873.5.12.466. Flores Island, 1862. Collected by Wallace's assistant Allen. Wallace Collection.

 There are several other syntypes in the collection.
- diluta—Saxicola diluta Blanford and Dresser 1874 31

 Proc. zool. Soc. Lond., 1874: 234, pl. 39. SYNTYPE, Adult male
 (in text; female on label). Reg. no. 1876.5.23.424. Hukomkap,
 Damaraland, 13 May 1866. Collected by C. J. Andersson.
 Sharpe Collection.

There are several other syntypes in the collection.

- diluta—Stachyris poliocephala diluta Robinson and Kloss, 1918. 32B Ibis, (10) 6:587. HOLOTYPE, Adult female. Reg. no. 1936.4. 12.1065. Taiping, Perak, Malaya, 4 Nov. 1914. Presented by H. C. Robinson and C. B. Kloss.
- diluta—Thringorhina guttata diluta Kinnear, 1924 32B Bull. Br. Orn. Cl., 45:11. HOLOTYPE, Adult male. Reg. no. 1924.12.21.139. Thai Nien, Tonkin, 23 Feb. 1924. Collected by H. Stevens and presented by the Percy Sladen Trust and Godman Exploration Fund.
- diluta—Volatinia jacarini diluta van Rossem, 1938 59
 Bull. Brit. Orn. Cl., 58:130. HOLOTYPE, Adult male. Reg. no.
 1899.2.1.908. San Blas, Tepic, (Nayarit), Mexico, 5 May 1889.
 Collected by W. B. Richardson. Salvin-Godman Collection.
- dilutior—Chlorocichla flaviventris dilutior White, 1946 21

 Ibis, 88:80. HOLOTYPE, Adult male. Reg. no. 1953.54.228.

 Balovale, Kaonde-Lunda Province, Northern Rhodesia, 12 June, 1943. Collected and presented by C. M. N. White.
- dilutior—Malacocincla fulvescens dilutior White, 1953 . . . 32B Bull. Br. Orn. Cl., 73:96. Holotype, Adult male. Reg. no. 1910.5.6.674. N'dala Tando (now Vila Salazar), north Angola, 29 Dec. 1908. Collected by Dr W. J. Ansorge. Ansorge Collection.
- dilutus—Stachyris nigriceps dilutus Robinson and Kloss, 1919 . 32B Ibis, (11) 1:584. SYNTYPE, Adult male. Reg. no. 1919.12.20. 302. Dram, 3,000′, south Annam, 1 April 1918. Collected by C. B. Kloss and presented by H. C. Robinson and Kloss.

This is the authors' selected male. Their selected female syntype is also in the collection. Renamed S. n. rileyi Chasen, 1936 on the grounds of its preoccupation by S. poliocephala diluta Robinson and Kloss, 1918.

dilutus—Turdinulus epilepidotus dilutus Robinson and Kloss, 1916. 32B J. Straits Br. roy. Asiat. Soc., 73:276. HOLOTYPE, Adult male. Reg. no. 1920.6.29.353. Sungei Kumbang, 4,700', Korinchi, west Sumatra, 13 April 1914. Collected and presented by H. C. Robinson and C. B. Kloss.

dimorpha—Pachyprora dimorpha Shelley, 1893
dinemelli—Textor dinemelli Rüppell, 1845
diops—Batis diops Jackson, 1905
disjuncta—Hirundo daurica disjuncta Bates, 1930
dispar—Turdus dispar Horsfield, 1821
disturbans—Dicrurus leucophaeus disturbans E. C. S. Baker, 1918 . 64 Novit. Zool., 25: 293. SYNTYPE, Adult male. Reg. no. 1883.3.1. 2044. Amherst, Tenasserim, Burma, 9 Dec. 1876. Collected by W. Davison. Hume Collection. The female syntype is also in the collection.
divaricatus—Lanius divaricatus Raffles, 1822
dives—Grallaria dives Salvin, 1865

dixoni—Geocichla dixoni Seebohm, 1881	31
djamdjamensis—Anthreptes collaris djamdjamensis Benson, 1942 . 4 Bull. Brit. Orn. Cl., 63:16. HOLOTYPE, Adult male. Reg. no. 1946.5.2642. Alghe, 6,000', Southern Abyssinia, 22 Sept. 1941. Collected and presented by C. W. Benson.	16
dodsoni—Cercomela familiaris dodsoni Macdonald, 1953 3 Ibis, 95: 72. HOLOTYPE, Adult male (printed in error as female). Reg. no. 1903.3.9.209. Deelfontein, east Cape Province, 5 May 1902. Collected by C. H. E. Seimund and C. H. B. Grant and presented by A. T. Sloggett.	31
dodsoni—Cisticola dodsoni Sharpe, 1895	35
dodsoni—Lanius algeriensis dodsoni Whitaker, 1898 2 Ibis, (7) 4:599. SYNTYPE, Adult. Reg. no. 1939.12.14.3. Rasil Ain, Haha District, south Morocco, 4 May 1897. Collected and presented by E. Dodson.	23
dodsoni—Pycnonotus dodsoni Sharpe, 1895	_
doggetti—Cinnyris doggetti Sharpe, 1902	16
dominicana—Certhiola dominicana E. C. Taylor, 1864 5 Ibis, (1) 6:167. HOLOTYPE, adult. Reg. no. 1884.7.31.181. Dominica, West Indies. Collected by E. C. Taylor. Sclater Collection.	54
donaldsoni—Cossypha donaldsoni Sharpe, 1895	
donaldsoni—Ploceipasser donaldsoni Sharpe, 1895 6 Bull. Br. Orn. Cl., 5:14. HOLOTYPE, Adult male. Reg. no.	

- 1895.7.7.23. ("near Lasamis, Lake Rudolf", on label), northern frontier of Kenya and Somaliland, ("14 Sept. 1895" on label). Collected and presented by A. Donaldson Smith.
- dorjei—Cinclus pallasi dorjei Ludlow and Kinnear, 1937 . . . 27 Ibis, (14) 1:263. HOLOTYPE, Adult male. Reg. no. 1935.4.5.12 (not 1936.4.5.12 as printed). Sakden, 9,000', east Bhutan, 18 Oct. 1934. Collected by F. Ludlow and G. Sherriff. Ludlow Collection.
- dorsalis—Automolus dorsalis P. L. Sclater and Salvin, 1880 3

 Proc. zool. Soc. Lond., 1880 : 158. HOLOTYPE, Adult. Reg. no. 1889.5.14.360. Sarayacu, Ecuador, 1880. Collected by C. Buckley. Salvin-Godman Collection.

dorsalis—Zosterops dorsalis Vigors and Horsfield, 1827 47 Trans. Linn. Soc. Lond., 15:235. Holotype, Adult. Reg. no. 1863.7.7.38. Australia. Collected by Caley. Presented by the Linnean Society as the type.
dorsatus—Parus dorsatus Rüppell, 1845
douglasi—Pitta douglasi Ogilvie-Grant, 1910
dowii—Calliste dowii Salvin, 1863
dryas—Malacocichla dryas Gould, 1854
dryas—Rhipidura dryas Gould, 1842
dubareasis—Lanius excubitor dubarensis C. H. B. Grant and Mackworth-Praed, 1951

dubia—Myrmecocichla dubia Blundell and Lovat, 1899 31 Bull. Br. Orn. Cl., 10: 22. HOLOTYPE, Adult. Reg. no. 1900.1.3. 112. Fontaly, south Abyssinia, 17 Jan. 1899. Collected and presented by H. Weld Blundell and Lord Lovat.
dubius—Budytes dubius Hodgson, 1844
dubius—Proparus dubius Hume, 1874
dubius—Pyrocephalus dubius Gould, 1839
dubius—Turdus dubius Latham, 1801
dukhunensis—Alauda dukhumensis Sykes, 1832
dukhunensis—Motacilla dukhunensis Sykes, 1832
dulitensis—Buchanga leucophaea dulitensis Harrisson and Hartley, 1934

1933.11.2.4. Mount Dulit, 4,500', Sarawak, Borneo, 22 Aug. 1932. Collected by T. H. Harrisson and C. H. Hartley. Presented by the Oxford University Borneo Expedition. of 1932.

dumasi—Poecilodryas brachyura dumasi Ogilvie-Grant, 1915 . . . 37

Ibis, (10) 2, Supplement: 163. HOLOTYPE, Adult male. Reg. no. 1900.4.10.21. North New Guinea. Collected by J. M. Dumas.

Purchased through Miss Sharpe.

Mayr, List of New Guinea Birds (1941): 142, stated that the locality was Humboldt Bay, but appeared to have overlooked Hartert's statement, Novit. Zool., 36 (1930): 118, that the Dumas collection was made on the eastern slopes of the Arfak Mountains and that he and Lord Rothschild had erroneously believed it to be Humboldt Bay.

syntypes in the collection.

- duncani—Myiodynastes solitarius duncani Chubb, 1919 . . . 9
 Bull. Brit. Orn. Cl., 40:62. SYNTYPE, Adult male. Reg. no.
 1922.3.5.2808. Supenaam River, British Guiana, 1913. Collected for F. V. McConnell. McConnell Collection.
 The female syntype is also in the collection.

Jardine used a Thompson MS. name. Subsequently, in the same volume, op. cit. 257 and pl. 30, he regarded this species as synony-

mous with N. adelberti Gervais, 1834.

eburneus—Phyllastrephus eburneus Bannerman, 1923 . . . 21

Bull. Br. Orn. Cl., 43:162. HOLOTYPE, Adult female. Reg. no.
1923.11.12.207. Béoumi, Ivory Coast, West Africa, 11 Dec.
1922. Collected and presented by W. P. Lowe and H. R. Hardy.

eclipes—Chatorhea eclipes Hume, 1877
edgari—Oriolus maculatus edgari Chasen, 1939
edithae—Corvus edithae Phillips, 1895
editus—Anthus richardi editus Vincent, 1951
cfulenensis—Eurillas efulenensis Sharpe, 1904
egertoni—Actinodura egertoni Gould, 1836
egregia—Dacnis egregia P. L. Sclater, 1855
elaeoprorus—Buarremon elaeoprorus P. L. Sclater and Salvin, 1879. 49 Proc. zool. Soc. Lond., 1879: 504. Syntype, Adult male. Reg. no. 1885.6.8.151. Santa Elena, Antioquia, Colombia. Collected by T. K. Salmon. Salvin-Godman Collection. There are two other syntypes in the collection.
elegans—Cistothorus elegans P. L. Sclater and Salvin, 1859 28 Ibis, (1) 1:8. HOLOTYPE, Adult female. Reg. no. 1885,3.6.467. Lake of Dueñas, Guatemala, 1 Feb. 1858. Collected by O. Salvin. Salvin-Godman Collection.
elegans—Dendrornis elegans Pelzeln, 1868

central Brazil, 8 Aug. 1826. Collected by J. Natterer. P. L. Sclater Collection. Listed as "authentic specimen" by Sclater, Cat. Bds. Brit. Mus.,
15 (1890) : 137.
elegans—Hypocnemis elegans P. L. Sclater, 1857
in Swainson and Richardson, Fauna Boreali-Americana, 2 (Birds), 1831 (publ. 1832): 122. HOLOTYPE, Adult (relaxed mount). Reg. no. Old Vellum Cat. 10.4. "The fur countries" (error, southern Algeria or Tunisia, vide Ogilvie-Grant, Novit. Zool., 9 (1902): 457, and Hartert Vog. Palaarkt. Fauna, 1 (1907): 427). Presented by the Hudson Bay Company.
elegans—Minus elegans Sharpe, 1881
elegans—Phoenicornis elegans McClelland, 1840
elegans—Synallaxis elegans P. L. Sclater, 1856
elegantior—Synallaxis elegantior P. L. Sclater, 1862
elegantula—Emberiza elegantula Swinhoe, 1870
elenae—Poecile atricapillus elenae P. R. Lowe, 1921

- year). Reg. no. 1921.8.2.7. (Giomeiu, on label) Breuil, 7,000', Valtournanche, Northern Italy, (27 July 1921, on label). Collected and presented by P. R. Lowe.
- elfriedae—Mirafra sabota elfriedae Hoesch and Niethammer, 1940 17 J. Orn., 88 (Sonderheft): 217. SYNTYPE, Adult male. Reg. no. 1937.2.17.56. Onguma, Etoscha Salt Pan, Damaraland, 10 Oct. 1936. Collected by W. Hoesch. Hoesch Collection. The female syntype is also in the collection.

Specimen Reg. no. 1893.12.1.27, from Kikuyu, mentioned by Sharpe, *Ibis*, (6), 4 (1892): 160, was incorrectly listed as type in Seebohm, *Monograph of the Turdidae*, 1 (1902): pl. 72 and Jackson, *Birds of Kenya and Uganda*, 2 (1938): 951. See the correspondence in the Brit. Mus. (Nat. Hist.) with Dr. H. Schouteden, 1962, 1964.

- elgonensis—Poliospiza elgonensis Ogilvie-Grant, 1912 59
 Bull. Brit. Orn. Cl., 31:17. HOLOTYPE, Adult male. Reg. no.
 1912.7.22.1. Mangiki, 6,000', Mount Elgon, Kenya, 14 June 1900.
 Collected and presented by F. J. Jackson.
- elizae—Cryptospiza elizae Boyd Alexander, 1903 60
 Bull. Brit. Orn. Cl., 13:38. Holotype, Adult male. Reg. no.
 1911.12.23.3293. Bakaki, 4,000', Fernando Po, 20 Nov. 1902.
 Collected by Boyd Alexander. Boyd Alexander Bequest.

made. In *Ibis*, (8) 3 (1903): 354 the description is repeated with a slight difference in the tarsal measurement, and it is stated that the type is the male, the date given being 14. Dec., although the date of collection of the male specimen is given as 12 Dec. on the label, the female having been collected on the 14 Dec.

eliae—Irena ellae Steere, 1890
ellisiana—Pyriglena ellisiana P. L. Sclater, 1855
elopurensis—Siphia elopurensis Sharpe, 1890
elwesi—Otocoris elwesi Blanford, 1872
emendata—Cisticola chiniana emendata Vincent, 1944 35 Bull, Brit. Orn. Cl., 64:63. Holotype, Adult male. Reg. no. 1933.3.1.1230. Mirrote, 900′, Mozambique Province, Portuguese East Africa, 13°50′ S, 39°35′E., 17 June 1932. Collected and presented by J. Vincent.
emigrans—Xiphocolaptes emigrans P. L. Sclater and Salvin, 1859 Ibis, (1) 1: 118. Syntype, Adult female. Reg. no. 1889.5.14.685. San Geronimo, Vera Paz, Guatemala, 23 Dec. 1859. Collected by O. Salvin. Salvin-Godman Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 145. There is another syntype in the collection.
emiliae—Nigrita emiliae Sharpe, 1869
empusa—Lagenoplastes empusa Blyth, 1866
enca—Fregilus enca Horsfield, 1821

Bull. Brit. Orn. Cl., 51:29. HOLOTYPE, Adult male. Reg. no. 1901.10.21.31. Enderby Island, near Auckland Island, New Zealand, 3 Jan. 1901. Collected by R. Henry (vide Ogilvie-Grant, Ibis, (8) 5 (1905): 586). Presented by the Earl of Ranfurly. enganensis—Siphia enganensis Ogilvie-Grant, 1895 Bull. Brit. Orn. Cl., 5: 2. SYNTYPE, Adult male. Reg. no. 1897. 5.13.316. Cape Engano, Luzon, Philippines, (29 April 1895, on label). Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. Listed as the type by Hachisuka, Birds of the Philippines, 2 (1935): 295, but other specimens are referred to in the original description. entebbe—Cisticola avresii entebbe Lynes, 1930 Ibis, (12) 6, Supplement: 154. HOLOTYPE, Adult male. Reg. no. 1928.10.25.9. Entebbe, Uganda, 6 June 1927. Collected by Admiral H. Lynes, and B. B. Osmaston. Presented by Admiral H. Lynes. eoa—Sylvicola eoa Gosse, 1847. Birds of Jamaica: 158. SYNTYPE, Adult male (relaxed mount). Reg. no. 1847.6.8.15. Crab Pond, Jamaica, 21 or 24 Jan. Collected by P. H. Gosse and purchased as the type. See also Peters, Proc. Biol. Soc. Wash., 40 (1927): 32-33. The specimen is faded, and the equally faded female syntype is also in the collection. epauletta—Cinnyris epauletta Hodgson, 1837 India Review, 1837: 272. SYNTYPE, Adult male. Reg. no. 1845. 1.9.523. Nepal. Collected and presented by B. H. Hodgson. There are other syntypes in the collection. epauletta—Pyrrhula? epauletta Hodgson, 1836 Asiat. Res., 19:156. SYNTYPE, Adult male (relaxed mount). Reg. no. 1843.1.13.838. Nepal. Collected and presented by B. H. Hodgson. There are other syntypes in the collection. ephippialis—Ramphocelus ephippialis P. L. Sclater, 1861. 51 Proc. zool. Soc. Lond., 1861:130. SYNTYPE, Adult male. Reg. no. 1885.6.12.553. Upper Amazon (error,=South Brazil, vide Sclater, Cat. Bds. Brit. Mus., 11 (1886): 171). Collected for Verreaux. Sclater Collection. The female syntype is also in the collection. ephippialis—Turdus ephippialis P. L. Sclater, 1862 31 Proc. zool. Soc. Lond., 1862: 109. HOLOTYPE, Adult. Reg. no. 1886.8.2.86. Bogota, Colombia. Collected by Trübner. Sclater Collection; obtained by the latter from the Bremen Museum.

enderbyi—Myiomoira macrocephala enderbyi Mathews, 1930.

epimacurus—Cinnyris epimacurus Hodgson, 1844 in J. E. Gray, Zoological Miscellanv: 82. This is a nomen nudum and the specimen referred to as the type by Gadow, Cat. Bds. Brit. Mus., 9 (1884): 25, is not valid. equatorialis—Rhopoterpe torquata equatorialis Chubb, 1918. Bull. Br. Orn. Cl., 38:85. SYNTYPE, Adult male. Reg. no. 1889. 9.20.567. Sarayacu, east Ecuador, 1880. Collected by C. Buckley. Sclater Collection. There are three other syntypes in the collection. equicaudata—Cisticola exilis equicaudata E. C. S. Baker, 1924 Bull. Brit. Orn. Cl., 44:39. HOLOTYPE, Adult male. Reg. no. 1916.12.27.954. Sambok, Siam, 29 August 1915. Collected and presented by E. G. Herbert. eremita—Nesocichla eremita Gould, 1855 31 Proc. zool. Soc. Lond., 1855: 164. HOLOTYPE, Adult. Reg. no. 1856.10.14.9. Tristan d'Acunha. Collected by J. Macgillivray on the voyage of H.M.S. Herald. eremodites—Pyrrhulauda eremodites R. Meinertzhagen, 1923. Bull. Br. Orn. Cl., 43:156. HOLOTYPE, Adult female. Reg. no. 1923.6.18.1. Sheikh Othman, Aden Protectorate, 15 Jan. 1922. Presented by Col. R. Meinertzhagen. erithacus—Liosceles erithacus P. L. Sclater, 1890 . Cat. Bds. Brit. Mus., 15:345. SYNTYPE, Adult male. Reg. no. 1889,9.20.707. Sarayacu, Ecuador, 1880. Collected by C. Buckley. Sclater Collection. There are three other syntypes in the collection; the fifth is now at the American Museum of Natural History, New York. ernesti-Oenanthe leucopyga ernesti R. Meinertzhagen, 1930 in Nicoll's Birds of Egypt, 1:280. HOLOTYPE, Adult male. Reg. no. 1965.M.18. Wadi Feiran, Sinai, 1 March 1928. Collected by R. Meinertzhagen. Meinertzhagen Collection. 1900.1.19.29. Camp 22, 13,000', Teliki Valley, Mount Kenya, 2 Sept. 1899. Collected by and purchased of H. J. Mackinder. erochroa—Abrornis erochroa J. E. Gray, 1846 Catalogue of mammalia and birds of Nepal and Thibet presented by B. H. Hodgson to the British Museum: 66, 152. SYNTYPE, Adult. Reg. no. 1843.1.13.282. Nepal. Collected and presented by B. H. Hodgson.

Listed with another specimen as the types by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 74. There are other syntypes in the collection. Gray used a Hodgson MS. name.

erromangae—Rhipidura erromangae Sharpe, 1900 Ibis. (7) 6:340. SYNTYPE. Adult male. Reg. no. 1900.1.10.19. Polenia Bay, Erromanga, New Hebrides, 25 May 1899. Collected and presented by A. M. Farguhar, from the voyage of H.M.S. Walleroo.

The second syntype is also in the collection.

- erwini—Zosterops palpebrosa erwini Chasen, 1934. Bull. Raffles Mus., 9:96. HOLOTYPE, Adult female (male, vide original label, and Gibson-Hill, Bull. Raffles Mus., 19 (1949): 150). Reg. no. 1947.60.60. Pulau Panjang, South Natuna Island, South China Seas, 9 August 1931. Collected by P. M. de Fontaine. Presented by the Raffles Museum, Singapore, as the type.
- erythacus—Siphia erythacus Jerdon and Blyth, 1861 Proc. zool. Soc. Lond., 1861: 210. (Not Siphia ervthaca of Blyth. J. Asiat. Soc. Bengal, 16 (1847): 126, from Malaya). SYNTYPE. Adult male. Reg. no. 1886.4.1.1913. Darjeeling, Sikkim, India. 1860. Collected by T. C. Jerdon. Hume Collection.

Jerdon, Birds of India, 1 (1862): 480, stated that he collected a specimen at Darjeeling. This specimen was labelled and registered as the type. Blyth appeared to have overlooked his earlier use of

this name for a different species (op. cit. 1847).

- erythrocephalum—Cinclosoma erythrocephalum Vigors, 1831 Proc. zool. Soc. Lond., 1831: 171. Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 361, lists a Hodgson specimen as the type; but Vigors stated that Gould presented the type to the Zoological Society Museum, and it was not received from the society when the museum was dispersed.
- erythrocephalus—Ploceus erythrocephalus Rüppell, 1845 Systematische Uebersicht der Vögel nord-ost Afrika's: 71. Holo-TYPE, Adult (relaxed mount). Reg. no. 1861.5.8.44. Shoa, Abyssinia. Collected by W. C. Harris. Presented by the Honourable East India Company Museum.
- erythrocerca—Formicivora erythrocerca P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858: 240, pl. 142. HOLOTYPE, Adult male. Reg. no. 1881.2.18.118. Brazil? Eyton Collection.
- erythrocercus—Myiarchus erythrocercus P. L. Sclater and Salvin, 1869 9 Proc. zool. Soc. Lond., 1868 (publ. 1869): 631. SYNTYPE, Adult. Reg. no. 1888.1.13.1082. Caraccas, Venezuela. Collected by A. Goering. Sclater Collection.

There are several other syntypes in the collection.

erythrocnemis—Pomatorhinus erythrocnemis Gould, 1863 Proc. zool. Soc. Lond., 1862 (publ. 1863): 281. SYNTYPE, Adult. Reg. no. 1898.10.3.614. North Formosa, April 1862 (on label). Collected by R. Swinhoe. Seebohm Bequest. There are several other syntypes in the collection.

- erythrolaema—Phoenicothraupis erythrolaema P. L. Sclater, 1862 51 Catalogue of a collection of American birds: 83. HOLOTYPE, Adult male. Reg. no. 1885.6.12.681. Sta. Marta, New Granada (error, see Todd and Carriker, Ann. Carnegie Mus., 14 (1922): 483. Corrected to "vicinity of the mouth of the Rio Sinú, Colombia", by Parkes (letter, MS. in prep.)). Collected for Verreaux. Sclater Collection.

Sclater used a Bonaparte MS. name.

- erythrolaema—Trochalopterum erythrolaema Hume, 1881 . . . 32B Stray Feathers, 10:153. HOLOTYPE, Adult male. Reg. no. 1886. 10.1.2785. Matchi, Manipur, India, 8 May 1881. Collected by A. O. Hume. Hume Collection.
- erythroleuca—Grallaria erythroleuca P. L. Sclater, 1873 4

 Proc. zool. Soc. Lond., 1873 : 783. HOLOTYPE, Adult. Reg. no. 1889.9.20.623. Huasampilla, Peru, Mar. 1872. Collected by H. Whitely. Sclater Collection.
- erythronota—Geocichla erythronota P. L. Sclater, 1859 31

 Ibis, (1) 1:113. SYNTYPE, Adult male. Reg. no. 1873.5.12.75.

 Macassar, Celebes, 1857. Collected by A. R. Wallace. Wallace Collection.

The second syntype is also in the collection.

- erythronota—Rhipidura erythronota Sharpe, 1879 37A Cat. Bds. Brit. Mus., 4:337. pl. 10. Holotype, Adult female. Reg. no. 1877.11.17.27. Vanua Levu, Fiji, 23 Oct. 1875. Collected by Kleinschmidt. Purchased of the Godeffroy Museum.
- erythronotus—Philydor erythronotus P. L. Sclater and Salvin, 1873 3 Nomencl. Av. Neotrop.: 160. HOLOTYPE, Adult. Reg. no. 1889. 5.20.362. Bogota, Colombia, 1870. Sclater Collection.
- erythrophrys—Pithys erythrophrys P. L. Sclater, 1855 4
 Proc. zool. Soc. Lond., 1854 (publ. 1855): 255, pl. 72. HOLOTYPE,
 Adult (female). Reg. no. 1889.9.20.519. Bogota, Colombia,
 1853. Collected for Argent. Sclater Collection.

Labelled as the type by the author, but three other syntypes were available and are in the collection.

- erythropleurus—Turdus erythropleurus Sharpe, 1887 31 Proc. zool. Soc. Lond., 1887 : 515. HOLOTYPE, Adult female. Reg. no. 1887.5.1.21. Christmas Island. Collected by Capt. Maclear of H.M.S. Flying Fish and presented by the Lords of the Admiralty.
- erythroprocta—Ruticilla erythroprocta Gould, 1855 3

 Proc. zool. Soc. Lond., 1855: 78. HOLOTYPE, Adult male. Reg. no. 1860.1.16.33. Erzeroum, Turkey. Collected by Dickson and Ross (vide Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 342) and purchased of J. Gould.

Labelled as "one of the type specimens". There are other syntypes in the collection, from the Eyton Collection; and two more in Philadelphia (vide de Schauensee, Proc. Acad. Sci. Nat. Philadelphia, 109 (1957): 197).

erythrops—Leptopogon erythrops P. L. Sclater, 1862 9

Proc. zool. Soc. Lond., 1862:111. SYNTYPE, Adult. Reg. no. 1888.1.13.417. Bogota, Colombia. Collected for Trübner. Sclater Collection.

The second syntype is in the collection.

- erythroptera—Drymoica erythroptera Jardine, 1849 35

 Contributions to Ornithology: 15, pl. 2. HOLOTYPE, Adult. Reg.
 no. 1886.6.24.210. Gold Coast. Collected by C. A. Gordon.
 Jardine Collection.

erythroptera—Mirafra erythroptera Blyth, 1845
erythroptera—Terpsiphone erythroptera Sharpe, 1879 38 Cat. Bds. Brit. Mus., 4:357. SYNTYPE, Adult male. Reg. no. 1875.4.9.45. Gambia River, Senegambia. Collected by Whitely. Purchased of R. B. Sharpe. According to Sclater, Systema Avium Aethiopicarum (1930):432, the specimens are examples of the Indian species, T. paradisi which have been wrongly labelled. A female syntype is in the collection.
erythropterus—Anabates erythropterus P. L. Sclater, 1856
erythropterus—Andropadus erythropterus Hartlaub, 1858 2 Proc. zool. Soc. Lond., 1858 : 292. HOLOTYPE, Immature. Reg. no. 1858.1.4.97. Ashanti, Gold Coast. Collected by S. T. D. Aubinn, a native collector and purchased of J. Gould. Hartlaub used G. R. Gray's MS. name.
erythropterus—Formicarius erythropterus Gould, 1855
erythropterus—Lanius erythropterus Vigors, 1831
erythropterus—Lichenops erythropterus Gould, 1839
erythropterus—Turdus (Monticola) erythropterus G. R. Gray, 1860 3. Proc. zool. Soc. Lond., 1860: 350. HOLOTYPE, Adult male. Reg. no. 1873.5.12.573. Gilolo, 1858. Collected by A. R. Wallace. Wallace Collection.

9.1.2373. Pakchan, Tenasserim, 26 Jan. 1875. Collected by W. Davison. Hume Collection.

There is a second male syntype in the collection.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 164. According to Kinnear (MS.) there were two males and two females in the Sykes Collection; only the above specimen was listed by Sharpe (loc. cit.).

- erythropygia—Saxicola erythropygia Cavendish Taylor, 1867. . . . 31

 Ibis, (2) 3:61. HOLOTYPE, Immature. Reg. no. 1905.6.28.501.

 River Nile, Egypt, March 1864. Collected and bequeathed by E. Cavendish Taylor.

The author was apparently unaware that the same name had been used by Sykes, *Proc. zool. Soc. Lond.*, 1832:92 (q.v.), for a different species.

- erythropygium—Edolisoma erythropygium Sharpe, 1888... 20
 Proc. zool. Soc. Lond., 1888: 184. SYNTYPE, Adult female. Reg.
 no. 1888.2.7.33. Guadalcanar, Solomon Archipelago, 15 April
 1887. Collected by and purchased of C. M. Woodford.
 The male syntype is also in the collection.
- erythrorhynchus—Arremon erythrorhynchus P. L. Sclater, 1855 . 49

 Proc. zool. Soc. Lond., 1855: 83, pl. 89. HOLOTYPE, Adult male.

 Reg. no. 1855.11.5.26. Bogota, New Colombia. Collected for and purchased of J. Gould.

erythrothorax—Leistes erythrothorax Pelzeln, 1870 Ornithologie Brasiliens: 197, 326. SYNTYPE, Adult male. Reg. no. 1885.7.12.165. Rio Madeira, Brazil. 16 Nov. 1829. Collected by J. Natterer. Sclater Collection, received by him in exchange from the Vienna Museum. Pelzeln used a Natterer MS. name. A female syntype is also in the collection. Listed as "typical specimens", by P. L. Sclater, Cat. Bds. Brit. Mus., 11 (1886): 349.	-
erythrotis—Grallaria erythrotis P. L. Sclater and Salvin, 1876. Proc. zool. Soc. Lond., 1876: 357. SYNTYPE, Adult. Reg. no. 1889.7.10.876. Tilotilo, Yungas Province, Bolivia, 1876. Collected by C. Buckley. Salvin-Godman Collection. There are two other syntypes in the collection.	
erythrotis—Merula erythrotis Davison, 1886	•
erythrura—Myrmotherula erythrura P. L. Sclater, 1890	4
euchrysea—Hirumdo euchrysea Gosse, 1847	
eugenei—Myiophoneus eugenei Hume, 1873	•
euophrys—Basileuterus euophrys P. L. Sclater and Salvin, 1876 Proc. zool. Soc. Lond., 1876: 352. SYNTYPE, Adult male. Reg. no. 1885.3.8.708. Tilotilo, Yungas of La Paz, Bolivia, 1876. Collected by C. Buckley. Salvin-Godman Collection. The female syntype is also in the collection.	
euophrys—Thryothorus euophrys P. L. Sclater, 1860	
everetti—Arachnoraphis everetti Sharpe, 1893	46

Mount Kinabalu, North Borneo, Nov. 1892. Collected by A. H.
Everett. Purchased of E. Gerrard. This specimen, the only one purchased, was registered as the type but the original description was based on eight specimens.
everetti—Criniger everetti Tweeddale, 1877
everetti—Dicaeum everetti Tweeddale, 1877
everetti—Edolisoma everetti Sharpe, 1893
everetti—Geocichla everetti Sharpe, 1892
everetti—Oxycerca everetti Tweeddale, 1877 60 Ann. Mag. nat. Hist., (4) 20:96. SYNTYPE, Adult. Reg. no. 1888.9.20.9. Monte Alban, Luzon, Philippines, Feb. 1877. Collected by A. H. Everett. Tweeddale Collection. The other three syntypes are also in the collection, all four being labelled as types by the author.
everetti—Prionochilus everetti Sharpe, 1877
everetti—Siphia everetti Sharpe, 1890
everetti—Zosterops everetti Tweeddale, 1877

examinatus—Rhynchocyclus sulphurascens examinatus Chubb, 1920 9 Bull. Brit. Orn. Cl., 40: 108. Holotype, Adult. Reg. no. 1922. 3.5.2655. Bartica Grove, British Guiana, 1912. Collected by F. V. McConnell. McConnell Collection.
excellens—Cinnyris excellens Ogilvie-Grant, 1895
excelsior—Cinclodes excelsior P. L. Sclater, 1860 3 Proc. zool. Soc. Lond., 1860: 77. SYNTYPE, Adult female. Reg. no. 1889.5.20.48. 14,000′, Pichincha, Ecuador, Jan. 1859. Collected by L. Fraser. Sclater Collection. This specimen, although labelled female, was listed as male by Sclater, Catalogue of American Birds, 1862: 148. There is another syntype in the collection.
exigua—Geothlypis exigua Ridgway, 1902
exile—Todirostrum exile P. L. Sclater, 1857
exilis—Malurus exilis Vigors and Horsfield, 1827 35 Trans. Linn. Soc. Lond., 15: 223. HOLOTYPE, Reg. no. 1863.7.7.24. Australia. Presented by the Linnean Society of London as the type. The author used a Latham MS. name.
eximia—Nectarinia eximia Horsfield, 1821
eximium—Dicaeum eximium P. L. Sclater, 1877
explorator—Zosterops explorator Layard, 1875

vide Finsch, Proc. zool. Soc. Lond., 1877: 734). From the voyage of H.M.S. Challenger. A fuller description was given in Finsch (op. cit.). The second female syntype is also in the collection.
exsul—Spindalis exsul Salvin, 1885
extensicauda—Drymoica extensicauda Swinhoe, 1860 35 Ibis, (1) 2:50. HOLOTYPE, Adult male. Reg. no. 1881.5.1.373. Amoy, China, 9 April 1859. Collected by R. Swinhoe. Gould Collection. Both Gould and later Sharpe (Cat. Bds. Brit. Mus., 7 (1881)) appear to have overlooked the fact that this was the type.
extrema—Malacocincla rufipennis extrema Bates, 1930 32B Bull. Br. Orn. Cl., 51:49. HOLOTYPE, Adult male. Reg. no. 1930.12.3.8. Near Nzerekoré, 1,900', French Guinea, 15 May 1930. Collected by and purchased of G. L. Bates.
exul—Pericrocotus exul Wallace, 1863
exultans—Aethopyga ignicauda exultans E. C. S. Baker, 1925 . 46 Bull. Brit. Orn. Cl., 46:13. HOLOTYPE, Adult male. Reg. no. 1921.7.15.626. Shueli-Salwin divide, west-central Yunnan, (8,000', on label), July 1919. Collected by G. Forrest. Presented by Col. Stephenson Clarke.
eytoni—Dendrocolaptes eytoni P. L. Sclater, 1853
eytoni—Juida eytoni Fraser, 1856
ezrai—Aethopyga ezrai Delacour, 1926

China.

faenorum—Ammomanes deserti faenorum R. Meinertzhagen, 1951. 17 Proc. zool. Soc. Lond., 1951: 131. HOLOTYPE, Adult female. Reg. no. 1965.M.17. Bahrain Island, Persian Gulf, 19 Jan. 1951. Collected by R. Meinertzhagen. Meinertzhagen Collection.
fagani—Accentor fagani Ogilvie-Grant, 1913
fagani—Malimbus fagani Ogilvie-Grant, 1907
fagani—Serinus fagani Sharpe, 1896
fagani—Tchitrea fagani Bannerman, 1921
fairbanki—Trochalopteron fairbanki Blanford, 1869 32B J. Asiat. Soc. Bengal, 38 pt. 2:175, pl. 17. HOLOTYPE, Adult. Reg. no. 1869.9.15.1. Pulney Hills, south India. Collected by the Rev. S. Fairbank and presented by him through W. T. Blanford. See Blanford, J. Asiat. Soc. Bengal, 49, pt. 2 (1880):142–3 for a note on this specimen.
falcularius—Dendrocopus falcularius Vieillot, 1822
fallax—Elainea fallax P. L. Sclater, 1861
fallax—Lanius fallax Finsch, 1872

- Reg. no. 1888.5.30.361. Rairo, north-east Abyssinia, 13 Aug. 1868. Collected by W. Jesse. Tweeddale Collection.

 Two female syntypes are also in the collection.
- - There are two other syntypes, in bad condition, in the collection.

- fantiensis—Eremomela badiceps fantiensis Macdonald, 1940 . . . 35

 Ibis, (14) 4:341. HOLOTYPE, Adult male. Reg. no. 1911.12.23.
 2200. Prahau, Gold Coast, 1 July 1900. Collected by Boyd Alexander. Boyd Alexander Bequest.
- fantisiensis—Alseonax fantisiensis Sharpe, 1879 3'
 Cat. Bds. Brit. Mus., 4:131. Syntype, Adult (relaxed mount).
 Reg. no. 1875.4.9.475. Fantee, West Africa. Collected by A. Swanzy. Sharpe Collection.
 There are three other syntypes in the collection.
- fasciatus—Salpinctes fasciatus Salvin and Godman, 1891 . . . 28

 Ibis, (6) 3:610. SYNTYPE, Adult. Reg. no. 1897.10.1.233.

 Volcan el Viejo, 6,500', near Chinandega, Nicaragua, 28 April 1891. Collected by W. B. Richardson. Salvin-Godman Collection.

The second, immature, syntype is also in the collection.

A number of specimens were received by Gould (vide, Handbook to the birds of Australia, 1 (1865): 507). Another syntype, which was listed as the type by Gadow, Cat. Bds. Brit. Mus., 9 (1884): 240, is also in the collection, but the measurements do not agree with the description.

 north-east of Latakoo. (South Africa, on label). Collected by

and purchased of A. Smith.

Registered as the type and listed as such by Sharpe, *Cat. Bds. Brit. Mus.*, 7 (1883): 134, but two other syntypes are referred to in the description.

- fasciolatus—Acrocephalus fasciolatus G. R. Gray, 1860 . . . 35

 Proc. zool. Soc. Lond., 1860: 349. HOLOTYPE, (Immature male, vide Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 110). Batchian, Molucca Islands, 1858. Collected by A. R. Wallace. Wallace Collection.
- feadensis—Calornis (Aplonis) feadensis Ramsay, 1881 62 J. Linn. Soc. Lond. Zool., 16:129. Holotype, Adult male. Reg. no. 1895.12.24.5. Fead Island, Solomons. Collected by the Rev. G. Brown. Received in exchange from E. P. Ramsay, Sydney.

- feminina—Hyphantornis feminina Ogilvie-Grant, 1907 61

 Bull. Br. Orn. Cl., 21:15. Syntype, Adult female. Reg. no. 1906.12.23.148. Mokia, 3,400′, south-east Ruwenzori, 2 May 1906. Collected by R. E. Dent and G. Legge. Presented by the subscribers to the Ruwenzori Expedition.

 There are other syntypes in the collection.
- feminina—Lophorhina superba feminina Ogilvie-Grant, 1915 . . . 70

 Ibis, (10) 2, Jubilee Supplement: 27. Syntype, Adult male. Reg. no. 1916.5.30.1043. Camp 6B, Utakwa River, 4,200', Dutch New Guinea, 2 Feb. 1913. Collected by C. B. Kloss. Presented by British Ornithologists' Union New Guinea Expedition.

This is the author's selected male. His selected female and several other syntypes are also in the collection.

femininus—Carpodacus femininus Rippon, 1906 59

Bull. Brit. Orn. Cl., 19:31. Syntype, Adult female. Reg. no.
1906.12.17.565. Yangtze River, 10,000', Big Bend, West Yunnan,
23 March 1906. Collected by G. Rippon and presented as the type.

There are three other syntypes in the collection.

- fernandonis—Orthotomus sutorius fernandonis Whistler, 1939 . . . 35

 Bull. Brit. Orn. Cl., 60:15. HOLOTYPE, Adult male. Reg. no. 1940.12.3.364. Ohiyo, 5,820', Central Hill Zone, Ceylon, 27 Nov. 1936. Collected by E. C. Fernando on the British Museum and Colombo Museum Avifaunal Survey.

There are several other syntypes in the collection.

Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 31., states, apparently in error, that the specimen came from the voyage of H.M.S. Sulphur.

Goulds, Birds of New Guinea 2 (1884): pl. 36, implies that the type was at Sydney, but no holotype was designated in the original description. The reference in Nature 20 (1879): 125 is a nomen

nudum.

- ferrugiceps—[Lanius] Collurio ferrugiceps Hodgson, 1837 . . . 23

 The Indian Review, 1:446. SYNTYPE, Adult. Reg. no. 1843.1.13.

 768. Almost confined to the lower region of Nepal. Presented by B. H. Hodgson.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 650.

ferruginea—Hemichelidon ferruginea Hodgson, 1845 Proc. zool. Soc. Lond., 1845: 32. SYNTYPE, Adult. Reg. 1848.6.4.120. Nepal. Presented by B. H. Hodgson. There are a number of syntypes in the collection.	. 37 no.
ferrugineifrons—Hylophilus ferrugineifrons P. L. Sclater, 1862. Proc. zool. Soc. Lond., 1862:110. HOLOTYPE, Adult. Reg. 1886.9.15.317. Interior of Colombia. (Collected by Trübner label.) Sclater Collection.	no.
fervidus—Accentor fervidus Sharpe, 1883	. 30 no. y H.
figlina—Pyranga figlina Salvin and Godman, 1883 Biol. CentrAmer., Aves 1:293. Syntype, Adult female. no. 1885.6.7.352. Poctum, Guatemala, March 1862. Colle by O. Salvin. Salvin-Godman Collection. The male syntype is in the United States National Muse vide Deignan, Type specimens of birds in the U.S. National Muse (1961):581.	ected eum,
filigera—Ptilotis filigera Gould, 1851	snea
finschi—Cassinia finschi Sharpe, 1870	. 31 .493. arpe
finschi—Emphonia finschi P. L. Sclater and Salvin, 1877 Proc. zool. Soc. Lond., 1877: 19. HOLOTYPE, Adult male. no. 1885.6.12.105. Demerara, British Guiana. Collected Wilkens. Sclater Collection, received by him in exchange from Bremen Museum.	Reg.
finschi—Trichostoma finschi Walden, 1876 Ibis, (3) 6:378. pl. 11. SYNTYPE, Adult male. Reg. 1873.5.12.48. Macassar, Celebes. Collected by A. R. Walwallace Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883):54	llace.
fitzgeraldi—Upucerthia fitzgeraldi Scott, 1900	Col-

flammea—Lexops flammea Wilson, 1890
Proc. zool. Soc. Lond., 1889 (publ. 1890): 445. SYNTYPE, (male, on label). Reg. no. 1939.12.9.307. Kalae, Molokai, Sandwich Islands, June 1888. Collected by S. B. Wilson. Rothschild
Bequest.
Label is printed "E Mus. Acad. Cantabrigiae". The plumage is less vivid than that described but the specimen was collected at the
same time as a pair of syntypes in the University Museum of Zoology, Cambridge.
flammeus—Macrosphenus kempi flammeus Marchant, 1950 35
Bull. Brit. Orn. Cl., 70: 26. HOLOTYPE, Adult male. Reg. no. 1950.9.1. Umuagwu, Owerri Division, Southern Nigeria, 21 Aug. 1949. Collected and presented by S. Marchant.
flammifer—Pericrocotus flammifer Hume, 1875
Stray Feathers, 3:321. SYNTYPE, Adult male. Reg. no. 1886. 4.1.393. Malewoon, Bankasoon, south Tenasserim, 25 March
1875. Collected by W. Davison. Hume Collection.
There is also a female syntype in the collection.
flammifera—Arachnothera flammifera Tweeddale, 1878 46
Proc. zool. Soc. Lond., 1878: 343. HOLOTYPE, Adult male. Reg.
no. 1888.7.1.492. South Leyte, Philippines, Sept. 1877. Collected
by A. H. Everett. Tweeddale Collection. The type is illustrated in Shelley, Monograph of the Nectariniidae,
(1879): pl. 115.
flammulata—Megabyas flammulata J. and E. Verreaux, 1855 38
Rev. Zool., 7:348. HOLOTYPE, Adult female. Reg. no. 1854. 5.23.3. Mouda, Gaboon, West Africa. Purchased of Verreaux.
flammulatus—Anabates flammulatus Eyton, 1849
in Jardine, Contributions to Ornithology, 1849: 131. Adult. Reg. no. 1881.2.18.204. Bogota, Colombia. Eyton Collection. There is another syntype in the collection.
flammulatus—Synalaxis flammulatus Jardine, 1850
Reg. no. 1886.6.24.452. 14,000', Andes of Quito, Ecuador. Col-
lected by W. Jardine. Jardine Collection.
flava—Nectarinia flava Bloxham, 1826
in Byron, Voyage of H.M.S. Blonde, Appendix: 249. SYNTYPE,
Reg. no. Old Vellum Cat. 26, no. 56b. Sandwich Islands. Col-
lected by A. Bloxham. Presented by Lord Byron, R.N. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 10.
Another syntype, a relaxed mount with broken bill, is also in the
Collection.

178 flavala
flavala—Hemixos flavala Hodgson, 1845
flaveolus—Regulus flaveolus G. R. Gray, 1848
flavescens—Aethopyga ignicauda flavescens E. C. S. Baker, 1921 . 46 Bull. Brit. Orn. Cl., 41:71. Holotype, Adult male. Reg. no. 1905.9.10.959. Mount Victoria, South Chin Hills, Burma, 19 March, 1904. Collected and presented by Col. G. Rippon.
flavescens—Hypocnemis flavescens P. L. Sclater, 1865 4 Proc. zool. Soc. Lond., 1864 (publ. 1865): 609. Holotype, Adult (female, vide Sclater, Cat. Bds. Brit. Mus., 15 (1880): 286). Reg. no. 1889.9.20.505. Marabitanas, Rio Negro, Brazil, May 1831. Collected by J. Natterer. Sclater Collection; obtained by Sclater in exchange with the Vienna Museum. Sclater used Natterer's MS name.
flavescens—Sylvia flavescens G. R. Gray, 1860 35 Proc. zool. Soc. Lond., 1860: 349. SYNTYPE, Adult (female, on label). Reg. no. 1873.5.12.221. Batchian, South-west Pacific, 1858. Collected by A. R. Wallace. Wallace Collection. This specimen is labelled as the type, but there is another syntype in the collection.
flavicans—Myiobius flavicans P. L. Sclater, 1861
flavicapillum—Heteropelma flavicapillum P. L. Sclater, 1860 8 Proc. zool. Soc. Lond., 1860: 466. SYNTYPE, Adult. Reg. no. 1888.1.13.1378. South-east Brazil, 1853. Sclater Collection. Sclater obtained it through Argent. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 322. There is another syntype in the collection.
flaviceps—Rhodocanthis flaviceps Rothschild, 1892 55 Ann. Mag. nat. Hist. (6) 10:111. SYNTYPE, adult (male, on label). Reg. no. 1939.12.9.133. (collector's no. 1443). Hawaii Island, Sandwich Group, (19 Oct. 1891, on label). Collected by H. C. Palmer. Rothschild Bequest. No holotype was designated. The specimen listed as the type by

- Hartert, Novit. Zool., 26 (1919): 169, was one of a series of syntypes which also included the above specimen and a female which is also in the collection
- flavicineta—Poecilodryas flavicineta Sharpe, 1879. Ann. Mag. nat. Hist., (5) 3:313. HOLOTYPE, Adult. Reg. no. 1879.3.6.61. (Port Moresby, on label), South-east New Guinea. Collected by K. Broadbent. Purchased of Gerrard.
- flavieollis—Mirafra flavicollis McClelland, 1840 in Horsfield, Proc. zool. Soc. Lond., 1839, (publ. 1840): 163. Syntype, Male (immature, vide Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 512). Reg. no. 1880.1.1.4639. Assam. Collected by J. McClelland. Presented by the Honourable East India Company Museum.

There is a second syntype in the collection.

- flavieollis—Pipra flavicollis P. L. Sclater, 1851 Contributions to Ornithology: 143. SYNTYPE, Adult male. Reg. no. 1888.1.13.1247. Upper Amazon, Barra do Rio Negro, Brazil, 1851. Collected by A. R. Wallace. Sclater Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 294.
- flavicollis—Yuhina? flavicollis Hodgson, 1836 Asiat. Res., 19:166. SYNTYPE, Adult (relaxed mount). Reg. no. 1843.1.13.369. Nepal. Collected and presented by B. H. Hodgson.

There is a second syntype in the collection.

- flavierissalis—Eremomela flavicrissalis Sharpe, 1895 Proc. zool. Soc. Lond., 1895: 481 (not 48, as quoted by many authors). Holotype, Adult. Reg. no. 1895.7.7.9. Shebeli, Somaliland, 6 Sept. 1894. Collected and presented by A. Donaldson Smith.
- flavierissus—Cassiculus flavierissus P. L. Sclater, 1860 . Proc. zool. Soc. Lond., 1860: 276. SYNTYPE, Adult male. Reg. no. 1885.7.12.42. Babahoyo, Ecuador, 1859. Collected by L. Fraser. Sclater Collection.

A female syntype is in the collection.

- flavida—Geothlypis flavida Ridgway, 1902 Birds of North and Middle America. pt. 2:678. HOLOTYPE, Adult male. Reg. no. 1906.12.7.3406. Nassau, New Providence Island, Bahamas. 30 May 1889. Collected by C. V. Maynard. G. S. Miller Jr. Collection.
- flavidieollis—Saltator flavidicollis P. L. Sclater, 1860 Proc. zool. Soc. Lond., 1860: 274. SYNTYPE, Adult. Reg. no. 1885.6.12.1058. Babahoyo, Ecuador, July-Sept. 1859. Collected by L. Fraser. Sclater Collection.

The two other syntypes are in the collection.

flavidifrons—Tyrannulus flavidifrons P. L. Sclater, 1860	9
flavidiventris—Estrelda flavidiventris Wallace, 1864 60 Proc. zool. Soc. Lond., 1863 (publ. 1864): 495. SYNTYPE, Adult male. Reg. no. 1873.5.12.1340. East Timor, 1861. Collected by A. R. Wallace. Wallace Collection. There are several other syntypes in the collection.	0
flavifrons—Muscicapa flavifrons Gmelin, 1789 4 Systema Naturae, 1, pt. 2:944. Holotype, Unsigned finished painting. Tanna, New Hebrides, 7 Aug. 1774. No. 158 in Forster, Unpublished water-colour drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 308.	7
flavigaster—Anthreptes flavigaster Eyton, 1839	6
flavigularis—Abrornis flavigularis Godwin-Austen, 1878 39 J. Asiat. Soc. Bengal., 47; pt. 2:19. HOLOTYPE, Adult. Reg. no. 1895.7.14.915. Sadiya, North-east India, Jan. 1877. Collected by M. T. Ogle. Godwin-Austen Collection. See also Ripley, Synopsis of the Birds of India and Pakistan (1961): 486.	5
flavigularis—Apalis flavigularis Shelley, 1893 35 Ibis, (6) 5:16, Syntype, Adult. Reg. no. 1892.9.10.89. Milanji Plateau, South Nyasaland, 26 Oct. 1891. Collected by A. Whyte. Presented by H. H. Johnstone. This specimen was presented as the type but there are two other syntypes in the collection.	5
flavigularis—Platyrhynchus flavigularis P. L. Sclater, 1862	9
flavimentalis—Abrornis schisticeps flavimentalis E. C. S. Baker, 1924 35 Bull. Brit. Orn. Cl., 44:63. HOLOTYPE, Adult. Reg. no. 1905. 9.10.816. Mount Victoria, South Chin Hills, Burma, 18 April 1904. Collected and presented by G. Rippon.	5
flavipectus—Aethopyga flavipectus Ogilvie-Grant, 1894	6

- 1897.5.13.111. La Trinadad, Benguet District, North-west Luzon, Philippines, 25 March 1894. Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition.

Listed as the type by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 320.

- flavirostris—Anaeretes flavirostris P. L. Sclater and Salvin, 1876 . 9

 Proc. zool. Soc. Lond., 1876: 355. SYNTYPE, Adult. Reg. no. 1888.1.1.493. Tilotilo, Yungas Province, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.

 There are two other syntypes in the collection.
- flavirostris—Chlorophonia flavirostris P. L. Sclater, 1861 . . . 51

 Proc. zool. Soc. Lond., 1861: 129. HOLOTYPE, Adult, ?female.

 Reg. no. 1885.6.12.23. Ecuador. Collected for J. Gould. Sclater

 Collection.

- flavirostris—Myzantha flavirostris Vigors and Horsfield, 1827. 48

 Trans. Linn. Soc. Lond., 15:319. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.68. Syndey, Australia. Collected by G. Caley. Presented by the Linnean Society of London as the type.

No holotype was designated. The specimen listed as the type by Hartert, *Novit. Zool.*, 26 (1919): 169, was one of a series of syntypes which also included the above specimen.

flaviventer—Machaerorhynchus flaviventer Gould, 1851. . . . 38

Birds of Australia, Supplement (March 1851): pl. 11. HOLOTYPE,

Adult (relaxed mount). Reg. no. 1857.10.16.6. Cape York,

Queensland. Collected by Captain O. Stanley R.N. Purchased

of J. Gould.

Gould stated that the single type was in the Museum of the Zoological Society, but appears not to have returned it after preparing the illustration, and to have sold it to the Museum. The label is inscribed "Type".

- flaviventre—Camptostoma flaviventre P. L. Sclater and Salvin, 1865 9

 Proc. zool. Soc. Lond., 1864 (Publ. 1865): 358. SYNTYPE, Adult.

 Reg. no. 1888.1.1.586. Panama. Collected by J. McLeannan.

 Salvin-Godman Collection.

The second syntype is also in the collection.

Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 157. The other syntype is also in the collection.

There is at least one other syntype in the collection.

Cat. Bds. Brit. Mus., 5 (1881): 132.

flaviventris—Muscipeta flaviventris J. and E. Verreaux, 1855. . . 38 J. Orn., 1855: 103. ?Syntype, Adult. Reg. no. 1854.5.23.26.

- Gaboon, West Africa. Collected by and purchased of J. and E. Verreaux. No single type was defined and it is not known if other specimens were used flaviventris—Specotheres flaviventris Gould, 1849. Proc. zool. Soc. Lond., 1849:1849:111. SYNTYPE, Adult male. Reg. no. 1850.7.20.149. Cape York, North East Australia, 7 Oct. 1848. Collected by J. Macgillivray. Presented by Captain O. Stanley from the voyage of H.M.S. Rattlesnake. There are two other syntypes in the collection. flavivertex—Elainea flavivertex P. L. Sclater, 1887 Proc. zool. Soc. Lond., 1887: 49. HOLOTYPE, Adult. Reg. no. 1888.1.13.628. Upper Ucayali River, East Peru. Collected by E. Bartlett. Sclater Collection. flavivertex-Heterocercus flavivertex Pelzeln, 1869 Ornithologie Brasiliens: 125, 186. SYNTYPE, Adult male. Reg. no. 1888.1.13.1393. Maribatanas, Rio Negro, Brazil, 25 March 1831. Collected by J. Natterer. Sclater Collection, received by him in exchange from Vienna. A female syntype is also in the collection. flavivertex—Setophaga flavivertex Salvin, 1887 884. Sierra Nevada of Santa Marta, Colombia, Nov. 1886. Collected for H. Whitely. Salvin-Godman Collection. The second syntype is in the collection. (21 April 1868, on label). Collected by W. T. Blanford. Presented by the Governor and Council of Bombay, through the author. A female syntype from the same source is also in the collection. label). Seebohm Bequest. Collected by Charles Darwin and purchased of the Zoological Society of London.
- flavolateralis—Acanthiza flavolateralis G. R. Gray, 1859 . . . 36

 Proc. zool. Soc. Lond., 1858: 161. HOLOTYPE, Adult. Reg. no. 1859.4.19.8. Island of Nu, Port de France, New Caledonia (20 April 1858, on label). Collected by J. Macgillivray. Purchased of Cuming.

flavolivacea—Neornis flavolivacea Hodgson, 1845
flavostriata—Nectarinia flavostriata Wallace, 1865 4 Proc. zool. Soc. Lond., 1865: 478. pl. 29. SYNTYPE, Adult male. Reg. no. 1873.5.12.859. Menado, North Celebes, 1859. Collected by A. R. Wallace. Wallace Collection. This specimen was listed as the type by Gadow, Cat. Bds. Brit. Mus., 9 (1884): 24. There are several other syntypes in the collection.
flavostriata—Ptilotis flavostriata Gould, 1875
flavostriatus—Andropadus flavostriatus Sharpe, 1876 2. Ibis, (3) 6:53. SYNTYPE, Adult female. Reg. no. 1875.10.7.51. Macamac, Lydenburg, Transvaal, 1 July 1874. Collected and presented by F. A. Barratt. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881):100.
flavotincta—Glyciphila flavotincta G. R. Gray, 1870
flavotincta—Grallaria flavotincta P. L. Sclater, 1877
flavotincta—Lalage flavotincta Sharpe, 1899
Bull. Brit. Orn. Cl., 10:28. SYNTYPE, Adult female. Reg. no. 1900.1.10.9. Mallicolo Island, New Hebrides, 1899. Collected and presented by A. M. Farquhar from the voyage of H.M.S. Wallaroo.
The author subsequently listed this specimen first (<i>Ibis</i> , (7) 6 (1900): 342), and also labelled it as the type. The other three syntypes are also in the collection and are from Espiritu Santo Island.
flavotincta—Pipra flavotincta P. L. Sclater, 1853 8 Proc. zool. Soc. Lond., 1852 (publ. 1853): 34, pl. 48. SYNTYPE, Adult (male; on label). Reg. no. 1888.1.13.1349. Santa Fé de Bogata, Colombia. (Collected by Parzudaki, on label.) Sclater Collection.
Listed as the type by Schater, Cat. Bds. Brit. Mus., 14 (1888): 315.

floris	185
flavovirescens—Microeca flavovirescens G. R. Gray, 1858 Proc. zool. Soc. Lond., 1858: 178. SYNTYPE, Adult male. F. no. 1858.3.10.64. Aru Islands. Collected by A. R. Walla Wallace Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 4 (1879): The female syntype is also in the collection.	ace.
flavoviridis—Orthotomus flavoviridis Moore, 1854 in Horsfield and Moore, Catalogue of birds in the Museum of Honourable East India Company, 1:314. SYNTYPE, Adult m Reg. no. 1880.1.1.2001. Malacca. Presented by the Honoura East India Company Museum as the type. The second syntype is also in the collection.	ale.
flavum—Dicaeum flavum Horsfield, 1821	. 47

- flavum—Malacopteron magnirostris flavum Chasen and Kloss, 1928 32B J. Malay roy. Asiat. Soc., 6:58. HOLOTYPE, Adult male. Reg. no. 1947.60.52. Pulau Siantan, Anamba Group, South China Sea, 21 Sept. 1925. Collected by F. N. Chasen and presented by the Raffles Museum, Singapore.

This specimen is clearly the male syntype but was erroneously labelled as a female. The two other syntypes are also in the collection.

- florida—Calliste florida P. L. Sclater and Salvin, 1869 51

 Proc. zool. Soc. Lond., 1869 : 416, pl. 28. HOLOTYPE, Adult female.

 Reg. no. 1885.6.4.374. Costa Rica, 1869. Collected by J. Carmiol. Salvin-Godman Collection.

no. 1873.5.12.1806. Flores, 1862. Collected by A. R. Wallace. Wallace Collection.

There is another syntype in the collection. In Peters, *Checklist* of Birds of the World, 9 (1960): 171, the reference is given as Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 9, 14.

- fluviatilis—Muscisaxicola fluviatilis P. L. Sclater and Salvin, 1866 9

 Proc. zool. Soc. Lond., 1866: 87. SYNTYPE, Adult. Reg. no. 1888.1.13.206. Lower Ucayali River, East Peru. Collected by E. Bartlett. Sclater Collection.

There are several other syntypes in the collection.

fohkiensis—Pericrocotus speciosus fohkiensis E. C. S. Baker, 1920 20 Bull. Br. Orn. Cl., 40:116. Syntype, Adult female. Reg. no. 1905.12.24.211. Yamahan, Fohkien, China. Collected and presented by C. B. Rickett.

There is another syntype in the collection.

- forbesi—Atlapetes rufigenis forbesi Morrison, 1947 49
 Bull. Brit. Orn. Cl., 67:88. HOLOTYPE, Adult male. Reg. no.
 1946.49.2. Pomayaco, 9,100', Apurimac Dept., Peru, 1 Oct.
 1939. Collected by and purchased of A. Morrison.
- forbesi—Lagonosticta rara forbesi Neumann, 1908 . . . 60

 Bull. Brit. Orn. Cl., 21:58. HOLOTYPE, Adult male. Reg. no.
 1884.1.10.254. Lokoja, Niger River, Northern Nigeria, 9 Sept.
 1882. Collected by W. A. Forbes. Forbes Collection.

- formosa—Fringilla formosa Gould, 1839 Zoology of the voyage of H.M.S. Beagle 3, Birds: 93. SYNTYPE, Adult male (relaxed mount). Reg. no. 1855.12.19.24. (Good Success Bay, on label), Tierra del Fuego. (The date, 4 Jan. 1837, on the label, probably refers to the date on which Gould received it). Collected by C. Darwin. Purchased of the Zoological Society of London. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 782. There are several other syntypes in the collection. formosa—Prinia inornata formosa Harington, 1913 35 1898.10.3.234. Laulong, Formosa, 6 Dec. 1893. Collected by P. A. Holst. Seebohm Bequest. Camphor Hills, Formosa, March 1862. Collected by R. Swinhoe. Seebohm Bequest. There are several other syntypes in the collection. formosae—Dendrocitta sinensis var. formosae Swinhoe, 1863. Ibis, (1) 5:387. SYNTYPE, Adult. Reg. no. 1897.6.1.229. Inner ranges of the Formosan Mountains, April 1862. Collected by R. Swinhoe. Seebohm Bequest. There is another syntype in the collection. Seebohm Collection. There are several other syntypes in the collection. 6.4.4. Mount Arizan, central Formosa, 25 Feb. 1909. Collected for and presented by Collingwood Ingram.
- formosanus—Carpodacus formosanus Ogilvie-Grant, 1911 . . . Bull. Brit. Orn. Cl., 27:51. New name for Carpodacus incertus Ogilvie-Grant, 1906 (q.v.), on the grounds of its preoccupation by Fringilla incerta Risso, 1826.
- formosanus—Proparus formosanus Ogilvie-Grant, 1906. . . . 32B Bull. Br. Orn. Cl., 16:120. HOLOTYPE, Adult male. Reg. no. 1907.12.12.62. Mount Morrison, 9,000', central Formosa, Jan. 1906. Collected by and purchased of W. Goodfellow.

formosum—Dicaeum formosum Ogilvie-Grant, 1912 4
Bull. Brit. Orn. Cl., 29:109. HOLOTYPE, Adult male. Reg. no.
1913.1.29.55. Mt. Arizan, 7,000', Formosa, March 1912. Col-
lected by W. Goodfellow. Purchased of E. J. Johnston.
The bill is damaged.
forreri—Chrysomitris forreri Salvin and Godman, 1886 5
Biol. CentrAmer., Aves, 1:429. Syntype, Adult (female, vide
Van Rossem, Bull. Brit. Orn. Cl., 58 (1938): 136). Reg. no. 1885.
12.14.1185. Durango, South-west Mexico, 19 June 1882. Col-
lected by A. Forrer. Salvin-Godman Collection.
The second syntype is also in the collection.
forreri—Vireo forreri Madarász, 1885
Termeszet Füzet, 1885: 9, 85, pl. 6. NEOTYPE? Adult. Reg. no.
1885.3.10.29. Tres Marias Islands, 4 May 1881. Collected by
Forrer. Salvin-Godman Collection.
The holotype appears to have been destroyed and P. R. Grant,
Acta. zool. Budapest 12 (1966): 79, proposes what is, in effect, a
neotype and appears to fulfil the requirements for this as laid down
in the International Code of Zoological Nomenclature, except that
he does not use the actual word "neotype" in the designation
and does not fully establish the loss of the original holotype.
forresti—Ianthocincla forresti Rothschild, 1921
Novit. Zool., 28: 35. SYNTYPE, Adult male. Reg. no. 1921.7.15.
237. Shweli-Salwin Divide, 7,000', May 1919. Collected by G.
Forrest and presented by Stephenson Clarke.
forresti—Phylloscopus proregulus forresti Rothschild, 1921 3
Novit. Zool., 28:45. HOLOTYPE, Adult female. Reg. no. 1921.
7.15.410. Lichiang Range, 9.000', North West Yunnan, May 1918.
Lat. 7°10′ N. Collected by G. Forrest. Presented by Col. R.
Stephenson Clarke.
·
forresti—Ptilotis forresti Ingram, 1906
Bull. Brit. Orn. Cl., 16:116. SYNTYPE, Adult. Reg. no. 1881.
5.1.5575. Australia. Collected for J. Gould. Gould Collection.
This is the Gould syntype referred to by the author. Hartert,
Novit. Zool. 26 (1919): 177, refers to a specimen in the Tring
Museum, now in the American Museum of Natural History, as the
"type".
forsythi—Pyrhocorax forsythi Stoliczka, 1874
Stray Feathers 2: 462. HOLOTYPE, ?Juvenile. Reg. no. 1891.7.22.
24. "On road from Lamaguru to the Indus" (Leh, Yarkand,
Kashmir, 4 Sept. 1873, on label). Collected by F. Stoliczka.
Received in exchange with the Indian Museum.
There had been some confusion in the registration numbers and
labels, but this specimen agrees with the original description.
fortipes—Horornis fortipes Hodgson, 1845
Proc. zool. Soc. Lond., 1845: 31. There is no evidence that
Hodgson's specimens in the collection can be considered as types or
o in the second of the second

- syntypes, although listed as such by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 137, since they were collected after the date of the description.
- fortirostris—Drymoica fortirostris Jardine and Fraser, 1852 . . . 35 in Jardine, Contributions to Ornithology: 60. HOLOTYPE, Adult male. Reg. no. 1886.6.24.204. Abomey, Dahomey, 1852. Collected by L. Fraser. Jardine Collection.

- fortis—Percnostola fortis P. L. Sclater and Salvin, 1867 . . . 4
 Proc. zool. Soc. Lond., 1867: 980, pl. 45. SYNTYPE, Adult male.
 Reg. no. 1889.9.20.461. Pebas, east Peru, 13 July 1866. Collected by J. Hauxwell. Sclater Collection.
 The two other syntypes are in the collection.

- forwoodi—Motacilla forwoodi Ogilvie-Grant and Forbes, 1899 . 19
 Bull. Liverpool Mus., 2:3. Holotype, Adult female. Reg. no. 1899.8.11.127. Abd-el-Kuri Island, Socotra, 23 Feb. 1899. Collected by W. R. Ogilvie-Grant and H. O. Forbes and presented by the Royal Society.

foxi—Calamornis foxi W. L. Sclater, 1927	
fraithii—Himatione fraithii Rothschild, 1892	
francisi—Motacilla francisi Swinhoe, 1870	19
fraseri—Anthreptes fraseri Jardine and Selby, 1843	46
fraseri—Basileuterus fraseri P. L. Sclater, 1884	54
fraseri—Conirostrum fraseri P. L. Sclater, 1858	54
fraseri—Muscicapa fraseri Strickland, 1844	31
fraseri—Oreomanes fraseri P. L. Sclater, 1860	51
frater—Amydrus frater P. L. Sclater and Hartlaub, 1881 6 Proc. zool. Soc. Lond., 1881: 171. HOLOTYPE, Adult male. Reg.	5 2

no. 1881.3.21.41. Socotra, 28 Feb. 1880. Collected by Prof. I. Bayley Balfour. Presented by the Socotra Committee of the British Association. frater—Herpsilochmus frater P. L. Sclater & Salvin, 1880 . Proc. zool. Soc. Lond., 1880: 159. SYNTYPE, Adult male. Reg. no. 1889.9.20.336. Sarayacu, Ecuador, Collected by C. Buckley. Sclater Collection. The female syntype is in the collection. frater—Oxyrhamphus frater P. L. Sclater and Salvin, 1868. Proc. zool. Soc. Lond., 1868: 326. SYNTYPE, Adult female. Reg. no. 1888.1.1.4. Calovevora, Veragua, 1867. Collected by E. Arcé. Salvin-Godman Collection. The other syntype is also in the collection. frater—Troglodytes frater Sharpe, 1881 Cat. Bds. Brit. Mus., 6:261. HOLOTYPE, Adult (relaxed mount). Reg. no. 1846,9.9.55. Bolivia. Collected by and purchased of T. Bridges. fratercula—Alcippe fratercula Rippon, 1900. Bull. Brit. Orn. Cl., 11:11. HOLOTYPE, Adult. Reg. no. 1900. 12.20.289. South Shan States, over 4,000', April 1899. Collected and presented by G. Rippon. fraterculus—Ammomanes fraterculus Tristram, 1864 Proc. zool. Soc. Lond., 1864: 434. SYNTYPE, Adult male. Reg. no. 1864.11.22.174. Engedi, Palestine. Collected by H. B. Tristram, Tristram Collection. fraterculus—Pericrocotus fraterculus Swinhoe, 1870 20 Ibis, (2) 6:244. SYNTYPE, Adult male. Reg. no. 1897.7.2.69. Hainan, China, Feb. 1868. Collected by R. Swinhoe. Seebohm Bequest. There is also a female syntype in the collection. fratrum—Pachyprora fratrum Shelley, 1900. 38 Ibis, (7) 6:522. SYNTYPE, Adult female. Reg. no. 1900.10.4.2. St. Lucia Lake, Zululand. Collected by R. B. and J. D. S. Woodward, Presented by P. L. Sclater. The second syntype is also in the collection. fredericki—Planesticus fredericki Chubb. 1918 Bull. Br. Orn. Cl., 38:53. HOLOTYPE, Adult. Reg. no. 1922. 3.5.3576. Bartica Grove, British Guiana, 1911. Collected by F. V. McConnell Collection. fremantlii—Calendula fremantlii Lort Phillips, 1897 Bull. Br. Orn. Cl., 6:46. SYNTYPE, Adult male. Reg. no. 1898. 7.27.37. Gedais, Goolis Mts., Somaliland, 18 Feb. 1897. Col-

lected by and purchased of E. Lort Phillips.

There are three other syntypes in the collection.

- fretensis—Munia punctulata fretensis Kloss, 1931 60
 Treubia, 13:363. HOLOTYPE, Adult male. Reg. no. 1936.4.12.
 2719. Kuala Lumpur, Selangor, Malaya (5 Nov. 1908, on label).
 Collected by C. B. Kloss. Presented by the Raffles Museum,
 Singapore.
- fretensis—Tephrodornis pelvica fretensis Robinson & Kloss, 1920 20 J. Straits Branch roy. Asiat. Soc., 81:109. HOLOTYPE, Adult male. Reg. no. 1936.4.12.139. Gunong Angsi, 2,600′, Negri Sembilan, Federated Malay States, Nov. 1904. Presented by the Raffles Museum, Singapore.
- friesi—Pomatorhinus tickelli friesi Delacour, 1927 32B Bull. Br. Orn. Cl., 47:160. HOLOTYPE, Adult female. Reg. no. 1927.6.5.14. Thua-Luu, central Annam, 13 Feb. 1927. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indochina.

A female syntype is also in the collection. Others are in the Vienna Museum (vide Hellmayr and Conover, Catalogue of birds of the Americas, 5 (1927): 222).

frontalis—Acanthiza frontalis Vigors and Horsfield, 1827 . . . 36
Trans. Linn. Soc. Lond., 15:226. Holotype, Adult, (relaxed mount). Reg. no. 1863.7.7.27. Australia. Presented as the type by the Linnean Society of London.

frontalis—Dendrocitta frontalis McClelland, 1839 7 in Horsfield, Proc. zool. Soc. Lond., 1839: 163. HOLOTYPE, Adult. Reg. no. 80.1.1.4527. Assam. Collected by J. McClelland. Presented by the Honourable East India Company Museum.
frontalis—Euphonia frontalis P. L. Sclater, 1851 5 in Jardine, Contributions to Ornithology: 89. HOLOTYPE, Adult male. Reg. no. 1885.6.12.17. Ecuador (=Venezuela, vide Sclater, Cat. Bds. Brit. Mus. 11 (1886): 55). Collected for Verreaux. Sclater Collection, received by him from E. Wilson. Sclater used a Bonaparte MS name.
frontalis—Oriolus frontalis Wallace, 1862
frontalis—Orthotomus frontalis Sharpe, 1877
frontalis—Sitta frontalis Swainson, 1820
frontata—Motacilla frontata Swinhoe, 1870
fruticola—Saxicola fruticola Horsfield, 1821
fucatus—Proparus fucatus Styan, 1899

fugensis—Hypsipetes fugensis Ogilvie-Grant, 1895
fugglescouchmani—Camaroptera brachyura fugglescouchmani Moreau, 1939
fugglescouchmani—Seicercus umbrovirens fugglescouchmani Moreau, 1941
fulgidum—Dicaeum fulgidum P. L. Sclater, 1883
fulignosa—Apalis melanocephala fulignosa Vincent, 1933 35 Bull. Brit. Orn. Cl., 53: 141. HOLOTYPE, Adult male. Reg. no. 1933.3.1.11. Cholo Mountain, 4,200′, Nyasaland, 16°02′ S., 35°03′ E., 2 Sept. 1932. Collected by J. Vincent. Presented by the Portuguese East Africa Expedition.
fuliginosa—Artomyias fuliginosa J. and E. Verreaux, 1855 37 J. Orn., 1855: 104. SYNTYPE, Adult. Reg. no. 1854.5.23.42. Gaboon, West Africa. Collected by and purchased of J. and E. Verreaux. Listed as the ?type by Sharpe, Cat Bds. Brit. Mus., 4 (1870): 144, and figured on pl. 3. There is another syntype in the collection.
fuliginosa—Certhia discolor fuliginosa E. C. S. Baker, 1922
fuliginosa—Geospiza fuliginosa Gould, 1837

mount). Reg. no. 1855.12.19.44. (Chatham Island, on label), Galapagos Archipelago. Collected by C. Darwin. Purchased of the Zoological Society of London.

Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 13 ("Geological Society" printed in error for "Zoological Society").

There are several other syntypes in the collection.

fuliginosa—Hemichelidon fuliginosa Hodgson, 1845 37

Proc. zool. Soc. Lond., 1845: 32. SYNTYPE, Adult. Reg. no. 1845.1.9.272. Nepal. Presented by B. H. Hodgson.

Hodgson's plate no. 427 is on the label. There are several other

syntypes in the collection. (See also appendix, Corrections and

Additions, vol. 2.)

fuliginosa—Pachycephala fuliginosa Vigors and Horsfield, 1827 . 39
Trans. Linn. Soc. Lond., 15: 241. SYNTYPE, Adult female (relaxed mount). Reg. no. 1863.7.7.42. South Coast of New Holland (=Port Lincoln, Eyre's Peninsula, vide Mathews, Birds of Australia, 8 (1920): 221). Collected by R. Brown. Presented by the Linnean Society.

This specimen was presented as the type and listed as such by Gadow, Cat. Bds. Brit. Mus. 8 (1883): 193, but two specimens were referred to in the description, only one coming to the museum.

The female syntype is in the collection.

- fuliginosus—Anthus fuliginosus Vigors and Horsfield, 1827 . . . 36
 Trans. Linn. Soc. Lond., 15:230. HOLOTYPE, Adult. Reg. no. 1863.7.7.33. Van Dieman's Land, 1804. Collected by R. Brown. Presented as the type by the Linnean Society of London.

fulvescens—Chamaeza fulvescens Salvin & Godman, 1882. Ibis, (4) 6:79. SYNTYPE, Adult male. Reg. no. 1889.7.10.825. Merume Mts., British Guiana, 22 June 1881. Collected by H. Whitely. Salvin-Godman Collection. There are two other syntypes in the collection.
fulvicauda—Suthora fulvicauda Campbell, 1892
fulviceps—Euscarthmus fulviceps P. L. Sclater, 1871 Proc. zool. Soc. Lond., 1871: 497. SYNTYPE, Adult (male, on label). Reg. no. 1888.1.13.316. 10 miles from Lima, Peru, 4 Dec. 1869. Collected by W. Nation. Sclater Collection. The "imperfect skin from Babahoyo", a syntype collected by Fraser, and listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 94, is also in the collection.
fulvicervix—Calliste fulvicervix P. L. Sclater and Salvin, 1876 5 Proc. zool. Soc. Lond., 1876: 354, pl. 30. Holotype, Adult (female, vide Hellmayr and Conover, Catalogue of birds of the Americas 9 (1936): 133). Reg. no. 1885.6.4.538. Tilotilo, Yungas Province, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.
fulvicrissa—Euphonia fulvicrissa P. L. Sclater, 1857 5 Proc. zool. Soc. Lond., 1856 (publ. 1857): 276. HOLOTYPE, Adult male. Reg. no. 1885.5.12.117. Santa Martha, New Granada. Collected for Verreaux. Sclater Collection.
fulvifacies—Abrornis fulvifacies Swinhoe, 1870
fulvifrons—Meliphaga [Myzomela] fulvifrons Vigors and Horsfield, 1827
fulvifrons—Temnoris fulvifrons Hodgson, 1845

fulvigularis—Myjobius fulvigularis Salvin and Godman, 1889 Biol. Centr.-Amer., Aves, 2:58. HOLOTYPE, Adult male. Reg. no. 1888.1.1.1224. Santa Fé, Veragua. Collected by E. Arcé. Salvin-Godman Collection fulvipectus—Cyclorhynchus fulvipectus P. L. Sclater, 1860 Proc. zool. Soc. Lond., 1860: 92. HOLOTYPE, Adult male. Reg. no. 1888.1.13.698. Nanegal, Ecuador. Collected by L. Fraser. Sclater Collection. fulvipennis—Cotyle fulvipennis P. L. Sclater, 1859 Proc. zool. Soc. Lond., 1859: 364. HOLOTYPE, (Immature, vide Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 208). Reg. no. 1884.5. 15.99. Jalapa, Vera Cruz, south Mexico. Collected by De Oca. Sclater Collection. fulviventer—Budytes fulviventer Hodgson, 1844. . 19 in J. E. Grav, Zoological Miscellany, 1844: 83 (nomen nudum). Specimens in the collection were listed as types by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 525. fulviventris—Grallaria fulviventris P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858:68. HOLOTYPE, Adult. Reg. no. 1889.9.20.642. Rio Napo, Ecuador. Probably collected by Villavicencio. Sclater Collection; obtained by Sclater through Verreaux. fulviventris—Horornis? fulviventris Hodgson 1845 Proc. zool. Soc. Lond., 1845:31. ?SYNTYPE, Adult. Reg. no. 1845.1.9.826. Nepal. Presented by B. H. Hodgson. Listed as the type by Seebohm, Cat. Bds, Brit, Mus., 5 (1881): 128. There are other syntypes in the collection. fulviventris—Myiagra fulviventris P. L. Sclater, 1883 Proc. zool. Soc. Lond., 1883:54. HOLOTYPE, Adult male. Reg. no. 1887.12.30.1128. Larat, Tenimber Islands, 2 Aug. 1882. Collected by H. O. Forbes. Tweeddale Collection. fulviventris—Turdus fulviventris P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1857: 273. HOLOTYPE, Adult male. no. 1886.8.2.121. Bogota, Colombia. Collected for Jamrach. Sclater Collection. Verreaux's MS label, mentioned by Sclater, is no longer attached. fulvolateralis—Acrocephalus fulvolateralis Sharpe, 1877 in Layard, Birds of South Africa, 2nd Edn., : 289. HOLOTYPE, Adult. Reg. no. 1876.5.23.22. Natal. Sharpe Collection. fulvotincta—Pachycephala fulvotincta Wallace, 1863 Proc. zool. Soc. Lond., 1863: 492. SYNTYPE, Adult male. Reg. no. 1863.12.16.4. Flores. Collected by Allen (op. cit., p. 480). Wallace Collection.

There are several other syntypes in the collection.

130 Mary Contest
fulvoventer—Reguloides fulvoventer Godwin-Austen, 1874 J. Asiat. Soc. Bengal, 43 pt. 2:167. HOLOTYPE, Adult. Reg. no.
1895.7.14.950. Donsiri Valley, Assam. Godwin-Austen Collection.
fulvus—Drymocataphus fulvus Walden, 1875
fulvus—Sphenoecus fulvus G. R. Gray, 1862
fulvus—Thryophilus fulvus P. L. Sclater, 1874
fumicolor—Ochthoëca fumicolor P. L. Sclater, 1856 9 Proc. zool. Soc. Lond., 1856 : 28, pl. 117. HOLOTYPE, Adult. Reg. no. 1888.1.13.50. Bogota, Colombia. Collected for Verreaux. Sclater Collection.
fumigata—Calamoherpe fumigata Swinhoe, 1863 35 Proc. zool. Soc. Lond., 1863 : 91. SYNTYPE, Adult female. Reg. no. 1886.7.8.1744. Amoy, China, 18 May 1861. Collected by R. Swinhoe. Hume Collection. This is one of the eight syntypes referred to; no male syntypes are in the collection.
fumigata—Munia fumigata Walden, 1873
fumigatus—Uragus sibiricus fumigatus Sowerby, 1920 59 Bull. Brit. Orn. Cl., 40:99. Holotype, Adult male. Reg. no. 1898.11.1.789. Tataschew, near Krasnoyarsk, South-central Siberia, 5 March 1882. Collected by J. Kibort. Seebohm Collection. This specimen was originally labelled with the date 21 Feb. 1882, but Seebohm's label has the date 5 March 1882, and this is clearly the specimen described, and the registration number is quoted.
fumosus—Automolus fumosus Salvin & Godman, 1891

- funebris—Apalis cinerea funebris Bannerman, 1936 . . . 35

 Bull. Brit. Orn. Cl., 57: 72. Holotype, Adult male. Reg. no. 1926.8.8.573. Oku, 7,000′, West of Kumbo, Banso Mountains, Cameroon, 8 Feb. 1925. Collected by and purchased of G. L. Bates
- - Two of the other syntypes are also in the collection.
- funereus—Oryzoborus funereus P. L. Sclater, 1859 59

 Proc. zool. Soc. Lond., 1859 : 378. HOLOTYPE, Adult male. Reg.
 no. 1885.12.10.92. Suchapam, Oaxaca, Mexico, April 1859. Collected by A. Boucard. Sclater Collection.

 The total length was erroneously stated to be 8.4", but is in fact 4.4".
- furensis—Amadina fasciata furensis Lynes, 1920 . . . 60

 Bull. Brit. Orn. Cl., 41: 17. HOLOTYPE, Adult male (=immature).

 Reg. no. 1920.12.22.617. Derrat, 50 miles east of El Fashar, Darfur,

 Sudan, 9 March 1920. Collected and presented by H. Lynes.
- furensis—Cercomela scotocerca furensis Lynes, 1926 31

 Ibis, (12) 2:391. SYNTYPE, Adult male. Reg. no. 1922.12.8.909.

 Jebel Marra, 7,900', central Darfur, Sudan, 6 March 1921. Collected and presented by Admiral H. Lynes.

 This is the author's selected male syntype. His selected female.
 - This is the author's selected male syntype. His selected female syntype is also in the collection.
- furensis—Ploceus [Sitagra] taeniopterus furensis Lynes, 1923 . . . 61

 Bull. Br. Orn. Cl., 43:94. HOLOTYPE, Adult male. Reg. no.
 1922.12.8.1905. Zalingei, west-central Darfur, Sudan, 2 Sept.
 1921. Collected by H. Lynes. Lynes-Lowe Expedition Collection.

Abyssinian highlands, 4 April 1868. Collected by W. T. Blanford and presented by the Governor and Council of the Bombay Government.

fusca—Alcippe fusca Godwin-Austen, 1876 J. Asiat. Soc. Bengal, 45, pt. 2:197. SYNTYPE, Adult. Reg. no. 1895.7.14.2103. Naga Hills, North-East Frontier, India, season", 1873-4. Collected by A. W. Chennell and presented by H H Godwin-Austen.

The three other syntypes are also in the collection.

fusca—Casiornis fusca P. L. Sclater and Salvin, 1873 Nomencl. Av. Neotrop., 57, 159. SYNTYPE, Adult. Reg. no. 1888. 1.13.1585 (labelled 1885 in error). Bahia, eastern Brazil. (Collected by Wucherer, on label). Sclater Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 366.

There is a second syntype in the collection.

- fusca—Certhidea fusca P. L. Sclater and Salvin, 1870. Proc. zool. Soc. Lond., 1870: 324. SYNTYPE, Adult male. Reg. no. 1885.4.1.190. Abingdon Island, Galapagos Archipelago. Collected by A. Habel. Salvin-Godman Collection. There are two other syntypes in the collection.
- fusca—Pachycephala fusca Vigors and Horsfield, 1827 Trans. Linn. Soc. Lond., 15: 240. SYNTYPE, Adult female (relaxed mount). Reg. no. 1863.7.7.40. Australia. Collected by G. Caley. Presented by the Linnean Society.

This specimen was presented as the type and listed as such by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 193, but two specimens are referred to in the description.

fusca—Petroica fusca Gould, 1842 Birds of Australia, 3: pl. 8. SYNTYPE, Adult male. Reg. no. 1841.1467. Tasmania, (near Wellington Mountain, Van Diemen's Land, Sept. 1839, in British Museum Register). Purchased of J.

Labelled "P. fusca" in Gray's hand. The author refers to many specimens and from the date of registration this seems certain to be a syntype.

- fusca—Rhinocrypta fusca P. L. Sclater & Salvin, 1873. Nomencl. Av. Neotrop.,: 161. HOLOTYPE, Adult. Reg. no. 1889. 9.20.715. Mendoza, Argentina, 1871. Collected by Weisshaupt. Sclater Collection.
- **fuscata**—Garrulax moniliger fuscata E. C. S. Baker, 1917 . . . Bull. Br. Orn. Cl., 38:64. HOLOTYPE, Adult male. Reg. no. 1886.10.1.3952. Tavoy, south Burma, 17 March 1878. Collected by A. O. Hume. Hume Collection.

- fuscata—Ortygospiza atricollis fuscata W. L. Sclater, 1932 Bull. Brit. Orn. Cl., 52:142. HOLOTYPE, Adult male. Reg. no. 1932.6.5.2. Kawambwa, Northern Rhodesia, 6 Nov. 1930. Collected and presented by H. Lynes.
- fuscicapilla—Cisticola fuscicapilla Wallace, 1864. Proc. zool. Soc. Lond., 1863 (publ. 1864): 489. SYNTYPE, Adult (male, on label). Reg. no. 1873.5.12.611. (Delhi, on label; this is a Timor locality, vide Ticehurst, Systematic review of the genus Phylloscopus (1938): 183) East Timor, (c. May 1861, on label). Collected by A. R. Wallace. Wallace Collection. There is another syntype in the collection.
- fuscicapillus—Copurus fuscicapillus P. L. Sclater, 1861. Proc. zool. Soc. Lond., 1861 (publ. 1862): 381. SYNTYPE, Adult. Reg. no. 1888.1.13.157. Bogota, Colombia. Collected for Verreaux. Sclater Collection. Sclater used a Verreaux MS. name. There are two other syntypes

in the collection.

- no. 1858.3.10.17. Aru Islands, 1857. Collected by A. R. Wallace. Purchased of S. Stevens. The female syntype is also in the collection.
- fuscicapillus—Hylophilus fuscicapillus P. L. Sclater and Salvin, 1880 56 Proc. zool. Soc. Lond., 1880:155. SYNTYPE, Adult. Reg. no. 1886.9.15.302. Sarayacu, Ecuador. Collected by C. Buckley. Sclater Collection.

There are two other syntypes in the collection.

fuscicapillus—Picolaptes fuscicapillus Pelzeln, 1868 Zur Ornithologie Brasiliens: 44, 63. SYNTYPE, Adult male. Reg. no. 1889.5.20.585. Engenho do Capitão Gama, Matto Grosso. Brazil, 20 July 1826. Collected by J. Natterer. P. L. Sclater Collection.

Listed as "authentic specimen" by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 154.

fuscicapillus—Tropidorhynchus fuscicapillus Wallace, 1862 Ibis, (1) 4:351. SYNTYPE, Adult. Reg. no. 1873.5.12.1010. Morty Island, near Gilolo, northern Molucca group, 1861. Collected by Allen. Wallace Collection.

There are several syntypes in the collection and one in the Cambridge University Zoological Museum.

fuscicaudata—Ruticilla? fuscicaudata Blanford, 1869 31 Ann. Mag. nat. Hist., (4) 4:329. HOLOTYPE, Adult male. Reg. no. 1869.10.16.215. Abyssinia (Mas'halet Pass, 4,500', near Bogos, 13 July 1868, on label). Collected by W. T. Blanford and presented by the Governor and Council of the Bombay Government.

- fusciceps—Thripophaga fusciceps P. L. Sclater, 1889 3

 Proc. zool. Soc. Lond., 1889 : 33. Syntype, Adult (relaxed mount).

 Reg. no. 1846.9.9.56. Bolivia. Collected by and purchased of T. Bridges.

 There is another syntype in the collection.
- fuscigularis—Apalis murina fuscigularis Moreau, 1937 35

 Bull. Brit. Orn. Cl., 58: 48. HOLOTYPE, Adult male. Reg. no. 1938.2.11.5. Taita Hills, 5,400′, Southern Kenya, 12 Nov. 1937. Collected by and purchased of R. E. Moreau.
- fuscipennis—Bathmocercus fuscipennis Sharpe, 1903 . . . 35

 Bull. Brit. Orn. Cl., 14: 19. HOLOTYPE, Adult male. Reg. no. 1903.7.16.16. Efulen, Cameroons, 15 Dec. 1902. Collected by G. L. Bates. Purchased of W. E. de Winton.
- fusciventris—Ptyonoprogne rufigula fusciventris Vincent, 1933. . 18
 Bull. Br. Orn. Cl., 53: 143. HOLOTYPE, Adult male. Reg. no. 1933.3.1.21. Namuli Mt., 6,200', Quelimane Province, Mozambique, 15°21' S., 37°04', E., 4 Aug. 1932. Collected and presented by J. Vincent.
- fuscoflava—Pachycephala fuscoflava P. L. Sclater, 1883 . . . 39

 Proc. zool. Soc. Lond., 1883: 198, pl. 28. SYNTYPE, Male (immature, vide Hartert, Novit. Zool., 8 (1901): 170). Reg. no. 1883.5.30.99. (allocated later). Larat Island, Timor-Laut, Tenimber Islands. 15 Sept. 1882. Collected by H. O. Forbes. Presented by the "Timor-Laut" Committee of the British Association.

 The female syntype is also in the collection.
- fuscogularis—Siphia strophiata fuscogularis E. C. S. Baker, 1923 . 37 Bull. Brit. Orn. Cl., 44:1. HOLOTYPE, Adult male. Reg. no. 1919.12.20.163. Langbian Peaks, 6,000-7,500′, South Annam, 21 April 1918. Collected by C. B. Kloss. Presented by H. C. Robinson and C. B. Kloss.

- fuscorufa—Rhipidurafuscorufa P. L. Sclater, 1883 37A

 Proc. zool. Soc. Lond., 1883: 197, pl. 27. Syntype, Adult male.

 Reg. no. 1883.5.30.81. Loetoe, Tenimber Group, 20 Sept. 1882.

 Collected by H. O. Forbes. Presented by the Timor Laut Committee of the British Association.

Fourteen syntypes were referred to, of which one other is in the

collection.

- fuscorufus—Ochthodiaeta fuscorufus P. L. Sclater and Salvin, 1876 9

 Proc. zool. Soc. Lond., 1876: 354. SYNTYPE, Adult. Reg. no.
 1888.1.1.55. Tilotilo, Yungas province, Bolivia. Collected by C.
 Buckley. Salvin-Godman Collection.

 There is a second syntype in the collection.
- gabonensis—Malaconotus gabonensis Shelley, 1894 . . . 23
 Bull. Br. Orn. Cl., 3: 43. SYNTYPE, Adult (relaxed mount). Reg.
 no. 1865.4.9.65. Gaboon, West Africa. Collected for Verreaux
 and purchased of R. B. Sharpe.

This name is based on the specimens called Laniarius hypopyrrhus (nec Hartlaub), by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 155.

There are several other syntypes in the collection.

gabonensis—Ortygospiza gabonensis Lynes, 1914 60
Bull. Brit. Orn. Cl., 33:131. Syntype, Adult male. Reg. no.
1878.12.31.880. Gaboon, West Africa. Collected by P. B. Du
Chaillu. Sharpe Collection.

This is the author's selected male syntype. The described female

syntype is also in the collection.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 148.

- gaboonensis—Estrilda astrild gaboonensis W. L. Sclater and 453. Landana, Congo, Jan. 1883. Collected by L. Petit, Shelley Collection. gabriellae—Cissa gabriellae Ogilvie-Grant, 1906 Bull. Brit. Orn. Cl., 19:12. HOLOTYPE, Adult. Reg. no. 1907. 12.13.24. Nhatrang, Annam. Collected by and purchased of J. J. Vassal gabun—Cisticola avresii gabun Lynes, 1931 Bull. Brit. Orn. Cl., 52:9. HOLOTYPE, Adult male. Reg. no. 1931.10.8.5. Port Gentil, coastal Gabun, 17 April 1931. Collected by Admiral H. Lynes and J. Vincent. Presented by Admiral H. Lvnes. gaikwari—Sylviella gaikwari Sharpe, 1901 . 35 Bull. Brit. Orn. Cl., 11:47. HOLOTYPE, Adult female. Reg. no. 1901.4.20.15. Ania, Somaliland, (3 Feb. 1899, on label). Collected by Dr. Donaldson Smith. Presented by the Gaekwar of Baroda, the collector having made the expedition to Somaliland on his behalf. galbanus—Garrulax galbanus Godwin-Austen, 1874 Proc. zool. Soc. Lond., 1874: 44, pl. 10. HOLOTYPE, Adult. Reg. no. 1895.7.14.2216. Munipur Valley, under the Koupru Range, Assam, Feb. 1873 (in text; Jan. on label). Collected by H. H. Godwin-Austen. Godwin-Austen Collection. galeata-Myiagra galeata G. R. Gray, 1860 Proc. zool. Soc. Lond., 1860: 352. HOLOTYPE, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 385). Reg. no. 1860. 2.4.53. Batchian, Moluccas. Collected by A. R. Wallace. Purchased of Stevens. gallio—Zosterops gallio Sharpe, 1884. in Gadow, Cat. Bds. Brit. Mus., 9:185. HOLOTYPE, Adult female. Reg. no. 1873.5.12.904. East Java, 1861. Collected by A. R. Wallace. Wallace Collection.
- gambiensis—Emberiza affinis gambiensis Bannerman, 1932 . . . 49
 Bull. Brit. Orn. Cl., 52:111. HOLOTYPE, Adult male. Reg. no. 1929.2.18.155. Old Nokunda, Gambia, 22 Jan. 1929. Collected and presented by W. P. Lowe.

Bull. Brit. Orn. Cl., 12:11. HOLOTYPE, Adult male. Reg. no. 1911.12.23.1373. Gambaga, Gold Coast, 27 April 1901. Col-

gambagae—Alseonax gambagae Boyd Alexander, 1901.

lected by Boyd Alexander. Boyd Alexander Bequest.

ganeesa—Hypsipetes ganeesa Sykes, 1832	. 21
Proc. zool. Soc. Lond., 1832:86. SYNTYPE, Adult. Reg.	no.
1880.1.1.4746. Dense woods of the Ghauts. Collected by	Lt.
Col. W. Sykes and presented by the Honourable East India Co	om-
pany Museum.	
The other syntype is also in the collection. See Jardine a	and

Selby, *Illustrations of Ornithology*, 2nd. ser., 4 (1836): pl. 2.

- garoensis—Turdinus garoensis Godwin-Austen, 1874 32B J. Asiat. Soc. Bengal, 43, pt. 2:160, pl. 8. Holotype, Adult. Reg. no. 1895.7.14.3399. Garo Hills, North East Frontier, India, 1872–3. Collected by W. Robert and presented by H. H. Godwin-Austen.

There are two other syntypes in the collection, and a fourth in the Academy of Natural Sciences, Philadelphia.

- gayeti—Melanochlora gayeti Delacour and Jabouille, 1925 . . . 42

 Bull. Brit. Orn. Cl., 46: 6. SYNTYPE, Adult male. Reg. no. 1926.

 9.8.958. Bana, 1,900 metres (1,400, on label), 12 miles west of Fourane, Central Annam, 24 May 1924. Collected by J. Delacour and P. Jabouille, from the French British Indo-China Expedition.

 The female syntype is also in the collection.

This specimen was labelled "n. sp." by the author, but a large number of syntypes collected at the same period are in the collection.

genderuensis—Cercomela familiaris genderuensis Bannerman, 1922 31 Bull. Br. Orn. Cl., 43:8. HOLOTYPE, Adult male. Reg. no. 1922. 11.25.133. Genderu, 4,600', Cameroon Highlands, 28 Aug. 1921. Collected by and purchased of G. L. Bates. geochrous—Malacocircus geochrous Hodgson, 1844 in J. E. Gray, Zoological Miscellany: 83. Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 393 refers to two specimens as types, but the name is a nomen nudum. gephyra—Sylvietta ruficapilla gephyra White, 1953 Bull. Brit. Orn. Cl., 73:68. HOLOTYPE, Adult male. Reg. no. 1953.54.322. Mwinilunga, Northern Rhodesia, 26 Jan. 1940. Collected and presented by C. M. N. White. germana—Cyanocitta germana P. L. Sclater and Salvin, 1876. Proc. zool. Soc. Lond., 1876: 270. SYNTYPE. Adult female. no. 1887.3.9.140. Belize River, British Honduras. Collected by Ouilter. Salvin-Godman Collection. A male syntype is in the collection. gigantea—Locustella lanceolata gigantea Johansen, 1954 J. Orn., 1954: 92. The author selected as the type specimen no. 1912.12.15.186. from the H. F. Witherby Collection, from Shaweishan Island, E. China. This cannot now be traced but other specimens from a paratypical series are in the collection. gigas—Elainea gigas P. L. Sclater, 1871 Proc. zool. Soc. Lond., 1870 (publ. 1871): 831. HOLOTYPE, Adult. Reg. no. 1888.1.13.537. Rio Napo, Ecuador. Collected for Verreaux. Sclater Collection. 49 Amoy, China, May 1866. Collected by R. Swinhoe. Seebohm Collection. There are several other syntypes in the collection. gilberti—Kupeornis gilberti Serle, 1949 Bull. Br. Orn. Cl., 69:50. HOLOTYPE, Adult male. Reg. no. 1949.14.1. Kupé Mountain, 5,000', Kumba Division, British Cameroons, 10 April 1948. Collected and presented by W. Serle. gili—Cyanopica cyanus gili Witherby, 1923. Bull. Brit. Orn. Cl., 43:74. HOLOTYPE, Adult male. Reg. no. 1924.12.20.4. Near Candeleda, north-west Avila Province, Spain, 29 Oct. 1922. Collected and presented by H. F. Witherby. gilletti—Mirafra gilletti Sharpe, 1895. Bull. Br. Orn. Cl., 4:29. HOLOTYPE, Adult. Reg. no. 1895.7.7.17. Sibbe, Ogaden Country, Somaliland (i.e. 25 miles west of Sasah Baneh, east Abyssinia, vide Smith, Geog. J., 1900:712), 4 Aug. 1894. Collected by F. Gillett and presented by Dr A. Donaldson Smith.

- glaber—Sublegatus glaber P. L. Sclater and Salvin, 1868 . . . 9

 Proc. zool. Soc. Lond., 1868: 172, pl. 13. Adult (male, on label).

 Reg. no. 1888.1.13.650. Caraccas, Venezuela (Aug. 1859, on label).

 Collected by O. Goering. Sclater Collection.

 Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 158.

 There are two other syntypes in the collection.

Gould appears to have retained the specimen, which is not listed by Stone, *Proc. Acad. Nat. Sci. Philadelphia* 1899: 62–63, as one of the types in the Wilson Collection at the Philadelphia Museum.

- glabrirostris--Melanoptila glabrirostris P. L. Sclater, 1857 . . . 29

 Proc. zool. Soc. Lond., 1857: 275. SYNTYPE, Adult. Reg. no. 1886.8.2.199. Omoa, Honduras. Collected by J. Leyland and purchased of Sclater.
- glauca—Diglossa glauca P. L. Sclater and Salvin, 1876 . . . 51 Proc. zool. Soc. Lond., 1876: 253. HOLOTYPE, Adult male. Reg. no. 1885.4.1.74. Nairapi, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.

Peters, Checklist of birds of the World 12 (1967): 55 accepts the description of the species under the name of Orites glaucogularis by Moore, Proc. zool. Soc. Lond., 1854: 140, but according to Baker, Fauna of British India, 2nd Bdn. 7 (1930): 16, Moore's account was not published until 11 April 1855 and Gould's was published on 1 April.

- gloriosa—Diglossa gloriosa P. L. Sclater and Salvin, 1871 . . . 51

 Proc. zool. Soc. Lond., 1870 (publ. 1871): 784, pl. 46. HOLOTYPE,

 Adult male. Reg. no. 1884.7.31.11. Parama de la Culata, north of

 Merida, Venezuela, 1869. Collected by A. Goering. Sclater

 Collection.
- gloriosus—Rhynchocyclus flaviventris gloriosus Chubb, 1920 . . . 9
 Bull. Brit. Orn. Cl., 40: 108. SYNTYPE, Adult male. Reg. no. 1892.1.16.254. Quonga, British Guiana, 21 Nov. 1887. Collected by H. Whitely, Salvin-Godman Collection.

 There are two female syntypes in the collection.
- goalpariensis—Certhia goalpariensis Royle, 1840 46
 Illustrations of Botany . . ., : 78, pl. 7. HOLOTYPE, Adult male.

Reg. no. 1880.1.1.4055. Dhera Doon, Himalayas. Collected by Dr. J. F. Royle. Presented by the Honourable East India Com-

pany Museum as Dr. Royle's type.

Royle made no written description but this is his only specimen and therefore assumed to be the one on which the plate was based. The name appears to be a latinised name of Latham's, *Gen. Hist. Bds.* 4 (1822): 221. The specimen was listed as the type by Gadow, *Cat. Bds. Brit. Mus.*, 9 (1884): 20. It is in bad condition with a broken beak.

The second syntype is in the collection.

- godwini—Pomatorhinus ruficollis godwini Kinnear, 1944 . . . 32B Ibis, 86:79. Holotype, Adult male. Reg. no. 1935.4.5.98. Chungkar, 6,000', Trashigong-Duwengiri road, south-east Bhutan, 8 Nov. 1934. Collected and presented by F. Ludlow and G. Sherriff.
- godwini—Trochalopterum erythrocephalum godwini Harington, 1914 32A Bull. Br. Orn. Cl., 33:92. HOLOTYPE, Adult. Reg. no. 1895.7.14. 2242. Hengdan Peak, North Cachar Hills. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.
- goeringi—Chlorospingus goeringi P. L. Sclater and Salvin, 1871 . 51 Proc. zool. Soc. Lond., 1870 (publ. 1871): 784, pl. 46. HOLOTYPE, Adult male. Reg. no. 1885.6.12.885. Paramos of Merida, Venezuela, 1870. Collected by A. Goering. Sclater Collection.
- golandi—Heterhyphantes golandi Clarke, 1913 61

 Bull. Br. Orn. Cl., 31:32. HOLOTYPE, Adult male. Reg. no. 1913.1.17.1. Mombasa, Kenya. Collected and presented by S. R. Clarke.
- goodenovii—Muscicapa goodenovii Vigors and Horsfield, 1827 . 37 Trans. Lim. Soc. Lond., 15:245. Holotype, Adult (male, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879) 172) (relaxed mount). Reg. no. 1863.7.7.44. Coast of South Australia, 1802. Collected by R. Brown. Presented by the Linnean Society as the type.

- goodfellowi—Cicinnurus goodfellowi Ogilvie-Grant, 1907 . . . 70

 Bull. Brit. Orn. Cl., 19: 39. HOLOTYPE, Adult male. Reg. no. 1908.5.27.74. Cyclops Mountains, 3,000', Humboldt's Bay, north Dutch New Guinea, Aug. 1906. Collected by W. Goodfellow.
- goodfellowi—Erythrura trichroa goodfellowi Ogilvie-Grant, 1911 60 Bull. Brit. Orn. Cl., 29:29. SYNTYPE, Adult male. Reg. no. 1913.3.6.358. Moroka Mountains, 4,000', British New Guinea, Feb. 1908. Collected by W. Goodfellow. Purchased of E. J. Johnstone.

The female syntype is also in the collection.

- goodfellowi—Ianthia goodfellowi Ogilvie-Grant, 1912 31

 Ibis, (9) 6:649, pl. 14. SYNTYPE, Adult male. Reg. no 1913.1.

 29.51. Mount Arizan, 7,000', Formosa. Feb. 1912. Collected by W. Goodfellow and purchased of E. J. Johnstone.

 The female syntype is also in the collection.
- goodfellowi—Pitta novaeguineae goodfellowi C. M. N. White, 1937. 12

 Bull. Brit. Orn. Cl., 57: 136. HOLOTYPE, Adult male. Reg. no. 1907.12.11.63. Silbattabatta, Aru Island, Feb. 1904. Collected by W. Goodfellow. Purchased of E. J. Johnstone.
- goodfellowi—Regulus goodfellowi Ogilvie-Grant, 1960 . . . 35

 Bull. Brit. Orn. Cl., 16: 122. SYNTYPE, Adult male. Reg. no. 1907.12.12.20. Mount Morrison, 10,000', Central Formosa, (Jan. 1906, on label). Collected by and purchased of W. Goodfellow. The female syntype is also in the collection.
- goodfellowi—Rhinomyias goodfellowi Ogilvie-Grant, 1905 . . . 37

 Bull. Brit. Orn. Cl., 16:17. HOLOTYPE, Adult female. Reg. no. 1905.11.26.106. Mount Apo, 8,000′, South-east Mindanao, Philippines, March 1905. Collected by and purchased of W. Goodfellow.

goslingi—Apalis goslingi Boyd Alexander, 1908 35 Bull. Brit. Orn. Cl., 21:89. Holotype, Adult male. Reg. no. 1911.12.23.2211. Guraba, River Welle (Uele), Belgian Congo, 17 May 1906. Collected by W. Boyd Alexander on the Alexander-Gosling Expedition. Boyd Alexander Bequest.
goslingi—Fringillaria goslingi Boyd Alexander, 1906
gossei—Embernagra gossei Chubb, 1918
gouldi—Calliste gouldi P. L. Sclater, 1886
gouldii—Meliphaga gouldii. See appendix, Corrections and Additions, vol. 2
gouldi—Zosterops gouldi Bonaparte, 1851
gouldii—Anthus gouldii Fraser, 1843
There is another specimen in the collection, labelled "type" and "West Africa", with "(Cap Palmas)" added later. This second specimen is typical of Liberian populations in being markedly paler. Both were listed as the types by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 559.
gouldii—Manucodia gouldii G. R. Gray, 1859
H.M.S. Rattlesnake. Gray's name is proposed for a specimen figured by Gould. Gould, Handbook of the birds of Australia, 1 (1865): 236, states that his plate was based on this specimen.
gouldii—Monarcha gouldii G. R. Gray, 1860

The other syntype is also in the collection. The specimen referred to as a paratype by Hindwood, *Records of the Australian Museum*, 21 (1946): 391, is not likely to have been seen by Gould.

- gracilipes—Tyranniscus gracilipes P. L. Sclater and Salvin, 1868

 Proc. zool. Soc. Lond., 1867 (publ. 1868): 981. HOLOTYPE, Adult male. Reg. no. 1888.1.13.491. Pebas, Peru (25 June 1866, on label). Collected by H. Hauxwell. Sclater Collection.

 Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 133.
- gracilirostris—Andropadus gracilirostris Strickland, 1844 . . . 2
 Proc. zool. Soc. Lond., 1844 : 101. HOLOTYPE, Adult male. Reg.
 no. 1847.1.18.11. Fernando Po, June. Collected by and purchased of L. Fraser.
- gracilirostris—Catharus gracilirostris Salvin, 1865 31 Proc. zool. Soc. Lond., 1864 (publ. 1865) : 580. SYNTYPE, Adult. Reg. no. 1885.3.2.50. Volcan de Cartago, Costa Rica, 1864. Collected by E. Acré. Salvin-Godman Collection. Listed as type by Hellmayr, Catalogue of Birds of the Americas, 7 (1934) : 475.
- gracilirostris—Chloropeta gracilirostris Ogilvie-Grant, 1906 . 35

 Bull. Brit. Orn. Cl., 19: 33. HOLOTYPE, Immature male. Reg.
 no. 1906.12.23.1465. (Mokia, on label), 3,400′, South-eastern
 slopes of Ruwenzori, Uganda, 12 June 1906. Collected by D.
 Carruthers. Presented by subscribers to the Ruwenzori Expedition.
 Subsequently designated as type of the Genus Calamonastides,
 vide Grant and Mackworth-Praed, Bull. Brit. Orn. Cl., 60 (1940): 91.

gracilirostris—Vireo gracilirostris Sharpe, 1890 56 J. Linn. Soc. Lond., (Zool) 20: 478. SYNTYPE, Adult. Reg. no. 1888.6.1.27. Fernando Noronha Island, Brazil, Aug. 1887. Collected by H. N. Ridley. Presented by the Royal Society. This specimen was labelled as the type. There are three other syntypes in the collection.
gracilis—Aphelocoma gracilis Miller, 1896
gracilis—Hypsipetes gracilis McClelland, 1840
gracilis—Ptilotis gracilis Gould, 1866
gracilis—Pyctorhis gracilis Styan, 1899
granadensis—Dendroeca granadensis Sharpe, 1885
grandior—Turdinus marmoratus grandior Voous, 1950 32E Limosa, 22 1949, (publ. 1950): 351. HOLOTYPE, Adult female. Reg. no. 1908.12.15.94. Semangko Pass, 2,500–4,500', Selangor Pahang Boundary, 28 Nov. 1908 (Feb. on label). Collected by H. C. Robinson and presented by the Kuala Lumpur Museum.
grandis—Acridotheres grandis Moore, 1856

grandis—Bradvoterus grandis Ogilvie-Grant, 1917 35 Ibis, (10) 5:78. HOLOTYPE, Adult male. Reg. no. 1920.6.26.506. Bitye, 2,000', Ja River, Cameroons, 24 March 1914. Collected and presented by G. L. Bates. 5.6.569. Yunnan, western China. Collected by F. W. Styan. Stvan Collection. There are several other syntypes in the collection. grandis—Hyphantornis grandis G. R. Gray 1849. The Genera of birds 2:351. New name for Ploceus collaris Fraser. 1842 (q.v.) which is preoccupied. grandis—Motacilla grandis Sharpe, 1885 Cat. Bds. Brit. Mus., 10:492. SYNTYPE, Adult male. Reg. no. 1884.1.25.4. Sapporo, Yezo, Japan, 30 Sept. 1882. Collected and presented by T. Blakiston. There are four other syntypes in the collection. no. 1882.3.8.71. Taburi, Astrolabe Mountains, South-east New Guinea. Collected by A. Goldie. Purchased of E. Gerrard. grandis—Stizorhina grandis Ogilvie-Grant, 1910. Bull. Br. Orn. Cl., 27:30. HOLOTYPE, Adult. Reg. no. 1910. 12.26.173. Gazi, Shimba Hills, near Mombasa, British East Africa, 28 Aug. 1910. Collected by R. Kemp and presented by C. D. Rudd. granti—Bradyornis granti Bannerman, 1911 Bull. Brit. Orn. Cl., 27:84. SYNTYPE, Adult male. Reg. no. 1912.10.15.551. Gibbe River, South-west Abyssinia, 28 April 1905. Collected by P. Zaphiro. Presented by W. N. McMillan. This is the described male. The described female syntype is also in the collection. granti—Bradypterus usambarae granti Benson, 1939 35 Bull. Brit. Orn. Cl., 59:110. HOLOTYPE, Adult male. Reg. no. 1933.3.1.1031. Lichenya Plateau, Mlanje Mountains, Nyasaland, 31 Dec. 1931. Collected and presented by J. Vincent. granti—Camaroptera granti Boyd Alexander, 1903 . 35 Bull. Brit. Orn. Cl., 13:36. HOLOTYPE, Adult male. Reg. no. 1911.12.23.2141. Badasou, Fernando Po, 11 Nov. 1902. Col-

lected by Boyd Alexander. Boyd Alexander Bequest.

2:537. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.2098. Sumatra (locality queried by Sharpe, *Cat. Bds. Brit. Mus.*, 13 (1890): 91, 92). Collected by Sir S. Raffles. Presented by the Honour-

able East India Company Museum.

There is another syntype in the collection.

granti—Cossypha insulana granti Serle, 1949 31 Bull. Br. Orn. Cl., 69:53. HOLOTYPE, Adult female. Reg. no. 1949.14.5. Kupé Mountain, 4,500', Essosong, Kumba Division, British Cameroons, 18 Nov. 1947. Collected and presented by W. Serle.
granti—Cyanomitra olivacea granti Vincent, 1934
granti—Malimbus malimbicus granti Bannerman, 1943 61 Bull. Br. Orn. Cl., 64:6. HOLOTYPE, Adult male. Reg. no. 1909.8.5.228. Ndalla Tando, north Angola, West Africa, 12 Dec. 1908. Collected by and purchased of W. J. Ansorge.
granti—Neocossyphus granti Boyd Alexander, 1908 31 Bull. Br. Orn. Cl., 23:15. Holotype, Adult female. Reg. no. 1911.12.23.2088. Beritio, Welle River, 14 Feb. 1906. Collected by Boyd Alexander on the Alexander-Gosling Expedition. Boyd Alexander Bequest.
granti—Pellorneum ruficeps granti Harington, 1913 32E Bull. Br. Orn. Cl., 33:81. HOLOTYPE, Adult male. Reg. no. 1886.10.1.1421. Mynall, Travancore, 10 March 1877. Hume Collection.
granti—Pyrenestes granti Sharpe, 1908
granti—Rhipidura rufifrons granti Hartert, 1918 37A Bull. Brit. Orn. Cl., 38:60. New name for R. harterti Ogilvie- Grant, 1915 (q.v.), on the grounds of its preoccupation by R. flabellifera harterti Mathews, 1912. Since this is a new name the type specimen designated by Hartert for it has no type status.
granviki—Melocichla mentalis granviki C. H. B. Grant and Mackworth-Paed, 1941
gratiosus—Mecocerculus gratiosus P. L. Sclater, 1862
1888.1.13.85. Highlands of Ecuador. Collected for J. Gould. Sclater Collection.

no. 1876.5.23.576. Omaruru River, Damaraland, 30 Oct. 1966.
Collected by C. J. Andersson. Sharpe Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 350. The second syntype referred to in the description is also in the collection.
grayi—Cisticola grayi Walden, 1872
grayi—Nectarinia grayi Wallace, 1865
grayi—Todopsis grayi Wallace, 1862
grayi—Zosterops grayi Wallace, 1864
graysoni—Quiscalus graysoni P. L. Sclater, 1884
greenwayi—Erythropygia barbata greenwayi Moreau, 1938 31 Bull. Br. Orn. Cl., 58: 64. HOLOTYPE, Adult Male. Reg. no. 1938.1.4.1. Kange, Mafia Island, Tanganyika, 11 Aug. 1937. Collected for and presented by R. E. Moreau.
gregoriana—Oreocincla gregoriana Hume, 1873
greyi—Poeoptera greyi Jackson, 1899

- grimwoodi—Cossypha polioptera grimwoodi C. M. N. White, 1954. 31 Bull. Br. Orn. Cl., 74:88. HOLOTYPE, Adult male. Reg. no. 1954.34.1. Source of the Zambesi River, Mwinilunga district, Northern Rhodesia, 23 July 1954. Collected and presented by Major I. R. Grimwood.
- griscomi—Spinus notatus griscomi van Rossem, 1938 59
 Bull. Brit. Orn. Cl., 58:135. Holotype, Adult male. Reg. no.
 1899.2.1.619. Volcan de Colima, Jalisco, Mexico, 31 Dec. 1889.
 Collected by W. B. Richardson. Salvin-Godman Collection.
- grisea—Scotocerca inquieta grisea Bates, 1936 35

 Bull. Brit. Orn. Cl., 57: 21. HOLOTYPE, Adult male. Reg. no. 1935.1.5.289. Mafraq Buraim, 4,000′, eastern edge of Taif Plateau, Arabia, 16 Nov. 1934. Collected and presented by Col. H. B. Philby.
- griseata—Ianthocincla subunicolor griseata Rothschild, 1921 . . . 32B Novit. Zool., 28:33. HOLOTYPE, Adult (male on label, female in text). Reg. no. 1921.4.21.225. Shweli-Salwin Divide, 10,000′, Yunnan, Aug. 1919. Collected by G. Forrest and presented by Stephenson Clarke.
- griseatus—Myiolestes griseatus G. R. Gray, 1858 39

 Proc. zool. Soc. Lond., 1858: 180. SYNTYPE, Adult (relaxed mount).

 Reg. no. 1850.7.20.194. Cape York, Australia. Collected by J. Macgillivray. Presented by Captain R. Stanley RN, from the voyage of H.M.S. Rattlesnake.

 The syntype from Dunk Island is also in the collection.
- griseicapillus—Lanius collurioides griseicapillus Delacour, 1926. 23

 Bull. Br. Orn. Cl., 47:13. SYNTYPE, Adult male. Reg. no. 1926.9.8.12. Xieng-Khouang, Laos, 21 Dec. 1925. Collected by J. Delacour on the Second Franco-British Expedition to French Indo china.

The female syntype is also in the collection.

- griseiceps—Basileuterus griseiceps P. L. Sclater and Salvin, 1868. 54
 Proc. zool. Soc. Lond., 1868: 170. HOLOTYPE, Adult male. Reg.
 no. 1884.5.15.437. Caripé, Venezuela, June 1867. Collected by A.
 Goering. Sclater Collection.

There is also a female syntype in the collection.

- 9.1.1443. East Pegu Hills, 15 April 1873. Collected by E. W. Oates. Hume Collection. There is another syntype in the collection. griseiceps—Pachycephala griseiceps G. R. Gray, 1858 Proc. zool. Soc. Lond., 1858: 178. HOLOTYPE, Adult male. Reg. no. 1858.3.10.70. Aru Islands. Collected by A. R. Wallace. Wallace Collection. griseiceps—Piprites griseiceps Salvin, 1865. Proc. zool. Soc. Lond., 1864 (Publ. 1865): 583. HOLOTYPE, Adult. Reg. no. 1888.1.20.10. Tucurrique, Costa Rica. Collected by E. Arcé. Salvin-Godman Collection. griseiceps—Platyrhynchus griseiceps Salvin, 1897. Bull. Brit. Orn. Cl., 7:15. SYNTYPE, Adult male. Reg. no. 1896. 11.27.30. Aunai, (Ourumee, on label). British Guiana, 9 Oct. 1890. Collected by H. Whitely. Salvin-Godman Collection. There are two other syntypes in the collection. griseiceps—Saxicola griseiceps Blanford & Dresser, 1874 Proc. zool. Soc. Lond., 1874: 233, pl. 37. SYNTYPE, Adult male. Reg. no. 1876.5.23.415. Colesberg, South Africa. Collected by T. C. Atmore. Sharpe Collection.

 Listed as type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 379.
- griseiceps—Tyranniscus griseiceps P. L. Sclater and Salvin, 1871 . 9
 Proc. zool. Soc. Lond., 1870 (publ. 1871): 841. SYNTYPE, Adult.
 Reg. no. 1888.1.13.497. Babahoyo, Ecuador (Aug. 1859, on label).
 Collected by L. Fraser. Sclater Collection.
 Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 134.
 The second syntype is also in the collection.
- griseigularis—Alseonax griseigularis Jackson, 1906 37 Bull. Brit. Orn. Cl., 19:19. HOLOTYPE, Adult male. Reg. no. 1907.6.19.3. Kibiran, Toro, Uganda (3 Oct. 1905, on label). Collected and presented as the type by F. J. Jackson.
- griseigularis—Corythocichla griseigularis Delacour & Jabouille, 1928 32B Bull. Br. Orn. Cl., 48:131. HOLOTYPE, Adult male. Reg. no. 1928.6.26.6. Bokor, 1,000 metres, south Cambodia, 8 Dec. 1927. Collected by J. Delacour and P. Jabouille on the Fourth Franco-British Expedition to Indochina.

- griseipectus—Thryothorus griseipectus Sharpe, 1881 28

 Cat. Bds. Brit. Mus., 6:236, pl. 15. Holotype, Adult male.

 Reg. no. 1869.4.10.3. Nauta, Loreto, Peru. Collected by and purchased of E. Bartlett.
- griseisticta—Hemichelidon griseisticta Swinhoe, 1861 37

 Ibis, (1) 3:330. Syntype, Adult female. Reg. no. 1897.7.2.254.

 Amoy, South China, 19 May 1859. Collected by R. Swinhoe. Seebohm Collection.

This specimen was labelled as the type when received and was the first specimen collected; but the author did not designate a type and other specimens, which may have been available, are in the collection.

- griseistriata—Cossypha griseistriata Sharpe, 1902 . . . 31

 Bull. Br. Orn. Cl., 13:8. HOLOTYPE, Adult female. Reg. no. 1902.12.8.9. Near Kangow's, Toro, west Uganda, 11 March 1902. Collected and presented by F. J. Jackson.
- griseiventer—Hemixus tickelli griseiventer Robinson & Kloss, 1919 21 Ibis, (11) 1:568. Syntype, Adult male. Reg. no. 1919.12.20.235. Langbian Peaks, 5,500-7,500', south Annam, 13 April 1918. Collected by C. B. Kloss and presented by the latter and H. C. Robinson.

This is the authors' selected male. Their selected female syntype is also in the collection.

griseiventris—Eupectes macrocercus griseiventris E. C. S. Baker, 1917 32A Bull. Br. Orn. Cl., 38:8. SYNTYPE, Adult male. Reg. no. 1916. 12.27.639. Tung Song Peninsular, Siam, 17 Sept. 1915. Collected and presented by E. G. Herbert.

The female syntype is also in the collection.

- griseocularis—Sublegatus griseocularis P. L. Sclater and Salvin, 1876 Proc. zool. Soc. Lond., 1876: 17. Syntype, Adult (male, on label). Reg. no. 1888.1.13.653. Maranura, 4,000', Urubamba, Peru, (23 Sept. 1874, on label). Collected by H. Whitely. Sclater Collection.

There are three other syntypes in the collection.

collection

- griseogenys—Leucosticte griseogenys Gould, 1843 59
 Proc. zool. Soc. Lond., 1843: 104. HOLOTYPE, Adult (relaxed mount). Reg. no. 1844.1.23.5. Vicinity of Sitka, Russian America, (vide Gould, in Hinds, Zoology of the voyage of H.M.S. Sulphur 1836-42 (1844): 49). Collected by Dr. Hinds. Presented by J. Gould.

The two other syntypes are in the collection.

- griseogularis—Monarcha griseogularis G. R. Gray, 1858 . . . 38

 Proc. zool. Soc. Lond., 1858: 177. HOLOTYPE, male (immature, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 423). Reg. no. 1858.

 3.10.51. Aru Islands, 1857. Collected by A. R. Wallace. Wallace Collection.
- griseogularis—Pericrocotus griseogularis Swinhoe, 1862. . . . 20
 Proc. zool. Soc. Lond., 1862: 282. Syntype, Adult male. Reg.
 no. 1863.2.16.12. North Formosa. Collected by R. Swinhoe and purchased of S. Stevens.

There are several other syntypes in the collection.

griseomurina—Synallaxis griseomurina P. L. Sclater, 1882 3

Proc. zool. Soc. Lond., 1882 : 578, pl. 43. HOLOTYPE, Adult. Reg. no. 1889.5.14.95. San Lucas, Ecuador. Collected by Villagomez for C. Buckley. Salvin-Godman Collection.

- griseonota—Pachycephala griseonota G. R. Gray, 1862 . . . 39

 Proc. zool. Soc. Lond., 1861 (publ. 1862): 429. HOLOTYPE, male (adult, vide Ogilvie-Grant, Bull. Brit. Orn. Cl., 25 (1910): 96). Reg. no. 1861.12.11.63. Mysol Island, Moluccas (=Ceram, vide Ogilvie-Grant, op. cit.). Collected by A. R. Wallace. Wallace Collection.
- griseonota—Zosterops griseonota G. R. Gray, 1859 47

 Proc. zool. Soc. Lond., 1859: 161. SYNTYPE, Adult male. Reg.
 no. 1859.4.19.5. Island of Nu, Port de France, New Caledonia.
 Collected by J. Macgillivray. Purchased of Cuming.
 A female syntype is also in the collection.
- griseonucha—Grallaria griseonucha P. L. Sclater & Salvin, 1870 4
 Proc. zool. Soc. Lond., 1870: 786. HOLOTYPE, Adult female. Reg.
 no. 1889.9.20.626. Paramo of La Culata, Merida, Venezuela,
 Sept. 1869. Collected by A. Goering. Sclater Collection.
- griseorufescens—Corydalla griseorufescens Hume, 1870 . . . 19

 Ibis, (2) 6: 286. SYNTYPE, Adult male. Reg. no. 1887.2.1.2837.

 Almorah, India, 29 Aug. 1868. Collected by W. E. Brooks.

 Hume Collection.
- griseotincta—Yuhina gularis griseotincta Rothschild, 1921 . . . 32B Novit. Zool., 28:42. HOLOTYPE, Adult male. Reg. no. 1921. 7.15.338. Shweli-Salwin Divide, 10-11,000′, 25°40′ N., Yunnan, June 1919. Collected by G. Forrest and presented by Stephenson Clarke.
- griseotincta—Zosterops griseotincta G. R. Gray, 1858 47

 Proc. zool. Soc. Lond., 1858: 175. Holotype, Adult male. Reg.
 no. 1851.10.11.13. Louisade Archipelago, New Guinea. Collected and presented by J. Macgillivray, from the voyage of H.M.S.

 Rattlesnake.
- grisescens—Drymocataphus tickelli grisescens Ticehurst, 1932. . . 32B Bull. Br. Orn. Cl., 53:18. HOLOTYPE, Adult male. Reg. no. 1932.10.20.2. Nyaunggyo stream, 2,500′, Taungup-Prome road, Arakan Yoma, Burma, 13 April 1929. Collected and presented by J. K. Stanford.

- grisescens—Rhynchocyclus grisescens Chubb, 1910 . . . Ibis, (9) 4:588. HOLOTYPE, Adult female. Reg. no. 1905.10. 12.640. Sapucay, Paraguay, 29 March 1903. Collected by and purchased of W. Foster. Presented by the Honourable East India Company Museum as the griseus—Sittasomus griseus Jardine, 1847 . Ann. Mag. nat. Hist., (1) 19:82. Syntype, Adult male. Reg. no. 1886.6.24.443. Tobago. Collected by Kirk. Jardine Collection: purchased of Gerrard. There are several other syntypes in the collection. grisiventris—Sitta europaea grisiventris Kinnear, 1920. Bull. Brit. Orn. Cl., 40: 142. HOLOTYPE, Adult. Reg. no. 1905. 9.10.488. Mount Victoria, Chin Hills, Burma, 24 March 1904. Collected and presented by G. Rippon. grotei-Serinus mozambicus grotei W. L. Sclater and Mackworth-Mackworth-Praed, 1918, (q,v), on the grounds of its preoccupation by Emberiza? [=Serinus] aurifrons Blyth, 1847. gualanensis—Icterus gualanensis C. F. Underwood, 1898 Bull. Brit. Orn. Cl., 7:59. HOLOTYPE, Adult male. Reg. no. 1898.12.15.1136. Gualen (on Motagua River), east Guatemala, 11 July 1897. Collected by C. F. Underwood, Salvin-Godman Collection. Now regarded as a mutant from of I. c. chrysater, vide Griscom, Bull. Amer. Mus. Nat. Hist. 64 (1932): 394. 1888.1.13.337. Gualaquiza, Ecuador, Jan. 1858. Collected by L. Fraser. Sclater Collection. The female syntype is also in the collection. guatemalae—Setophaga picta guatemalae Sharpe, 1885. Cat. Bds. Brit. Mus., 10:417. SYNTYPE, Adult male. Reg. no. 1880.12.31.42. Guatmala. Collected by T. Bridges. Eyton Collection.
- guatemalensis—Chlorophanes guatemalensis P. L. Sclater, 1861 . 51 Proc. zool. Soc. Lond., 1861:129. SYNTYPE, Adult male. Reg. no. 1884.7.31.126. Guatemala, (Choctum, Vera Paz. Collected by O. Salvin, vide Sclater; Catalogue of a collection of American birds (1862): 52). Sclater Collection. A female syntype is also in the collection.

guayabambae—Myrmotherula guayabambae Sharpe, 1900 4 Bull. Br. Orn. Cl., 11:2. Holotype, Adult female. Reg. no. 1899.6.10.207. Guayabamba (=Huayabamba), 4,000′, north Peru, 26 Aug. 1894. Collected by O. T. Baron. Salvin-Godman Collection.
guerrerensis—Aimophila acuminata guerrerensis van Rossem, 1938. 49 Bull. Br. Orn. Cl., 58: 128. Holotype, Adult male. Reg. no. 1899.2.1.3664. Acaguazotla, 3,500 ft., Guerrero, Mexico, 18 Oct. 1888. Collected by D. W. Smith. Salvin-Godman Collection.
guerrerensis—Automolus guerrerensis Salvin & Godman, 1891 . 3 Biol. CentrAmer., Aves, 2:157. Syntype, Adult male. Reg. no. 1899.6.1.109. 8,000', Omilteme, near Chilpancingo, Guerrero, Mexico, July 1888. Collected by Mrs H. H. Smith. There is a female syntype in the collection; the male specimen was fully described and could be regarded as the holotype.
guerrerensis—Certhia familiaris guerrerensis van Rossem, 1939 . 44A Proc. Biol. Soc. Washington 52:5. HOLOTYPE, Adult male. Reg. no. 1892.1.11.56. (not 1899 as stated in description). Omilteme (8,000', on label), Guerrero, Mexico, July 1888. Collected by D. W. Smith. Salvin-Godman Collection.
guerrerensis—Lepidocolaptes souleyetii guerrerensis van Rossem, 1939
guerrerensis—Pipilo torquatus guerrerensis van Rossem, 1938 49 Bull. Brit. Orn. Cl., 58: 131. HOLOTYPE, Adult, probably male. Reg. no. 1899.2.1.3849. Omilteme, 8,000', Guerrero, Mexico, July. Collected by D. W. Smith. Salvin-Godman Collection.
guerrerensis—Troglodytes brunneicollis guerrerensis van Rossem, 1938
guentheri—Myzomela guentheri Gadow, 1884
guillelmi—Alauda arvensis guillelmi Witherby, 1921

- guianensis—Dendrocincla longicauda guianensis Chubb, 1919. Bull. Br. Orn. Cl., 39:61. HOLOTYPE, Adult. Reg. no. 1889.5. 14.790. Bartica Grove, British Guiana, 20 Dec. 1880. Collected by H. Whitely. Salvin-Godman Collection. guianensis—Rhynchocyclus olivaceus guianensis McConnell, 1911 Bull. Brit. Orn. Cl., 27: 106. HOLOTYPE, Adult. Reg. no. 1922. 3.5.2638. (Abary River, on label), British Guiana, (Sept. 1906, on label). Collected by F. V. McConnell. McConnell Collection. guimarasensis—Cinnyris guimarasensis Steere, 1890 List of Birds and Mammals collected by the Steere Expedition to the Philippines, : 22. SYNTYPE, Adult male. Reg. no. 1896 6.6.462. Guimaras, Philippines, 24 Dec. Collected by J. B. Steere. Steere Collection. There is a second syntype in the collection. the Philippines: 19. SYNTYPE, Adult male. Reg. no. 1896.6.6.312. Guimaras, Philippines, 30 Dec. 1887. Collected by and purchased of J. B. Steere, (reprinted in *Ibis*, (6) 3 (1891): 313). guinanensis—Philydor erythrocercus guinanensis Chubb, 1919. Bull. Br. Orn. Cl., 39:60. HOLOTYPE, Adult male. Reg. no. 1922.3.5.2443. Ituribisi River, British Guiana, 1909. Collected by F. V. McConnell. McConnell Collection. guinea—Cisticola ruficeps guinea Lynes, 1930 1911.12.23.2528. Kintampo, Ashanti Province, Gold Coast, May 1902. Collected and presented by Boyd Alexander. guineae—Fraseria cinerascens guineae Bannerman, 1922 37 Bull. Brit. Orn. Cl., 42:69. HOLOTYPE, Adult male. Reg. no. 1910.5.6.516. Gunnal, Portuguese Guinea, West Africa, 30 May 1909. Collected by and purchased of W. J. Ansorge. guincensis—Cyanomitra obscura guineensis Bannerman, 1921 Bull. Brit. Orn. Cl., 41:135. HOLOTYPE, Adult male. Reg. no. 1911.12.18.291. Nanna Kru, Liberia, 1 Jan. 1911. Collected by W. P. Lowe. Presented by Capt. E. C. Hardy, RN. guineensis—Gerygone chrysogaster guineensis Mathews, 1928 . 36 Bull. Brit. Orn. Cl., 48:91. HOLOTYPE, Adult male. Reg. no. 1911.12.20.1124. Wakatimi, Mimika River, Dutch New Guinea. 10 May 1911. Collected by C. H. B. Grant. Presented by the subscribers to the British Ornithologists Union Expedition to Dutch New Guinea.
- guineensis—Parus leucomelas guineensis Shelley, 1900 . Birds of Africa 2: 222, 229. SYNTYPE, Adult. Reg. no. 1876.5.23. 470. Volta River, Gold Coast. Collected by Ussher. Sharpe Collection.

The other syntype is in the collection.

guirina—Nemosia guirina P. L. Sclater, 1856	
of a collection of American birds (1862): 87. gularis—Calornis gularis G. R. Gray, 1861	62
Proc. zool. Soc. Lond., 1861:431. HOLOTYPE, Adult female (vide Sharpe, Cat. Bds. Brit. Mus., 13 (1890):142). Reg. no. 1873.5.12.1901. Mysol, 1860. Collected by A. R. Wallace Wallace Collection.	
gularis—Campylorhynchus gularis P. L. Sclater, 1860 Proc. zool. Soc., Lond., 1860 : 462. HOLOTYPE, Adult. Reg. no. 1886.9.15.18. Mexico. Collected by Floresi (vide Hellmayr, Catalogue of Birds of the Americas, 7 (1934) : 151). Sclater Collection; obtained by Sclater from J. Gould.	,
gularis—Cichlopsis gularis Salvin & Godman, 1882 Ibis, (4) 6:76. SYNTYPE, Adult male. Reg. no. 1885.3.2.541. Merume Mountains, British Guiana, 22 June 1881. Collected by H. Whitely. Salvin-Godman Collection. The two female syntypes are also in the collection.	31
gularis—Ianthocincla gularis McClelland, 1840	2R
in Horsfield, <i>Proc. zool. Soc. Lond.</i> , 1839 (publ. 1840): 159. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4366. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.	
gularis—Lanius gularis Raffles, 1822	20
Trans. Linn. Soc. Lond., 13 (1821, publ. 1822): 304. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.2137. Sumatra (restricted to Benkulen, west Sumatra, by Kinnear and Robinson,	
Bull. Br. Orn. Cl., 47 (1927): 130). Collected by Sir Stamford Raffles and presented by the Honourable East India Company Museum.	
gularis—Linaria gularis Smith, 1836	59
Report of the Expedition for exploring Central Africa, 1834: 49. SYNTYPE, Adult (relaxed mount). Reg. no. 1845.7.6.141. South Africa. Collected by and purchased of A. Smith.	
There are two other syntypes in the collection, and others are at Philadelphia, vide Stone, Proc. Acad. Nat. Sci., Philadelphia, 1889: 562.	
gularis—Nicator gularis Finsch & Hartlaub, 1870	21
Die Vögel Ost-Afrikas,: 360. SYNTYPE, Adult. Reg. no. 1860. 12.31.194. Shupanga, Zambesi River (east of the mouth of Shiré River, Portuguese East Africa, fide C. H. B. Grant, MS). Col-	

lected by John Kirk of the Livingstone Expedition and received

from Lord Russell of the Foreign Office.

The authors stated Shupanga to be the type locality. The specimen from Tette on the Rivi River, listed as type by Gadow, *Cat. Bds. Brit. Mus.*, 8 (1883): 167 is also in the collection.

See also Gould, *Birds of Asia*, vol. 3, pt. 6 (1854): pl. 77. Gray used Horsfield's MS name.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 79. There is another syntype, in poor condition, in the collection.

gularis—Yuhina gularis Hodgson, 1836 32B Asiat. Res., 19: 166. Syntype, Adult (relaxed mount). Reg. no. 1843.1.13.166 Nepal. Collected and presented by B. H. Hodgson.

There is a second syntype in the collection.

- gulmergi—Hemichelidon sibirica gulmergi E. C. S. Baker, 1923 . 37 Bull. Brit. Orn. Cl., 43: 155. HOLOTYPE, Adult female. Reg. no. 1894.8.13.13. Gulmerg, Kashmir, 1 July 1893. Collected and presented by Dr. J. E. T. Aithison.

This specimen is labelled as the type. There are several other syntypes in the collection.

gurneyi—Ploceus gurneyi Shelley, 1887 61

Ibis, (5) 5:17, pl. 1. HOLOTYPE, Adult. Reg. no. 1889.7.25.11.

Caconda, Angola, July 1878. Collected by J. Anchieta. Shelley

Collection, the specimen having been received from Barboza du Bocage.

gurneyi—Promerops gurneyi J. Verreaux, 1871
gurue—Alethe anomala gurue Vincent, 1933
gustavi—Anthus gustavi Swinhoe, 1863
guttata—Aethomyias guttata Sharpe, 1882
guttata—Alauda guttata Brooks, 1873
guttata—Chlamydera guttata Gould, 1862
guttaticollis—Proparus guttaticollis Rickett & La Touche, 1897 32B Bull. Br. Orn. Cl., 6:50. SYNTYPE, Adult. Reg. no. 1905.12. 24.686. Kuatun, Fohkien Province, south-east China, Oct. 1896. Collected by C. B. Rickett and J. D. La Touche, and presented by the former. The second syntype is also in the collection.
guttaticollis—Turdinulus guttaticollis Ogilvie-Grant, 1895 32B Ibis, (7) 1:432. Syntype, Adult. Reg. no. 1895.7.14.649. Miri Hills, north of Sadiya, northern Assam. Collected by W. Robert. H. H. Godwin-Austen Collection. The three other syntypes are also in the collection.
guttatus—Dryoscopus guttatus Hartlaub, 1865 23 Proc. zool. Soc. Lond., 1865: 86. HOLOTYPE, Adult. Reg. no. 1873.12.10.165. Benguela, Angola. Collected by and purchased of J. J. Monteiro.

habessinicus	227
guttatus—Enicurus guttatus Gould, 1865	
guttatus—Salpinetes guttatus Salvin & Godman, 1891	j
guttuligera—Thripophaga guttuligera P. L. Sclater, 1864 Proc. zool. Soc. Lond., 1864: 167. HOLOTYPE, Adult. Reg. no 1889.5.20.306. Bogota, Colombia. Collected for Verreaux Sclater Collection.	٠.
gutturalis—Conopophaga gutturalis P. L. Sclater, 1868	5

- gutturalis—Conopophaga gutturalis P. L. Sclater, 1868 5

 Proc. zool. Soc. Lond., 1868: 574. HOLOTYPE, Adult (female, vide Sclater, Cat. Bds. Brit. Mus., 15 (1890): 332). Reg. no. 1889.9. 20.665. Bogota, Colombia. Sclater Collection.
 - tturalis—Falcunculus gutturalis Vigors and Horsfield, 1827. . . . 39
 Trans. Linn. Soc. Lond., 15: 212. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1863.7.7.17. "Kent's Group", Australia. Dec. 1803. Collected by R. Brown. Presented by the Linnean Society. The upper mandible is damaged.
- gutturalis—Myrmotherula gutturalis P. L. Sclater, 1881 . . . 4

 Ibis, (4) 5:269. Syntype, Adult male. Reg. no. 1889.9.20.254.

 Bartica Grove, British Guiana, 13 Jan. 1880. Collected by H.

 Whitely. Sclater Collection.

 There are five other syntypes in the collection.
- gutturalis—Phoenicothraupis gutturalis P. L. Sclater, 1854 . . . 51 Hnn. Mag. Nat. Hist. (2) 13:25. HOLOTYPE, Adult male. Reg. no. 1853.5.2.7. Colombia? Purchased of Parzudaki.
- gymnogenys—Crateropus gymnogenys Hartlaub, 1865 32B Proc. zool. Soc. Lond., 1865:86. HOLOTYPE, Adult. Reg. no. 1873.12.10.180. Benguela, Angola. Collected and presented by J. J. Monteiro.
- habeli—Camarhynchus habeli P. L. Sclater and Salvin, 1870. . . . 53

 Proc. zool. Soc. Lond., 1870: 325, fig. 4. SYNTYPE, Adult male.

 Reg. no. 1885.12.14.546. Abingdon Island, Galapagos Archipelago, (14 Dec. 1868, on label). Collected by A. Habel. Salvin-Godman Collection.

There are two other syntypes in the collection.

- 1869.10.16.80. (Senafó, on label), Abyssinia. Collected by W. T. Blanford. Presented by the Government of Bombay. Seebohm, *Cat. Bds. Brit. Mus.*, 5 (1873): 63, listed the female syntype, which is also in the collection, as the type.
- hachisuka—Cyrtostomus frenatus hachisuka Mathews, 1926. . . 46

 Bull Brit. Orn. Cl., 47: 67. HOLOTYPE, Adult male. Reg. no. 1911.1.14.9. Obi Island, 27 May 1902. Collected by and received in exchange from G. M. Mathews.
- hadii—Geokichla piaggiae hadii Macdonald, 1940 . . . 31

 Bull. Br. Orn. Cl., 60:98. Holotype, Adult male. Reg. no. 1939.10.1.173. Emogadung, 7,200', Dongotona Mountains, southeast Sudan, 9 April 1939. Collected by J. D. Macdonald for the British Museum (Natural History).
- haemalea—Pyranga haemalea Salvin and Godman, 1883 . . . 51

 Ibis, (5) 1:205. SYNTYPE, Adult male. Reg. no. 1885.6.7.421.

 Roraima, British Guiana, 4 Jan. 1882. Collected by H. Whitely, Salvin-Godman Collection.

 The female syntype is also in the collection.
- haematonota—Formicivora haematonota P. L. Sclater, 1857.

 Proc. zool. Soc. Lond., 1857: 48. HOLOTYPE, (Juvenile male, vide Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 146). Reg. no. 1855.12.17.43. Chamicuros, east Peru. Collected by J. Hauxwell. Gould Collection.
- haematopygia—Montifringilla haematopygia Gould, 1853 . . . 59

 Proc. zool. soc. Lond., 1851 (publ. 1853): 115. SYNTYPE, Adult (relaxed mount). Reg. no. 1851.7.2.20. Tibet. Collected and presented by Lord Gifford.

 There is another syntype in the collection.
- haematostictum—Dicaeum haematostictum Sharpe, 1876 . . . 45
 Nature, 14: 298. SYNTYPE, Adult male. Reg. no. 1876.8.28.46.
 Guimaras, Philippines. Collected by and purchased of J. B. Steere.
 as "one of the types".
- haesitata—Drymoeca haesitata P. L. Sclater and Hartlaub, 1881. 35
 Proc. zool. Soc. Lond., 1881: 166. Holotype, Male. Reg. no. 1881.3.21.16. Socotra (23 March 1880, on label). Collected by Prof. Balfour. Presented by the British Association.
- hainana—Siphia hainana Ogilvie-Grant, 1900

 Bull. Brit. Orn. Cl., 10: 36. SYNTYPE, Adult male. Reg. no. 1900.1.2.84. Five Finger Mountains, Hainan, (7 May 1899, on label). Collected by J. Whitehead. Purchased of E. Gerrard.

 This is the described male. The described female syntype is also in the collection.

- hainanum—Dicaeum cruentatum hainanum Hachisuka, 1926. . 45

 Bull. Brit. Orn. Cl., 47: 69. HOLOTYPE, Adult male. Reg. no. 1898.10.20.27. Hainan, China, May 1898. Collected by R. Swinhoe. Seebohm Collection.
- hainanus—Tephrodornis hainanus Ogilvie-Grant, 1910 20

 Proc. zool. Soc. Lond., 1910; Abstract no. 81:18 (after p. 588).

 SYNTYPE, Adult male. Reg. no. 1909.8.30.29. Seven-Fingers

 Mountains, central Hainan, 17 Nov. 1906. Collected and presented
 by R. Douglas.

 The female syntype is also in the collection.
- hallae—Acrocephalus boeticatus hallae C. M. N. White, 1960 . . . 35

 Bull. Brit. Orn. Cl., 80: 21. HOLOTYPE, Adult male. Reg. no. 1950.50.564. Brandberg Mountains, 2,500′, South-west Africa, 11 April 1950. Collected by J. D. Macdonald. Presented by the British Museum South-West Africa Expedition.
- hallae—Anthus similis hallae C. M. N. White, 1957 19
 Bull. Br. Orn. Cl., 77: 30. Holotype, Adult male. Reg. no.
 1923.8.7.2494. Lake Karange, Ankole, Uganda, 13 Dec. 1910.
 Collected by T. V. Fox and presented by Col. S. R. Clarke.
- hallae—Cisticola fulvicapilla hallae, Benson, 1955 35

 Bull. Brit. Orn. Cl., 75: 105. HOLOTYPE, Adult female. Reg. no.
 1932.5.5.84. Tsotoroghe Pan, North-eastern Bechuanaland, 26

 June 1930. Collected for the Verney-Lang Kalahari Expedition.
 Presented by A. S. Verney.
- hallae—Criniger ochraceus hallae Deignan, 1956. 21
 Smiths. Misc. Coll., 134 (2): 6. HOLOTYPE, Adult male. Reg. no. 1928.6.26.834. 11°18′ N., 106°07′ E., Tay Ninh Province, Cochin-China, 15 Jan. 1928. Collected by J. Delacour and W. P. Lowe on the Fourth Franco-British Expedition to Indo-China.
- hamadryas—Rhipidura hamadryas P. L. Sclater, 1883 37A Proc. zool. Soc. Lond., 1883 : 54. HOLOTYPE, Adult female. Reg. no. 1883.5.30.86. Larat Island, Tenimber Group (11 August 1882, on label). Collected by H. O. Forbes. Presented by the Timor-Laut Committee of the British Association.

hamertoni—Erythropygia hamertoni Ogilvie-Grant, 1906 31 Bull. Br. Orn. Cl., 19: 24. SYNTYPE, Adult male. Reg. no. 1908.5. 28.62. Beira, Somaliland, 14 May 1903 ("120 miles north-west of Obbia" on label). Collected by and purchased of A. E. Hamerton.
hanadori—Dicaeum dorsale hanadori Hachisnka, 1941
hanapepe—Hemignathus hanapepe Wilson, 1889
hancii—Oreocincla hancii Swinhoe, 1863
haplonota—Grallaria haplonota P. L. Sclater, 1877 4 Ibis, (4) 1:442. Syntype, Adult. Reg. no. 1889.9.20.611. Venezuela, 1873. Collected by Spence. Sclater Collection. There is another syntype in the collection. The selected specimen is the one upon which the author's diagnosis was based, and as such could be regarded as the holotype.
hardwickii—Garrulux leucolophus hardwickii Ticehurst, 1926 32B Bull. Br. Orn. Cl., 46:113. HOLOTYPE, Adult male. Reg. no. 1895.7.14.2301. Naga Hills, Assam, 29 March 1876. Godwin- Austen Collection.
hardyi—Apalis hardyi Bannerman, 1923
hardyi—Sylviella hardyi Bannerman, 1911
hariabica—Sitta hariabica Whistler, 1944

haringtoni—Acrocephalus agricola haringtoni Witherby, 1920. Bull. Brit. Orn. Cl., 41:26. HOLOTYPE, Adult male. Reg. no. 1924.12.20.8. Buttakundi, Kaghan, North-west India, 12 July 1914. Collected by Col. H. H. Harington. Presented by H. F. Witherby. haringtoni—Anthus trivialis haringtoni Witherby, 1917. 19 Bull. Br. Orn. Cl., 37:44. HOLOTYPE, Adult male. Reg. no. 1924.12.19.2. Gittidas, 11,000', Kaghan Valley, North-West Frontier, India, 2 July 1914. Collected by H. H. Harington and received in exchange with H. F. Witherby. haringtoni—Garrulus haringtoni Rippon, 1905 Bull. Brit. Orn. Cl., 15:97. HOLOTYPE, Adult. Reg. no. 1905.9. 10.2. Mount Victoria, 5,200', south Chin Hills, 25 April 1904. Collected and presented by Col. G. Rippon. haringtoni-Pomatorhinus haringtoni E. C. S. Baker, 1914 Bull. Br. Orn. Cl., 33: 124. HOLOTYPE, Adult. Reg. no. 1886.10. 1.3688. Mongphoo (on label), near Darjeeling, India, Feb. 1873. Hume Collection. harrisoni—Pyrrhulauda harrisoni Ogilvie-Grant, 1900 Bull, Br. Orn. Cl., 11:30. SYNTYPE, Adult male. Reg. no. 1901. 3.15.18. 1.800' above the south end of Lake Rudolf, 21 April 1900. Collected and presented by J. J. Harrison. The female syntype is in the collection. harterti—Acanthis cannabina harterti Bannerman, 1913 59 Bull. Brit. Orn. Cl., 33:39. HOLOTYPE. Adult male. Reg. no. 1913.10.22.27. Haria, (1,100', on label), Lanzarote, Eastern Canary Islands, 25 May 1913. Collected by and purchased of D. A. Bannerman. harterti—Alauda arvensis harterti Whitaker, 1904 Bull. Br. Orn. Cl., 15:19. HOLOTYPE, Adult male. Reg. no. 1939.12.14.20. El Oubira, north Tunisia, May 1897. Presented by J. Whitaker. harterti—Amydrus fulgidus harterti Neumann, 1903 Orn. Mber., 11:183. HOLOTYPE, Adult male. Reg. no. 1873.5. 24.21. Fort of Denera, Fantee, Gold Coast. Collected by St. T. D. Aubinn for Governor Ussher. Sharpe Collection. harterti—Cisticola cisticola harterti Witherby, 1920 Bull. Brit. Orn. Cl., 40:120. HOLOTYPE, Adult male. Reg. no. 1924.12.19.3. Platea, Greece, 9 Dec. 1905. Collected by H. Lynes. Received in exchange from H. F. Witherby. harterti—Diaphorophyia ansorgei harterti Bates, 1925. 38 Bull. Brit. Orn. Cl., 46: 105. HOLOTYPE, Adult female. Reg. no. 1926.8.8.8. Bitye, 2,000', Ja River, South Cameroon, 8 April 1925.

Collected by and purchased of G. L. Bates.

in the collection.

- harterti—Malacocincla sepiaria harterti Chasen & Kloss, 1929 . 32B J. Orn., 1929 (2): 116. Holotype, Adult male. Reg. no. 1947.60. 49. Bettotan, near Sandakan, North Borneo, 25 July 1927. Collected by F. N. Chasen and C. B. Kloss and presented by the Raffles Museum, Singapore.
- harterti—Pnoepyga pusilla harterti Robinson & Kloss, 1918 . . . 32B J. F. M. S. Mus., 8:205. Holotype, Adult male. Reg. no. 1927.4.18.1014. Gunong ljau, 4,700', Larut Range, Perak, Federated Malay States, 16 Aug. 1909. Collected by C. B. Kloss and presented by H. C. Robinson & Kloss.

Renamed R. rufifrons granti by Hartert, 1918 on the grounds of its preoccupation by R. flabellifera harterti Mathews, 1912.

- hartlaubii—Onychognathus hartlaubii Hartlaub, 1858 62

 Proc. zool. Soc. Lond., 1858 : 291. SYNTYPE, Adult male (relaxed mount). Reg. no. 1851.11.27.12. Fernando Po. Collected by L. Fraser. Purchased of H. Cuming.

 Hartlaub used a G. R. Gray MS. name. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890) : 166. There is a second syntype
- hartleyi—Stachyris nigriceps hartleyi Chasen, 1935 32B Bull. Raffles Mus., 10:43. HOLOTYPE, Adult male. Reg. no. 1935.12.27.2. Mount Poi, 5,000', west Sarawak, Borneo, 20 Oct. 1923. Collected by E. Mjöberg and presented by the Raffles Museum, Singapore.
- hauxwelli—Formicivora hauxwelli P. L. Sclater, 1857 4

 Proc. zool. Soc. Lond., 1857: 131, pl. 126. SYNTYPE, Adult male.

 Reg. no. 1855.12.17.56. Chamicuros, east Peru. Collected by
 J. Hauxwell and purchased of J. Gould.

 The other syntype is in the collection.
- hauxwelli—Furnarius leocopus hauxwelli Chubb, 1918 3

 Bull. Br. Orn. Cl., 38:87. HOLOTYPE, Adult female. Reg. no. 1889.5.20.28. Pebas, Rio Marañon, east Peru, 29 April 1867. Collected by J. Hauxwell. P. L. Sclater Collection.

hauxwelli—Icterus hauxwelli P. L. Sclater, 1885
hauxwelli—Pipromorpha eleaginea hauxwelli Chubb, 1919 9 Ann. Mag. Nat. Hist., (9) 4:302. HOLOTYPE, Adult male. Reg. no. 1888.1.1.538. Pebas, eastern Peru, 18 July 1866. Collected by J. Hauxwell. Salvin-Godman Collection.
hawkeri—Cinnyris mariquensis hawkeri Neumann, 1899 46 Orn. Mber., 7: 24. Holotype, Adult male. Reg. no. 1898.6.13.61. Jifa Medir, Somaliland. Collected and presented by R. Hawker.
hawkeri—Granatina hawkeri Lort Phillips, 1898 60 Bull. Brit. Orn. Cl., 8: 23. Syntype, Adult (male, on label). Reg. no. 1895.7.12.74. Somaliland. (Bari, Shebeli River, on label.) Collected by and purchased of E. Lort Phillips. The two other syntypes are also in the collection.
hawkeri—Saxicola hawkeri Ogilvie-Grant, 1908
hayii—Mirafra hayii Jerdon, 1844
haynesi—Crateropus haynesi Sharpe, 1871
hebridium—Prunella modularis hebridium R. Meinertzhagen, 1934. 30 Ibis, (13) 4:57. HOLOTYPE, Adult male. Reg. no. 1965.M.50. South Uist, Outer Hebrides, 17 Nov. 1920. Collected by R. Meinertzhagen. Meinertzhagen Collection.
helenae—Cyanomyias helenae Steere, 1890
helenae—Cyanomyias helenae Steere, 1890
helenae—Laniarius helenae Kelsall, 1913

helenae—Neopipo helenae McConnell, 1911 8 Bull. Brit. Orn. Cl., 27: 105. HOLOTYPE, Adult. Reg. no. 1922.3. 5.3093. Iturbisce, British Guiana, Oct. 1908. Collected and presented by F. V. McConnell.
helenae—Serpophaga helenae Chubb, 1919
helenorae—Eremomela helenorea Boyd Alexander, 1899 35 Bull. Brit. Orn. Cl., 8:48. Holotype, Adult (female, on label). Reg. no. 1911.12.23.2198. Mesanangue, Zambesi River (25 August 1898, on label). Collected by W. Boyd Alexander. Boyd Alexander Bequest.
helenorae—Poliolais helenorae Boyd Alexander, 1903 35 Bull. Brit. Orn. Cl., 13:36. HOLOTYPE, Adult. Reg. no. 1911.12. 23.2215. Bakaki, Fernando Po, 20 Nov. 1902. Collected by Boyd Alexander. Boyd Alexander Bequest.
helianthea—Muscicapa helianthea Wallace, 1865
hellmayri—Sisopygis hellmayri Chubb, 1907 Bull. Brit. Orn. Cl., 19:63. Holotype, Adult male (Immature, vide Hellmayr and Conover, Catalogue of Birds of the Americas, 5 (1927):96). Reg. no. 1902.3.13.649. Tapacari, (10.000', Dept. Cochabamba, 66° W, 17.5° S, on label). Bolivia, 23 March 1901. Collected by P. O. Simons. Simons Collection.
hemileuca—Hypocnemis hemileuca P. L. Sclater & Salvin, 1866. Proc. zool. Soc. Lond., 1866: 186. HOLOTYPE, Adult male. Reg. no. 1889.9.20.539. Lower Ucayali River, east Peru. Collected by E. Bartlett. Sclater Collection.
hemileucus—Passer hemileucus Ogilvie-Grant and Forbes, 1900 . 69 Bull. Liverpool Mus., 2:3. SYNTYPE, Adult male. Reg. no. 1899. 8.11.131. Abd-el-Kuri Island, 1,400′, off Sokotra, 5 Dec. 1898. Collected by W. R. Ogilvie-Grant and H. O. Forbes. Presented by the Royal Society and the British Association. The female syntype is also in the collection.
hemimelaena—Myrmeciza hemimelaena P. L. Sclater, 1857

- label). Collected by H. O. Forbes. Presented by the Timor-Laut Committee of the British Association. This specimen was labelled as the type. The syntype from Loetoe Island is also in the collection. lected by and purchased of G. Henderson. From the Yarkand Expedition. hendersoni—Saxicola hendersoni Hume, 1871 Ibis, (3) 1:408. SYNTYPE, Adult male (only 3 tail feathers). Reg. no. 1873.6.30.4. Koshtak, Yarkand, Turkestan, 10 Sept. 1870. Collected by G. Henderson on the Yarkand Expedition, and purchased of E. Gerrard. The second syntype is also in the collection. henrici—Mirafra africana henrici Bates, 1930 Bull. Br. Orn. Cl., 51:47. HOLOTYPE, Adult male. Reg. no. 1930.12.3.2. Bossu, 2,000', near foot of Mount Nimba, French Guinea, 23 May 1930. Collected by and purchased of G. L. Bates. herberti-Corythocichla brevicaudata herberti E. C. S. Baker, 1917. 32B Bull. Br. Orn. Cl., 38:10. HOLOTYPE, Adult male. Reg. no. 1916.12.27.867. Tung Song, Siam, 16 Sept. 1915. Collected by C. Chungat for E. G. Herbert and presented by the latter.
- herberti—Cryptolopha herberti Boyd Alexander, 1903. Bull. Brit. Orn. Cl., 13:35. HOLOTYPE, Adult female. Reg. no. 1911.12.23.1570. Near Bakaki, Fernando Po, 20 Nov. 1902. Collected by Boyd Alexander. Boyd Alexander Bequest.
- 1920.10.27.1. Ban Na Sao, Camp 42, 40 miles north-east of Pak Hin Bun, Laos, 24 Feb. 1920. Collected and presented by E. J. Herbert.
- herberti-Prinia inornata herberti E. C. S. Baker, 1918. Bull. Brit. Orn. Cl., 38:39. SYNTYPE, Adult male. Reg. no. 1916.12.27.974. Bangkok, Siam, 5 July 1915. Collected and presented by E. G. Herbert. This is the described male, the female syntype is also in the collection.
- herberti—Sycobrotus herberti Boyd Alexander, 1907 . . 61 Bull. Br. Orn. Cl., 19:88. HOLOTYPE, Adult female. Reg. no.

1911.12.23.3641. Pompari, Welle River, West Africa, 27 March 1906. Collected and bequeathed by Boyd Alexander. Alexander-Gosling Expedition.

- herioti—Cvornis herioti Wardlaw-Ramsav, 1886 Ibis, (5) 4:159. HOLOTYPE, Adult female. Reg. no. 1887.12.30. 1295. Manilla, Philippines. Collected by F. Maitland-Heriot. Tweeddale Collection.
- hermonensis—Calandrella hermonensis Tristram, 1864 Proc. zool. Soc. Lond., 1864: 434. SYNTYPE, Adult male. Reg. no. 1879.4.5.600. Foot of Mount Hermon, Palestine, 3 June 1864. Collected by H. B. Tristram. Salvin-Godman Collection.

There is another syntype in the collection. Tristram, Catalogue of a Collection of Birds belonging to H. B. Tristram, (1889): 245, stated the types to be in his possession, but according to Mr R. Wagstaffe of the Liverpool Museum (in litt. 1953), no specimens in the Tristram Collection are labelled as such.

- heuglini—Argya heuglini Sharpe, 1883 . 32B Cat. Bds. Brit. Mus., 7: 391. SYNTYPE, Adult. Reg. no. 1881.5.26.1. Mombasa. Collected and presented by the Rev. A. Buxton. The syntype from Zanzibar, also in the collection, is the holotype of Argva saturata Sharpe, 1895, q.v.
- hibernans—Pratincola torquata hibernans Hartert, 1910. J. Orn., 58: 173. HOLOTYPE, Adult male. Reg. no. 1936.10.15.9. Tring, Hertfordshire, England, 14 Nov. 1898. Collected by E. Hartert and presented by the American Museum of Natural History. New York, from the Rothschild Collection.
- hibernicus—Cinclus cinclus hibernicus Hartert, 1910 Die Vögel der palaarktischen Fauna, 1:790. HOLOTYPE, Adult male. Reg. no. 1936.10.15.11. Near Mallow, County Cork, Ireland, 25 Aug. 1896. Collected for the Brazenor brothers, naturalists of Brighton and presented from the Rothschild Collection by the American Museum of Natural History, New York.
- hibernicus—Garrulus glandarius hibernicus Witherby and Hartert, 15.1. County Wexford, Ireland, Nov. 1910. Collected by W. J. Williams. Presented by the American Museum of Natural History, from the Rothschild Collection.
- hibernicus—Parus hibernicus Ogilvie-Grant, 1910. Bull. Brit. Orn. Cl., 27:37. HOLOTYPE, Adult male. Reg. no. 1910.12.3.1. Boyle, Co. Sligo, Ireland, 3 Dec. 1910. Collected and presented by Collingwood-Ingram.

- hibernicus—Prunella modularis hibernicus R. Meinertzhagen, 1934. 30 Ibis. (13) 4:57. HOLOTYPE, Adult female. Reg. no. 1965.M.37. Curragh, Ireland, 6 Jan. 1907. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- hijazensis—Ammomanes deserti hijazensis Bates, 1935 Bull. Br. Orn. Cl., 56:8. HOLOTYPE, Adult male. Reg. no. 1934. 8.10.3. Hadda, Wadi Fatima, Arabia, 14 Sept. 1934. Collected and presented by H. Philby.
- hildebrandi—Hemixus hildebrandi Hume, 1874 Stray Feathers, 2:508. HOLOTYPE, Adult male. Reg. no. 1886. 9.1.1013. Banks of Younzaleen River, 3,000', Salween District, Tenasserim, 15 Feb. 1874. Collected by W. Davison. Hume Collection.
- hildebrandti— Urobrachva hildebrandti Sharpe, 1890 Cat. Bds. Brit. Mus., 13:225. HOLOTYPE, Adult male. Reg. no. 1879.3.4.38. Lake Bomba, Mombasa, East Africa. July 1876. Collected by and purchased of J. M. Hildebrandt.
- hildegardae—Euprinodes hildegardae Sharpe, 1899 35 Bull. Brit. Orn. Cl., 10:28. HOLOTYPE, Adult male. Reg. no. 1900.2.6.1. Athi River, Masailand, Kenva. Collected and presented by S. L. Hinde.
- himalayanus—Fregilus himalayanus Gould, 1862. Proc. zool. Soc. Lond., 1862: 125. SYNTYPE. Reg. no. 1876.1.7.6. Himalayas (Kumaon (=Almora), on label), (Restricted to Kumaon by Meinertzhagen, *Ibis*, (12) 3 (1927): 372). Collected by Strachey. Purchased of J. Gould.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 3 (1877): 148. There is another syntype in the collection.

- himalayensis—Loxia himalayensis Blyth, 1844 59 J. Asiat. Soc. Bengal, 13:952. Although Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 440, listed several specimens as types these were from Hodgson. The description was by Blyth, using a Hodgson MS. name. The specimens listed by Sharpe have no type status and the two specimens collected by Hodgson in Nepal in 1843 and in the Museum of the Asiatic Society (vide Blyth, Catalogue of the birds in the Museum of the Asiatic Society, (1849, publ. 1852): 123) were probably the syntypes.
- himalayensis—Regulus himalayensis Bonaparte, 1856 . 35 C. r. Acad. Sci. Paris, 42:767. HOLOTYPE, Adult. Reg. no. 1881.5.1.1440. Himalayan Mountains, (Nepal, on label). Purchased of Gould.

See also Deignan, Bull. Brit. Orn. Cl., 74 (1954): 104, and 76 (1956): 106; and Vaurie, Bull. Brit. Orn. Cl., 75 (1955): 99–101.

himalayensis—Sitta himalayensis Jardine and Selby, 1835		
Bull. Brit. Orn. Cl., 11: 29. SYNTYPE, Adult male. Reg. no. 1901.5.7.2. Athi River, Kenya. Collected and presented by S. L. Hinde. The second syntype is in the collection. hindii—Cisticola hindii Sharpe, 1896	Illustrations of Ornithology, 3: pl. 144. SYNTYPE, Adult. Reg. no. 1886.6.24.269. Himalayas. (Collected by W. Murray, on label.) Purchased from the Jardine Collection.	
Bull. Brit. Orn. Cl., 6:7. SYNTYPE, Adult, (male, on label). Reg. no. 1896.9.11.30. Machakos, East Africa, 3 May 1896. Collected and presented by L. L. Hinde. There is another syntype in the collection. hirtensis—Troglodytes hirtensis Seebohm, 1884	Bull. Brit. Orn. Cl., 11:29. SYNTYPE, Adult male. Reg. no. 1901.5.7.2. Athi River, Kenya. Collected and presented by S. L. Hinde.	•
Zoologist, (3) 8:333, pl. Holotype, Adult female. Reg. no. 1898.10.2.200. St. Kilda Island, 7 June 1884. Collected by C. Dixon. Seebohm Bequest. hodgsoni—Abrornis? hodgsoni Moore, 1854	Bull. Brit. Orn. Cl., 6:7. SYNTYPE, Adult, (male, on label). Reg. no. 1896.9.11.30. Machakos, East Africa, 3 May 1896. Collected and presented by L. L. Hinde.	
in Horsfield and Moore, Catalogue of birds in the Museum of the Honourable East India Company, 1: 412. Syntype, Adult, (relaxed mount). Reg. no. 1880.1.1.3011. Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company Museum. Moore used a Hodgson MS. name. hodgsoni—Callene hodgsoni Jerdon, 1872	Zoologist, (3) 8:333, pl. HOLOTYPE, Adult female. Reg. no. 1898.10.2.200. St. Kilda Island, 7 June 1884. Collected by C.	
hodgsoni—Callene hodgsoni Jerdon, 1872	in Horsfield and Moore, Catalogue of birds in the Museum of the Honourable East India Company, 1:412. SYNTYPE, Adult, (relaxed mount). Reg. no. 1880.1.1.3011. Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company Museum.	? !
Proc. Asiat. Soc. Bengal, 1871: 209. SYNTYPE, Adult (male, in museum register). Reg. no. 1886.11.1.1716. Gulmurg, Kashmir (4 June 1871, in museum register). Collected by W. E. Brooks. Hume Collection. There is one other syntype in the collection. hodgsoni—Nectarinia hodgsoni J. E. Gray, 1846	hodgsoni—Callene hodgsoni Jerdon, 1872	
Catalogue of Mammalia and birds of Nepal and Thibet presented by B. H. Hodgson : 59. Gray was referring to N. hodgsonis of Jardine, Naturalists' Library 13, Ornithology 5, Sunbirds (1843) : 240. Since the original description is Jardine's, the specimens listed by Gadow Cat. Bds. Brit. Mus., 9 (1884) : 16 as Gray's types	Proc. Asiat. Soc. Bengal, 1871: 209. SYNTYPE, Adult (male, in museum register). Reg. no. 1886.11.1.1716. Gulmurg, Kashmir (4 June 1871, in museum register). Collected by W. E. Brooks. Hume Collection.	1
	Catalogue of Mammalia and birds of Nepal and Thibet presented by B. H. Hodgson : 59. Gray was referring to N. hodgsonis of Jardine, Naturalists' Library 13, Ornithology 5, Sunbirds (1843) : 240. Since the original description is Jardine's, the specimens listed by Gadow Cat. Bds. Brit. Mus., 9 (1884) : 16 as Gray's types	5 : :

hodgsoni—Nemura hodgsoni Moore, 1854 In Horsfield & Moore, Catalogue of birds in the museum of the Honourable East India Company, 1 (Nov. 1854): 300. (Also in Proc. zool. Soc. Lond., 1854 (publ. April 1855): 76, pl. 72). Syntype, Adult male. Reg. no. 1880.1.1.805. (Hodgson's pl. no. 949.) Nepal. Collected by B. H. Hodgson. With the female syntype (also in the collection), presented as the types by the Honourable East India Company Museum. hodgsoni—Ruticilla hodgsoni Moore, 1854. in Horsfield & Moore, Catalogue of Birds in the museum of the Honourable East India Company, 1:303. SYNTYPE, Adult male. Reg. no. 1880.1.1.1437. Nepal. Collected by B. H. Hodgson and presented by the Honourable East India Company Museum. There are several other syntypes in the collection. hokrae—Acrocephalus concinens hokrae Whistler, 1930. Bull. Brit. Orn. Cl., 50:71. HOLOTYPE, Adult male. Reg. no. 1930.6.16.1. Hokra Jheel, 5,000', Kashmir, 10 Aug. 1928. Collected and presented by H. Whistler. holerythrops—Trochalopterum holerythrops Rippon, 1904 Bull. Brit. Orn. Cl., 14:83. HOLOTYPE, Adult. Reg. no. 1905. 9.10.80. Mount Victoria, 6,800', South Chin Hills, Burma, 15 Mar. 1904. Collected and presented by G. Rippon. holerythrus—Lipaugus holerythrus P. L. Sclater and Salvin, 1860 . 7 Proc. zool. Soc. Lond., 1860: 300. SYNTYPE, Adult. Reg. no. 1888.1.20.624. (Choctum, on label), Vera Paz, Guatamala, Jan. 1860. Collected by O. Salvin. Salvin-Godman Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 357. hollandi—Hapalocercus hollandi P. L. Sclater, 1896 Ibis, (7) 2:317. HOLOTYPE, Adult female. Reg. no. 1897.11.14.63. Sta. Elana, (Entrerios), Argentine, 15 Jan. 1895. Collected by and purchased of A. H. Holland. (See also Hellmayr, Novit. Zool., 13 (1906) : 320.) holochlora—Chloropipo holochlora P. L. Sclater, 1888 . Cat. Bds. Brit. Mus., 14:287. HOLOTYPE, Adult. Reg. no. 1888.1.13.1220. Bogota, Colombia. Collected by Boucard. Sclater Collection. holomelas—Dryoscopus holomelas Jackson, 1906 Bull. Br. Orn. Cl., 16:90. HOLOTYPE, Adult male. Reg. no. 1906.6.3.1. Ruwenzori, Uganda, 21 Feb. 1902. Collected and presented by F. J. Jackson. holopolius—Graucalus holopolius Sharpe, 1888 Proc. zool. Soc. Lond., 1888: 184. HOLOTYPE, Adult female. Reg. no. 1888.2.7.31. Aola, Guadalcanar, Solomons, 21 April 1887. Collected by and purchased of C. M. Woodford.

holospodia—Taenioptera holospodia P. L. Sclater, 1887
holospodium—Parisoma holospodium Bates, 1908 37 Bull. Brit. Orn. Cl., 25: 27. SYNTYPE, Adult male. Reg. no. 1911.4.27.3. Bitye, 2,000′, Ja River, Southern Cameroon, 14 May 1908. Collected and presented by G. L. Bates. This is the described male syntype. The described female syntype is also in the collection.
holostictus—Automolus holostictus P. L. Sclater, 1875
holsti—Parus holsti Seebohm, 1894
holtii—Hypsipetes holtii Swinhoe, 1861
homochroa—Catamenia homochroa P. L. Sclater, 1859 95 Proc. zool. Soc. Lond., 1858 (publ. 1859): 552. Holotype, Adult male. Reg. no. 1885.2.10.408. Matos, Ecuador, (June 1858, on label). Collected by L. Fraser. Sclater Collection.
homochroa—Larvivora homochroa Hodgson, 1844 31 in J. E. Gray, Zoological Miscellany, 1844: 83 (nomen nudum). Two specimens in the collection were listed as types by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 27.
homochrous—Dendromanes homochrous P. L. Sclater, 1859
homochrous—Pachyrhamphus homochrous P. L. Sclater, 1859. 7 Proc. zool. Soc. Lond., 1859: 142. SYNTYPE, Adult male. Reg. no. 1888.1.13.1428. Pallatanga, Ecuador (Oct. 1859, on label). Collected by L. Fraser. Sclater Collection. The female syntype is also in the collection.
hopwoodi—Dicrurus leucophaeus hopwoodi E. C. S. Baker, 1918 . 64 Novit. Zool. 25: 294. HOLOTYPE, Adult. Reg. no. 1886.3.1.2012. Dacca, East Pakistan, Jan. 1871. Hume Collection.

horsfieldi—Oreocincla horsfieldi Bonaparte, 1857
horsfieldii—Myophonus horsfieldii Vigors, 1831
horsfieldii—Pomatorhinus horsfieldii Sykes, 1832 32B Proc. zool. Soc. Lond., 1832 : 89. Syntype, Adult (relaxed mount). Reg. no. 1880.1.1.1934. Dense woods of the Ghauts, India (in text; Mahableshwar, vide Salim Ali, The Birds of Travancore and Cochin, 1953 : 35). Collected by W. H. Sykes and presented by the Honourable East India Company Museum. There is a second syntype in the collection.
hortulorum—Turdus hortulorum P. L. Sclater, 1863 31 Ibis, (1) 5:196. SYNTYPE, Adult female. Reg. no. 1896.6.1.836. Amoy, south China, 2 April 1861. Collected by R. Swinhoe. Seebohm Bequest. No male syntype can be traced in the collection.
hosei—Pitta caerulea hosei E. C. S. Baker, 1918
hosii—Calyptomena hosii Sharpe, 1892
hosii—Dicaeum hosii Sharpe, 1897
hosii—Oriolus hosii Sharpe, 1892
Bull. Brit. Orn. Cl., 1: 4. SYNTYPE, Adult male. Reg. no. 1892.8.25. 6. Mount Dulit, 4,000', Sarawak, Borneo, June 1892. Collected by C. Hose. Purchased of E. Gerrard. The female syntype is also in the collection.
houghtoni—Sheppardia cyornithopsis houghtoni Bannerman, 1931 . 31
Bull. Br. Orn. Cl., 51:128. HOLOTYPE, Adult male. Reg. no. 1931.6.7.1. Sugarloaf Forest, near Freetown, Sierra Leone, (26 Oct. 1913, on label). Collected by and purchased of G. J. Houghton.

huambo—Cisticola natalensis huambo Lynes, 1930 35 Ibis, (12) 6, Supplement, 441, pl. 15. Holotype, Adult male. Reg. no. 1928.10.25.1. Lepe, 4,900', Benguela District, Angola, 18 Jan. 1927. Collected by Admiral H. Lynes and B. B. Osmaston. Presented by Admiral H. Lynes.
huancavelicae—Asthenes dorbignyi huancavelicae Morrison, 1938 . 3 Ibis, (14) 2:774. HOLOTYPE, Adult male. Reg. no. 1939.6.17.2. 11,000', Yauli, Huancavelica Dept., Peru, 31 Dec. 1937. Collected by A. Morrison. Morrison Collection.
huancavelicae—Upucerthia serrana huancavelicae Morrison, 1938 . 3 Ibis, (14) 2:774. Holotype, Adult male. Reg. no. 1939.6.17.3. 13.650', Lachocc, Huancavelica, central Peru, 10 Oct. 1937. Collected by A. Morrison. Morrison Collection.
hudsoni—Cnipolegus hudsoni P. L. Sclater, 1872
hudsoni—Synallaxis hudsoni P. L. Sclater, 1874
hufufae—Passer domesticus hufufae Ticehurst and Cheesman, 1924 61 Bull. Br. Orn. Cl., 45:19. HOLOTYPE, Adult male. Reg. no. 1924.11.30.20. Hufuf Town, Hasa Province, Central Arabia, 9 Dec. 1923. Collected by R. E. Cheesman, Cox-Cheesman Collection.
hufufae—Prinia gracilis hufufae Ticehurst and Cheesman, 1924 . 35 Bull. Brit. Orn. Cl., 44:19. Holotype, Adult male. Reg. no. 1924.11.30.59. Hufuf, Hasa Province, Central Arabia, 28 Nov. 1923. Collected by R. E. Cheesman. Cox-Cheesman Collection.
humei—Aegithina tiphia humei E. C. S. Baker, 1922 22 Fauna of British India (2nd ed.); Birds, 1:342. HOLOTYPE, Adult female. Reg. no. 1886.9.1.143. Raipur, Central Provinces, 12 May 1870. Collected by A. O. Hume. Hume Collection.
humei—Heterorhynchus humei Mandelli, 1873
humei—Lophophanes humei Brooks, 1873

1886.11.1.553. Sikkim, 10 Feb. 1873. Collected by L. Mandelli. Hume Collection.

There is another syntype in the collection.

humei—Reguloides humei Brooks, 1878
humeralis—Dessornis humeralis Smith, 1836
humeralis—Penthetriopsis humeralis Sharpe, 1901 61 Bull. Br. Orn. Cl., 11:57. HOLOTYPE, Adult. Reg. no. 1901.10.9.1. Mount Elgon, Kenya, (Nandi, Kenya, on label). Collected and presented by F. J. Jackson.
humeralis—Terenura humeralis P. L. Sclater & Salvin, 1880 4 Proc. zool. Soc. Lond., 1880: 159. SYNTYPE, Adult male. Reg. no. 1889.7.10.510. Sarayacu, east Ecuador. Collected by C. Buckley. Salvin-Godman Collection. There are two other syntypes in the collection.
humii—Coccothraustes humii Sharpe, 1886
humii—Microscelis psaroides humii Whistler & Kinnear, 1932 . 21 J. Bombay nat. Hist. Soc., 35: 754. HOLOTYPE, Adult male. Reg. no. 1886.9.1.964. Ceylon, 7 Feb. 1872 (1892 printed in error). Hume Collection.
humii—Molpastes humii Oates, 1889
humii—Pyrrhospiza humii Sharpe, 1888
humii—Sturnus humii Brooks, 1876

(1871): 410, (q.v.), which is preoccupied by S. nitens Brehm, Ibis, 1828: 282.
Since this is only a change of name the specimen listed as type by Sharpe, <i>Cat. Bds. Brit. Mus.</i> , 13 (1890) : 35, is not valid.
humii—Suthora humii Sharpe, 1883
Cat. Bds. Brit. Mus., 7: 487. SYNTYPE, Adult. Reg. no. 1859.3.4. 294. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
humilis—Ixulus humilis Hume, 1877
Stray Feathers, 5:106. HOLOTYPE, Adult male. Reg. no. 1886.10.1.6321. Moolyit, central Tenasserim hills, 16 Feb. 1877. Collected by W. Davison (on label). Hume Collection.
humilis—Podoces humilis Hume, 1871
humilis—Podoces humilis Hume, 1871
humphreysi—Orthotomus atrogularis humphreysi Chasen and Kloss,
1929
Bettotan, near Sandakan, North Borneo, 15 Aug. 1927. Collected by F. N. Chasen and C. B. Kloss. Presented by the Raffles
Museum, Singapore.
hunsteini—Donacicola hunsteini Finsch, 1886 60
Ibis, (5) 4: 1, pl. 1. SYNTYPE, Adult male. Reg. no. 1887.1.26.3. North New Ireland Island, Pacific Ocean, 6 June 1885. Collected by O. Finsch. Purchased of G. Schneider. The two other syntypes are also in the collection.
hunteri—Cinnyris hunteri Shelley, 1889
Proc. zool. Soc. Lond. 1889: 365 pl. 41. Syntype, Adult male.
Reg. no. 1889.3.21.142. Useri River, Kilimanjaro, Tanganyika, (July, vide Shelley, Birds of Africa, 2 (1900): 103). Collected and
presented by H. C. V. Hunter.
This specimen was labelled as the type. The female syntype is
also in the collection.
hunteri—Cisticola hunteri Shelley, 1889
Proc. zool. Soc. Lond., 1889: 364. HOLOTYPE, Adult male. Reg. no. 1889.3.21.185. Mt. Kilimanjaro (above forest belt), between
June and August 1888. Collected and presented by A. C. V. Hunter.
huriensis—Galerida theklae huriensis C. W. Benson, 1947
Bull. Br. Orn. Cl., 68:9. HOLOTYPE, Adult male. Reg. no.
1946.5.2057. Huri Hills, 3,000', between Mega and Marsabit,
5 miles south of the Abyssinia-Kenya border, 1 April 1942. Col-
lected and presented by C. W. Benson.

There is a second syntype in the collection. This name was based on Blyth's description (J. Asiat. Soc. Bengal, 18 (1849): 811) of the bird he listed as Accentor atrogularis Brandt.

Listed as the type by Sharpe, *Hist. Coll. Brit. Mus.*, 2 (1906): 394. Although Blyth refers to this as a new species he also relates it to *E. buchanani* Blyth, 1844, and it is not wholly clear as to his intentions. According to W. L. Sclater, (*Ibis*, (5) 1 (1892): 73) the type of *E. buchanani* is lost.

- hyacinthinus—Gyanocorax hyacinthinus Cabanis, 1848 71 in Schomburgk, Reisen im Britisch Guiana, 3:683. SYNTYPE, Adult. Reg. no. 1840.7.3.42. Canuku Mountains, British Guiana. Collected by and purchased of Sir R. Schomburgk.

 There was another syntype, at present untraced, in the collection.
- hybrida—Pterythius pallidus hybrida Harington, 1913 32B Bull. Br. Orn. Cl., 33:82. Syntype, Adult. Reg. no. 1905.9.10. 399. Mount Victoria, South Chin Hills, 25 March 1904. Collected and presented by G. Rippon.

 This is the author's selected male. His selected female is also in

This is the author's selected male. His selected female is also in the collection.

- hypernephala—Cisticola hunteri hypernephala Elliott, 1947 35 Bull. Brit. Orn. Cl., 68: 10. Holotype, Adult male. Reg. no. 1947.71.1. Olosirwa Mountain, 10,950', Crater Highlands, Northern Tanganyika, 27 Jan. 1945. Collected and presented by H. I. F. Elliott.

- hyperythra—Rhipidura hyperythra G. R. Gray, 1858 37A Proc. zool. Soc. Lond., 1858 : 176. HOLOTYPE, Adult, ?male. Reg. no. 1873.5.12.455. Aru Islands, 1857. Collected by A. R. Wallace. Wallace Collection.
- hyperythrus—Thamnophilus hyperythrus P. L. Sclater, 1855 . . . 4

 Edinb. New philos. J., (n.s.) 1:235. Holotype, Adult female
 (relaxed mount). Reg. no. 1855.11.5.35. Chamicuros, Peruvian
 Amazon. Collected by J. Hauxwell and purchased of J. Gould.
 Sclater used Gould's MS. name.
- hyperythrus—Thryothorus hyperythrus Salvin & Godman, 1880 . 28 Biol. Centr.-Amer.; Aves, 1:91. Holotype, Adult. Reg. no. 1885.3.6.364. Paraiso Station, Panama Railway, Panama, 1867. Collected by A. Hughes. Salvin-Godman Collection.
- hypochloris—Stelgidillas hypochloris Jackson, 1906 . . . 21 Bull. Br. Orn. Cl., 19:20. HOLOTYPE, Adult male. Reg. no. 1907.6.19.4. Kibiran, Toro, Uganda, 7 Oct. 1905. Collected and presented by F. J. Jackson.

This is the author's selected male; the female syntype is also in the collection.

- hypochryseus—Vireo hypochryseus Sclater, 1862 5

 Proc. zool. Soc. Lond., 1862 : 369, pl. 46. HOLOTYPE, Adult. Reg. no. 1886.9.15.283. Mexico. Collected for F. Parzudaki. Sclater Collection.
- hypoglaucus—Oxyrhamphus hypoglaucus Salvin and Godman, 1883 10 Ibis, (5) 1:208. Syntype, Adult male. Reg. no. 1888.1.1.7. Roraima Mountains, British Guiana, 7 Dec. 1881. Collected by H. Whitely. Salvin-Godman Collection.

There are several other syntypes in the collection.

hypogrammica—Butalis hypogrammica Wallace, 1862 . . . 37 Ibis, (1) 4:350. SYNTYPE, Adult male. Reg. no. 1873.4.12.640. Ceram, Northern Moluccas, 1859. Collected by A. R. Wallace. Wallace Collection.

Wallace used a G. R. Gray MS. name. There are several other syntypes in the collection.

- hypogrammica—Pytelia hypogrammica, Sharpe, 1870 60

 Ibis, (2) 6:56. HOLOTYPE, Adult male. Reg. no. 1877.7.11.118.

 Fantee, Gold Coast. Sharpe Collection; obtained by him from the dealer Higgins.
- hypoleuca—Petroica hypoleuca G. R. Gray, 1859. 37

 Proc. zool. Soc. Lond., 1859: 155. HOLOTYPE, Adult male. Reg.
 no. 1859.4.8.26. Dorey, North-west New Guinea, 1858. Collected
 by A. R. Wallace. Purchased of Stevens.
- hypoleuca—Serpophaga hypoleuca P. L. Sclater and Salvin, 1866.

 Proc. zool. Soc. Lond., 1866: 188. SYNTYPE, Adult. Reg. no. 1888.1.13.354. Lower Ucayali River, East Peru, 1865. Collected by E. Bartlett. Sclater Collection.

 Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 104.

There are several other syntypes in the collection.

Labelled "Type" in an early hand, and listed as such by Mathews, *Austral Avian Record*, 1 (1913): 166. Another syntype is in the collection.

- hypomelaena—Ceromacra hypomelaena P. L. Sclater, 1890 . . . 4 Cat. Bds. Brit. Mus., 15: 268. Holotype, Adult male. Reg. no. 1889.7.10.574. Cosnipata, south-west Peru, 2 June 1871. Collected by H. Whitely. Salvin-Godman Collection.
- hypophaeus—Chlorospingus hypophaeus P. L. Sclater and Salvin, 1868 51 Proc. zool. Soc. Lond., 1868: 389. SYNTYPE, Adult female. Reg.

- no. 1885.6.8.41. Calovevora, Veragua, 1868. Collected by E. Arcé. Salvin-Godman Collection.

 There are several other syntypes in the collection.
- hypospodia—Elainea hypospodia P. L. Sclater, 1887 9

 Proc. zool. Soc. Lond., 1887 : 49. HOLOTYPE, Adult male. Reg.
 no. 1888.1.13.588. Near Valencia, Venezuela, 1875. Collected by A. Goering. Sclater Collection.
- hypospodia—Pinarochroa hypospodia Shelley, 1885 31 Proc. zool. Soc. Lond., 1885 : 226. pl. 13. HOLOTYPE, Adult female. Reg. no. 1885.6.14.18. Kilimanjaro, 14,000', 1884. Collected by H. H. Johnston and presented by the British Association.

Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 45. There are several other syntypes in the collection.

- hypospodius—Thryothorus hypospodius Salvin & Godman, 1880. 28

 Biol. Centr.-Amer.; Aves, 1:92. Holotype, Adult. Reg. no.
 1885.3.6.357. Vicinity of Bogota, Colombia. Collected by
 Trübner. Salvin & Godman Collection.
- hypostictus—Campylorhynchus hypostictus Gould, 1855 . . . 28

 Proc. zool. Soc. Lond., 1855 : 68. HOLOTYPE, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 6 (1881) : 190). Reg. no. 1855.11.5.15. Ucayali River, Peru. Collected by H. W. Bates (vide Sharpe, loc. cit.), and purchased of J. Gould.
- hypoxantha—Drymoica hypoxantha Sharpe, 1877 35
 in Layard, Birds of South Africa, 2nd Edn., : 260. SYNTYPE,
 Adult male. Reg. no. 1876.5.23.141. Eland's Post, Cape Province,
 June 1870. Collected by T. C. Atmore. Sharpe Collection.
 Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883) : 191.
 There are two female syntypes in the collection.
- hypoxantha—Neodrepanis hypoxantha Salomonsen, 1933 . . . 14
 Bull. Brit. Orn. Cl., 53:182. HOLOTYPE, Adult male. Reg. no.
 1882.2.27.20. East of Antanarivo, Madagascar, July 1881. Collected by and purchased of the Rev. W. D. Cowan.
 The bill is broken.

hypoxanthus—Andropadus hypoxanthus Sharpe, 1876. in E. L. Layard, Birds of South Africa: 205. SYNTYPE, Adult. Reg. no. 1860.12.31.75. Tette, Zambesi. Collected by J. Kirk and presented by the Foreign Office; supposedly from the Livingstone Expedition. There are two other syntypes in the collection. hypoxanthus—Attila hypoxanthus Salvin and Godman, 1891. Biol. Centr.-Amer., Aves, 2:135. HOLOTYPE, Adult. Reg. no. 1888.1.20.658. Vera Paz, Guatamala. Collected by O. Salvin. Salvin-Godman Collection hyrcanus—Erithacus hyrcanus Blanford, 1874 . . Ibis, (3) 4:79. SYNTYPE, Adult male. Reg. no. 1874.11.23.126. (Resht, Persia, on label.) Collected by Col. St. John and received in exchange with W. T. Blanford of the Indian Museum. hyrcanus—Garrulus hyrcanus W. T. Blanford, 1873 71 Ibis, (3) 3:225. SYNTYPE, Adult male. Reg. no. 1874.11.23.182. Mazendaran, North Persia. Collected by Major St. John from the collection of the Boundary Commission. Received in exchange from the Indian Museum, Calcutta. No type was designated. This specimen was collected with the one in the Indian Museum subsequently referred to as the type by W. L. Sclater, *Ibis*, (6) 4 (1892) : 73. ibadanensis—Malimbus ibadanensis Elgood, 1958 Ibis, 100:621. HOLOTYPE, Adult male. Reg. no. 1958.9.2. Ibadan, Nigeria, 28 Feb. 1958. Collected and presented by J. H. Elgood. iberiae—Muscicapa hypoleuca iberiae Witherby, 1928 . Ibis, (12) 4:591. HOLOTYPE, Adult male. Reg. no. 1929.1.15.1. San Ildefonso, Segovia, Central Spain, 30 May 1927. Collected and presented by H. F. Witherby. ibericus—Coloeus monedula ibericus Kleiner, 1939 Bull. Brit. Orn. Cl., 60:12. HOLOTYPE, Adult male. Reg. no. 1897.11.10.605. Mountains of Granada, Spain, April 1871. Collected by Howard Saunders. Purchased of E. Hargitt. ibericus—Phylloscopus collybita ibericus Ticehurst, 1936 35 Bull. Brit. Orn. Cl., 57:64. HOLOTYPE, Adult male. Reg. no. 1934.1.1.5045. Paul D'Argila, near Coimbra, Portugal, 23 May 1920. Collected by H. F. Witherby. Witherby Collection. ichangensis-Hodgsonius phoenicuroides ichangensis E. C. S. Baker, 31 Bull. Br. Orn. Cl., 43:18. HOLOTYPE, Adult male. Reg. no. 1914.7.16.551. Ichang, Upper Yangtse Valley, China, 5 Aug. 1895. Collected by F. W. Styan. Styan Collection.

icterica—Prinia icterica Strickland, 1894
icterioides—Coccothraustes icterioides Vigors, 1830 59 Proc. zool. Soc. Lond., 1830 : 8. SYNTYPE, Adult female (relaxed mount). Reg. no. 1855.12.19.254. Himalayas. Purchased of the Zoological Society of London, having been received by them, through J. Gould, from India. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888) : 45. The male syntype can no longer be traced.
icterphrys—Vireolanius icterphrys Bonaparte, 1854
ignicauda—Cinnyris ignicauda Hodgson, 1837
igniceps—Heteropelma igniceps P. L. Sclater, 1872 8 Proc. zool. Soc. Lond., 1871 (publ. 1872): 750. SYNTYPE, Adult. Reg. no. 1888.1.13.1386. Oyapok, Cayenne, 1871. Collected by E. Jelski (vide Brit. Mus. register). Scalter Collection, received by him from Mde. Verdey of Paris. The second syntype is in the collection.
ignicolle—Dicaeum ignicolle G. R. Gray, 1858
igniferum—Dicaeum igniferum Wallace, 1863
ignipectus—Myzanthe ignipectus Blyth, 1843

- Company appear to have been sent before the date of the description; and a Hodgson specimen on which the description was probably based (vide Blyth, Catalogue of birds in the Museum of the Asiatic Society, (1849): 227) is still in India.
- ignobilis—Automolus ignobilis P. L. Sclater & Salvin, 1879 3

 Proc. zool. Soc. Lond., 1879: 522. HOLOTYPE, Adult. Reg. no. 1889.5.20.324. Frontino, Antioquia, Columbia. Collected by T. K. Salmon. Sclater Collection.

See also Sclater and Salvin, *Proc. zool. Soc. Lond.*, 1870: 784. The specimen is imperfect.

There is a second syntype in the collection, and two more stated by Sclater to be in the Academy of Natural Sciences, Philadelphia.

- imatong—Cisticola ayresii imatong Cave, 1938 35

 Bull. Brit. Orn. Cl., 59:8. HOLOTYPE, Adult male. Reg. no.
 1938.8.5.1. Lamoling, 8,000', Imatong Mountains, Equatorial
 Province, Sudan, 29 April 1938. Collected and presented by F. O.
 Cave.
- imbricata—Zoothera imbricata Layard, 1854 31

 Ann. Mag. nat. Hist., (2) 13:212. HOLOTYPE, Adult (relaxed mount). Reg. no. 1854.1.13.48. Ceylon. Collected by Thwaites and purchased of H. Cuming.
- immaculata—Ceromacra cinerascens immaculata Chubb, 1918. . 4
 Bull. Br. Orn. Cl., 38:84. Syntype, Adult male. Reg. no. 1922.

3.5.2163. Supenaam River, British Guiana, March 1910. Collected by F. V. McConnell. McConnell Collection.

This is the author's selected male. His selected female and two other syntypes are also in the collection.

- immaculata—Mirafra immaculata Hume, 1873 . 17 Stray Feathers, 1:12. HOLOTYPE, Adult. Reg. no. 1887.7.1.2644. Deobund, 9,000', Mussoorie, Himalayas, June 1968. Hume Collection
- immaculata—Myrmeciza immaculata P. L. Sclater & Salvin, 1864 Proc. zool. Soc. Lond., 1864: 357. SYNTYPE, Adult male. Reg. no. 1889.7.10.637. Lion Hill, Panama. Collected by J. M'Leannan. Salvin-Godman Collection. There are two other syntypes in the collection.
- immaculatus—Accentor immaculatus Hodgson, 1845 30 Proc. zool. Soc. Lond., 1845: 34. ?SYNTYPE, Adult (relaxed mount). Reg. no. 1843.1.13.353. Nepal. Collected and presented by B. H. Hodgson.

There are several other specimens in the collection which may be

syntypes.

- immaculatus—Brachypodius immaculatus Sharpe, 1876 Ibis, (3) 6:39. SYNTYPE, Adult. Reg. no. 1873.5.12.2463. Sumatra, 1861. Collected by A. R. Wallace. Wallace Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 66. The immature syntype from Sibu Island is also in the collection.
- immodestus—Pericrocotus immodestus Hume, 1877 20 Stray Feathers, 5:177. SYNTYPE, Adult male. Reg. no. 1886. 4.1.859. Mergui, Tenasserim, 8 Nov. 1874. Collected by W. Davison. Hume Collection.

There are several other syntypes in the collection.

immundus—Lipaugus immundus P. L. Sclater and Salvin, 1873. Nomencl. Av. Neotrop., : 57, 159. SYNTYPE, Adult male. Reg. no. 1888.1.13.1552. Oyapoc, Cayenne, 1871. Collected by Verdey. Sclater Collection.

The second syntype is in the collection.

impetuani—Emberiza impetuani Smith, 1836 Report of an Expedition for the exploring of Central Africa, 1834: 48. SYNTYPE, Adult (relaxed mount). Reg. no. 1845.7.6.143. Between the Nu Gariep and tropic, (South Africa, on label). Collected by and purchased of A. Smith.

There is another syntype in the collection and others are in the Philadelphia Museum, vide, Stone, Proc. Acad. Nat. Sci. Philadel-

phia 1899: 56.

impiger—Euscarthus impiger P. L. Sclater and Salvin, 1868 Proc. zool. Soc. Lond., 1868: 171, pl. 13. SYNTYPE, Adult male.

- Reg. no. 1888.1.13.286. Caraccas, Venezuela. "Latter part of summer 1867." Collected by A. Goering. Sclater Collection.

 Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 84.

 The second syntype is in the collection.
- improbus—Tyrannicsus improbus P. L. Sclater and Salvin, 1871 . 9

 Proc. zool. Soc. Lond., 1870 (publ. 1871): 841, pl. 53. SYNTYPE,

 Adult (male, on label). Reg. no. 1888.1.13.488. South of Merida,

 Venezeula. Collected by A. Goering. Sclater collection.

 Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 133.

 The second syntype is also in the collection.
- incae—Pseudochloris aureiventris incae Chubb, 1919 59

 Bull. Brit. Orn. Cl., 39:71. SYNTYPE, Adult male. Reg. no.
 1899.1.20.54. Puente del Inca, 9,170′, Aconcagua Valley, Chilean
 Andes, 16 Dec. 1896. Collected by P. H. Gosse. Presented by
 E. A. Fitzgerald.

 The female syntype is in the collection.
- incana—Cisticola incana P. L. Sclater and Hartlaub, 1881 35

 Proc. zool. Soc. Lond., 1881: 166. SYNTYPE, Adult male. Reg.
 no. 1881.2.21.17. Socotra Island, 20 March 1880. Collected by
 1. B. Balfour. Presented by the British Association.

 There are two other syntypes in the collection.
- - See also, Hartlaub, Abhand. Nat. Ver. Bremen, 7 (1881): 91. There is a female syntype in the collection and other syntypes are at Bremen and Berlin.

incertus—Carpodacus incertus Ogilvie-Grant, 1906. Bull. Brit. Orn. Cl., 16: 122. HOLOTYPE, Adult female. Reg. no. 1907.12.12.14. Mount Morrison, 8,000', Central Formosa, Jan. 1906. Collected by and purchased from W. Goodfellow. Renamed C. formosanus by Ogilvie-Grant, Bull. Brit. Orn. Cl., 27 (1911): 51, on the grounds of its preoccupation by Fringilla incerta Risso, 1826.	
incii—Muscipeta incii Gould, 1852	
incognita—Sturnia incognita Hume, 1879	62
indica—Kittacincla macroura indica E. C. S. Baker, 1924	,
indica—Riparia riparia indica Whistler, 1916	18
indicator—Criniger indicator J. & E. Verreaux, 1885 J. Orn. 1855: 105. HOLOTYPE, Adult. Reg. no. 1854.5.23.1. Gaboon. Purchased of Verreaux.	21
indicus—Sturnus indicus Hodgson, 1844	62
indigo—Muscicapa indigo Horsfield, 1821	1
indigotica—Diglossa indigotica P. L. Sclater, 1856	51

indistincta—Meliphaga (Melithreptus) indistincta Vigors 48 mount). Reg. no. 1863.7.7.65. King George's Sound, Australia. Collected by R. Brown. Presented by the Linnean Society of London as the type. The bill is damaged. The description states that the specimen is in bad condition. indochina—Cyornis rufigastra indochina Chasen and Kloss, 1928 . 37 Bull. Brit. Orn. Cl., 48:73. HOLOTYPE, Adult female. Reg. no. 1947.60.34. Daban, South Annam, 22 March 1918. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore. indochinensis—Horizillas rufifrons indochinensis Robinson & Kloss, J. F. M. S. Mus., 10: 205. SYNTYPE, Adult male. Reg. no. 1947. 60.54. Trangbom, Cochin China, 4 June 1918. Collected by H. C. Robinson and C. B. Kloss and presented by the Raffles Museum, Singapore. The female syntype is also in the collection. indochinensis—Lalage fimbriata indochinensis Kloss, 1915 . . 20 Bull. Br. Orn. Cl., 46: 7. HOLOTYPE, Adult male. Reg. no. 1947. 6.33. Dran, 3,000', south Annam, 9 May 1918. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore. indochinensis—Tchitrea affinis indochinensis Salomonsen, 1933 Ibis, (13) 3:734. HÖLOTYPE, Adult male. Reg. no. 1928.6.26. 1052. Angkor, Cambodia, 22 Dec. 1929. Collected by J. Delacour, on the fourth Franco-British Indo-China Expedition of J. Delacour and W. P. Lowe. indochinensis—Trochalopterum milnei indochinensis Delacour, 1927 . 32B Bull. Br. Orn. Cl., 47:158. HOLOTYPE, Adult male. Reg. no. 1927.5.6.11. Tam Dao, central Tonkin, 5 Dec. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China. indrapurae—Turdus indrapurae Robinson & Kloss, 1916 J. Straits Br. roy. Asiat. Soc., 73:277. SYNTYPE, Adult male. Reg. no. 1920.6.29.446. Korinchi Peak, west Sumatra, 27 April 1914. Collected and presented by H. C. Robinson and C. B. Kloss. This is the author's selected male. Their selected female syntype is also in the collection.

1888.1.13.684. South America (Restricted to Bahia, vide Berlepsch and Hartert, Novit. Zool., 9 (1902): 42).

Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 125.

Sclater states on the specimen label "type belongs to Bremen

J. Orn. 1853: 35. HOLOTYPE, Adult (relaxed mount). Reg. no.

inerme—Ornithion inerme Hartlaub, 1853

Museum". It would appear that he received the specimen from Bremen but that it subsequently came to the British Museum with his collection.
infelix—Monarcha infelix P. L. Sclater, 1877
infumata—Leucocirca infumata Hume, 1873
infuscata—Saxicola infuscata Smith, 1839
infuscatus—Aethiopsar fuscus infuscatus E. C. S. Baker, 1918 69 Bull. Br. Orn. Cl., 38:70. HOLOTYPE, Adult male. Reg. no. 1908.5.30.131. Legan, Lower Chindwin, Burma, 20 Jan. 1906. Collected by A. Mears. Purchased of E. Gerrard.
infuscatus—Anabates infuscatus P. L. Sclater, 1856
inglisi—Munia inglisi Hume, 1877
inglisi—Prinia socialis inglisi Whistler and Kinnear, 1933 35 J. Bombay nat. Hist. Soc., 36: 574. HOLOTYPE, Adult. Reg. no. 1886.10.1.1660. Bhotan Duars, India, Jan. 1876. Collected by L. Mandelli. Hume Collection.
innesi—Scotocerca inquieta innesi Nicoll, 1908 35 Bull. Brit. Orn. Cl., 21:98. HOLOTYPE, Adult male. Reg. no. 1908.5.22.1. Wadi Hof, N'Giza, near Cairo, Egypt, 14 Feb. 1908. Collected and presented by M. J. Nicoll.
innexa—Buchanga innexa Swinhoe, 1870

inornata—Myrmotherula inornata P. L. Sclater, 1890
inornata—Nemosia inornata Taczanowski, 1879
inornata—Newtonia brunneicauda inornata Salomonsen, 1934 37 Ibis, (13) 4:382. Holotype, Adult male. Reg. no. 1931.8.18. 2884. Ampotaka, Western Madagascar, 13 March 1930. Presented by the Franco-Anglo-American Archbold-Vernay Expedition.
inornata—Parula inornata Baird, 1864
inornata—Pitta inornata G. R. Gray, 1860
inornata—Prinia inornata Sykes, 1832 Proc. zool. Soc. Lond., 1832: 89. SYNTYPE, Adult female (relaxed mount). Reg. no. 1880.1.1.3005. Deccan, India. Collected by W. H. Sykes. Presented by the Honourable East India Company. This specimen was presented as the type but N. B. Kinnear (M.S. note, c. 1933) states that two specimens, of which this is one, were listed in the catalogue of the collection of the Honourable East
India Company. inornata—Zosterops inornata Layard, 1878
inornatum—Dicaeum inornatum Sharpe 1885

no. 1843.1.13.245. Nepal. Collected and presented by B. H.

Hodgson.

The name originated as a Hodgson MS name. It appeared as a nomen nudum in J. E. Gray, List of Mammals and Birds of Nepal... of B. H. Hodgson, (1846): 60,151, and in Sharpe, Proc. zool. Soc. Lond., 1883 (publ. 1884): 580.

- inornatus—Buarremon inornatus P. L. Sclater and Salvin, 1879 . 49

 Ibis, (9) 3:427. SYNTYPE, Adult (female, on label). Reg. no.
 1885.6.12.902. Pallatanga, Ecuador. (Sept. 1858, on label.)
 Collected by L. Fraser. Sclater Collection.

 There are three other syntypes in the collection.

- inornatus—Tropidorhynchus inornatus G. R. Gray and Mitchell, 1846 48 Genera of Birds, 1: pl. 39. Holotype, Adult (relaxed mount). Reg. no. 1846.7.16.190. (Timor, South-west Pacific, on label.) Collected and presented by E. Wilson.

Although labelled as the type this specimen was not referred to by Gadow, Cat. Bds. Brit. Mus., 9 (1884): 275, and was probably on display in the public gallery as a mounted specimen at that period.

- inquietus—Myiarchus inquietus Salvin and Godman, 1889 . . . 9
 Biol. Centr. Amer., Aves 2:88. Holotype, Adult female. Reg.
 no. 1899.4.20.1479. Acaguisotla, 3,500', Guerrero, Mexico, 18
 Oct. 1888. Collected by H. H. Smith, and presented by his wife to the Salvin-Godman Collection.

200 inquisitor
inquisitor—Rhynchocyclus poliocephalus inquisitor Chubb, 1920 . 9 Bull. Brit. Orn. Cl., 40: 108. Holotype, Adult male. Reg. no. 1922.3.5.2661. Bartica Grove, British Guiana, 1911. Collected by F. V. McConnell. McConnell Collection.
insignis—Artanus insignis P. L. Sclater, 1877 67 Proc. zool. Soc. Lond., 1877 : 101, pl. 15. Holotype, Adult female. Reg. no. 1888.9.20.574. New Ireland, March 1876. Collected by G. Brown. Tweeddale Collection.
insignis—Drymoipus insignis Hume, 1872
insignis—Euphonia insignis P. L. Sclater and Salvin, 1877 51 Proc. zool. Soc. Lond., 1877: 521. pl. 52. SYNTYPE, Adult male. Reg. no. 1885.6.4.58. Jina (= Jima), Ecuador. Collected by C. Buckley. Salvin-Godman Collection. The other syntype is in the collection.
insignis—Nemosia insignis P. L. Sclater, 1856 51 Proc. zool. Soc. Lond., 1856: 110. Syntype, Adult male. Reg. no. 1885.6.12.806. South Brazil (vide Hellmayr and Conover, Catalogue of birds of the Americas, 9 (1936): 379). (Collected for Argent, on label). Sclater Collection. The female and immature syntypes are also in the collection.
insignis—Rhinomyias insignis Ogilvie-Grant, 1895 37. Bull. Brit. Orn. Cl., 4: 40, (and Ibis, (7) 1: 442; pl. 12.) SYNTYPE, Adult male. Reg. no. 1897.5.13.245. Mount Data, 8,000' Lepanto District, North-west Luzon, Philippines, (29 Jan. 1895, on label). Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. A female syntype is also in the collection, and another in the American Museum of Natural History, vide Hachisuka, Birds of the Philippines, 2 (1935): 331 who listed the male as the type.
insignis—Saxicola insignis J. E. Gray, 1846
insignis—Sycobrotus insignis Sharpe, 1891 61 Ibis, (6) 3:117, pl. 6. HOLOTYPE, Adult female. Reg. no. 1893. 12.1.6. Mount Elgon, East Africa. Collected and presented by F. J. Jackson.

Ibis, (5) 2:450. SYNTYPE, Adult male. Reg. no. 1889.7.10.131. Roraima, 5,000', British Guiana, 13 Sept. 1883. Collected by H. Whitely. Salvin-Godman Collection. There are two other syntypes in the collection.	
insolens—Agriornis insolens P. L. Sclater and Salvin, 1869 9 Proc. zool. Soc. Lond., 1869: 153. SYNTYPE, Adult male. Reg. no. 1888.1.13.14. Tinta, South Peru, 21 May 1868. Collected by H. Whitely. Sclater Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 7.	
There are several other syntypes in the collection. insolens—Corvus insolens Hume, 1874	
insperatus—Parus insperatus Swinhoe, 1866 42	
Ibis, (2) 2:308. SYNTYPE, Adult. Reg. no. 1872.5.27.54. Fungshan Mountains, Formosa, Dec. 1865. Collected for R. Swinhoe. Purchased of H. Whitely, who obtained it from Swinhoe. A second syntype, from the Tweeddale Collection, is also in the collection.	
insulae—Pycnonotus luteolus insulae Whistler & Kinnear, 1932 . 21	
J. Bombay nat. Hist. Soc., 35:759. HOLOTYPE, Adult female. Reg. no. 1928.6.25.46. Tellula, Ceylon, 8 April 1928. Collected and presented by W. P. Lowe.	
insulana—Cossypha insulana. See appendix, Corrections and Additions, vol. 2	
insularis—Acrocephalus insularis Wallace, 1862	
insularis—Aethopyga siparaja insularis Delacour and Jabouille, 1928 46 Bull. Brit. Orn. Cl., 48: 134. HOLOTYPE, Adult male. Reg. no. 1928.6.26.11. Phu-Quoc Island, Coast of Cochin China, 25 Dec. 1927. Collected by J. Delacour and P. Jabouille on the 4th Expedition to Indo-China. Presented by J. Spedan Lewis.	
insularis—Amadina insularis Wallace, 1864 60 Proc. zool. Soc. Lond., 1863 (publ. 1864) : 495. SYNTYPE, Adult	
male. Reg. no. 1873.5.12.1381. East Timor, 1861. Collected by A. R. Wallace. Wallace Collection. Listed as the type by Sharpe, <i>Cat. Bds. Brit. Mus.</i> , 13 (1890): 312. There are several other syntypes in the collection.	

insignis—Thamnophilus insignis Salvin & Godman, 1884

- insularis—Buchanga insularis Sharpe, 1877. Cat. Bds. Brit. Mus., 3:253. SYNTYPE, Adult male. Reg. no. 1876.6.6.7. (M.P. Galla, Uva, 4,000', on label), East Ceylon, 20 Sept. 1875. Collected and presented by W. V. Legge. There are two other syntypes in the collection. insularis—Chloropsis aurifons insularis Whistler & Kinnear, 1932 . 22 J. Bombay nat. Hist. Soc., 35:752. HOLOTYPE, Adult male. Reg. no. 1886.9.1.740. Cotta, N.P., Ceylon, 18 June 1878. Collected by the Hart brothers. Hume Collection. insularis—Dissemurus paradiseus insularis Hachisuka. 1926 Bull. Brit. Orn. Cl., 47:58. HOLOTYPE, Adult male. Reg. no. 1886,3,1,2416. Sarawak, Borneo, 20 Sept. 1877. Collected for A. O. Hume. Hume Collection. insularis—Fringillaria insularis Ogilvie-Grant and Forbes, 1889 Bull. Liverpool Museum, 2:2. SYNTYPE, Adult male. Reg. no. 1899.8.11.60. Sokotra. (Adho Dimellus, 3,500', 10 Feb. 1899, on label.) Collected by W. R. Ogilvie-Grant and H. O. Forbes. Presented by the Royal Society. This is the authors' selected male syntype and the label is inscribed "type." The selected female and other syntypes are also in the collection. insularis—Hylophilus insularis P. L. Sclater, 1861 Proc. zool. Soc. Lond., 1861:128. HOLOTYPE, Adult. Reg. no. 1886.9.15.310. Tobago. Collected by J. Kirk. Sclater Collection, obtained by him from Sir W. Jardine. insularis—Myiophoneus insularis Gould, 1862 31 Proc. zool. Soc. Lond., 1862: 280. SYNTYPE, Adult. Reg. no. 1863.2.16.6. Northern Formosa. Collected by R. Swinhoe. Swinhoe Collection. There is a second syntype in the collection. insularis—Passer insularis P. L. Sclater and Hartlaub, 1881 Proc. zool. Soc. Lond., 1881: 169, pl. 16. SYNTYPE, Adult male. Reg. no. 1881.3.21.20. Socotra, 22 March 1880. Collected by I. B. Balfour. Presented by the Socotra Committee. There are three other syntypes in the collection. insularis—Phoenicothraupis insularis Salvin, 1888. Ibis, (5) 6:259. SYNTYPE, Adult male. Reg. no. 1885.9.9.753. Maco Island, off Northern Yucatan, Nov. 1885. Collected by G. F. Gaumer. Salvin-Godman Collection. There are several other syntypes in the collection. insularis—Pycnonotus plumosus insularis Chasen & Kloss, 1929 . 21
- insularis—Pycnonotus plumosus insularis Chasen & Kloss, 1929 . 21 J. Orn., 1929 (2): 115. HOLOTYPE, Adult male. Reg. no. 1947. 60.48. Banguey Island, British North Borneo, 3 Sept. 1927. Collected by F. N. Chasen & C. B. Kloss and presented by the Raffles Museum, Singapore.

in	Itacta—Pachycephala intacta Sharpe, 1900
in	Ritensa—Pogonocichla intensa Sharpe, 1901
in	tensiflava—Arachnothera chrysogenys intensiflava E. C. S. Baker,
	1925
in	tensum—Dicaeum chrysorrhaeum intensum E. C. S. Baker, 1921 . 45
	Bull. Brit. Orn. Cl., 41:108. HOLOTYPE, Adult. Reg. no. 1897. 12.10.942. Sikkim, nothern India, Nov. 1873. (probably collected by L. Mandelli). Presented by Col. J. Biddulph.
in	itensus—Serilophus intensus Robinson & Kloss, 1916 1
	J. Straits Branch Roy. Asiat. Soc., 73: 276. SYNTYPE, Adult male. Reg. no. 1920.6.29.192. 3,100', Siolak Dras, Korinchi Valley, west Sumatra, 19 March 1914. Collected and presented by H. C. Robinson and C. B. Kloss. There is a female syntype in the collection.
in	tercedens—Ptilorhis intercedens Sharpe, 1882
	J. Linn. Soc. Lond., 16: 444. SYNTYPE, Adult female. Reg. no. 1880.9.13.49. East Cape, South-east New Guinea. Collected by C. Hunstein. Purchased of E. Gerrard. The male syntype is in the collection.
in	terioris—Monticola imerina interioris Salomonsen, 1934
	Novit. Zool., 39:211. HOLOTYPE, Adult male. Reg. no. 1931. 8.18.2467. Manjakatompo, Ankaratra Mountains, east-central Madagascar, 14 May 1929. Collected by J. Delacour on the Archbold-Vernay Expedition.

interjector—Saltator cayanus interjector Chubb, 1921 50

Ann. Mag. nat. Hist., (9) 8:445. Holotype, Adult male. Reg.
no. 1903.12.20.100. Serra da Chapada, Matto Grosso, Brazil,
29 June 1902. Collected by A. Robert. Presented by Mrs Sladen,
from the Percy Sladen Expedition to Central Brazil.

 1890.1.29.80. Shanghai, Jan. Collected by Capt. Blakiston. Seebohm Collection.

There is another syntype in the collection.

There are several other syntypes in the collection.

Renamed C. familiaris ripponi by Kinnear, op. cit. 49 (1929): 109 on the grounds of preoccupation by C. intermedia of Müller, in

Linnaeus Syst. Nat. Suppl., (1776): 98.

- intermedia—Cryptolopha intermedia La Touche, 1898 35

 Bull. Brit. Orn. Cl., 7:37. Adult female. Reg. no. 1898.8.12.1.

 This specimen was one of the two original syntypes, vide La Touche, Ibis, (11) 5 (1923): 408–9, but was subsequently recognised as belonging to another population, and La Touche appears to have designated the male as a lectotype, reducing this specimen to a paralectotype.

This specimen was labelled as the type but there are several other

syntypes in the collection.

intermedia—Euphonia nigricollis intermedia Chubb, 1910 . . . 51

Ibis, (9) 4:624. ?Holotype. Adult male. Reg. no. 1885.6.4.68.

Roraima, 3,500′, British Guiana, 9 May 1883. Collected by H.

Whitely. Salvin-Godman Collection.

Listed as type by Hellymayr and Conover, Catalogue of birds of the Americas, 9 (1936): 18. W. L. Sclater (MS. notes, now destroyed) stated "this bird must obviously have been the type of

Chubb's race, though not specifically mentioned."

- intermedia—Paradigalla intermedia Ogilvie-Grant, 1913 . . . 70 Bull. Brit. Orn. Cl., 31:105. Holotype, Sub-adult male. Reg. no. 1916.5.30.1072. Camp 6C, Pundak-Padang, 5,500', Utakwa River, Dutch New Guinea, 25 Feb. 1913. (Collected by C. B. Kloss, on label). Presented by the British Ornithologists' Union New Guinea Expedition.

intermedia—Pitta soror intermedia Delacour, 1928 12 Bull. Brit. Orn. Cl., 49: 49. HOLOTYPE, Adult male. Reg. no. 1928.6.26.90. Phuqui, nothern Annam, southern Indo-China, 9 March 1928. Collected by J. Delacour, W. P. Lowe, and P. Jabouille, on the 4th Anglo-French Expedition to Indo-china.
intermedia—Prinia intermedia Jackson, 1910
intermedia—Prionops intermedia Sharpe, 1901 23 Bull. Br. Orn. Cl., 11:47. HOLOTYPE, Adult. Reg. no. 1897.12. 9.12. Teita, British East Africa, 9 June 1897. Collected and presented by C. Stuart Betton.
intermedia—Saxicoloides fulicata intermedia Whistler & Kinnear, 1933
intermedia—Stizorhina vulpina intermedia Clarke, 1913 31 Bull. Br. Orn. Cl., 31:107. HOLOTYPE, Adult male. Reg. no. 1906.12.11.52. Bunjako, Entebbe, Uganda, 10 Nov. 1905. Collected by and purchased of E. Degen.
intermedia—Strepera intermedia Sharpe, 1877
intermedia—Suthora poliotis intermedia Kinnear, 1944 33 Ibis, 86: 70. HOLOTYPE, Adult female. Reg. no. 1935.4.5.386. Yonpu La, 8,000', near Trashigong, east Bhutan, 2 Nov. 1934. Collected and presented by F. Ludlow and G. Sherriff. Renamed Paradoxornis poliotis crocotius by Kinnear, 1954 (q.v.) on grounds of preoccupation.
intermedia—Timalia pileata intermedia Kinnear, 1924 32B Bull. Br. Orn. Cl., 45: 9. HOLOTYPE, Adult male. Reg. no. 1888. 4.20.938. Tonghoo, Burma, 19 Dec. 1874. Collected by R. G. Wardlaw Ramsay. Tweeddale Collection.

There are several other syntypes in the collection.

- intermedia Yuhina nigrimentum intermedia Rothschild, 1923 . 32B Bull. Brit. Orn. Cl., 43:11. HOLOTYPE, Adult male. Reg. no. 1922.12.7.280. Mekong-Salween Divide, Yunnan, 27 July 1921. Collected by G. Forrest. Presented by S. R. Clarke.
- male. Reg. no. 1873.5.12.111. Macassar, South Celebes Group, 1856. Collected by A. R. Wallace. Wallace Collection. Wallace used a G. R. Gray MS. name. This specimen was listed as the type by Sharpe, Cat. Bds. Brit. Mus., 9 (1884): 185, but

Wallace referred to specimens from two localities.

- intermedium—Pellorneum intermedium Sharpe, 1883 Cat. Bds. Brit. Mus., 7:519, pl. 13. SYNTYPE, Adult female. Reg. no. 1881.10.17.6. Dilkooska, Cachar, India, 11 May 1878. Collected by J. Inglis and purchased of Watkins and Doncaster. The immature syntype is also in the collection.
- intermedius—Allotrius intermedius Hume, 1877 . 32B Stray Feathers, 5:112, 115. SYNTYPE, Adult male. Reg. no. 1886.11.1.855. Halfway between Panaduba and Mooleyit, Tenasserim, Burma, 28 Jan. 1877 (on label). Collected by W. Davison (on label). Hume Collection.
- intermedius—Parus afer intermedius Shelley, 1900 42 Birds of Africa, 2:223, 241. HOLOTYPE, Adult female. Reg. no. 1890,12,20,85. Potchefstroom, Transvaal, 7 June 1877. Collected by T. Ayres for J. H. Gurney. Presented by A. G. Gurney Jnr. for J. H. Gurney.
- intermedius--Philentoma intermedius Hume, 1880 Stray Feathers, 9:113. HOLOTYPE, Adult female. Reg. no. 1886. 4.1.3081. Gunong Palai, Johore, Malaya, 6 March 1880. Collected by W. Davison. Hume Collection.
- intermedius—Ploceus intermedius Rüppell, 1845. Systematische Uebersicht der Vögel nord-ost Afrika's: 71. Holo-TYPE, Adult male (relaxed mount). Reg. no. 1861.5.8.41. Shoa, Abyssinia. Collected by W. C. Harris. Presented by the Honourable East India Company Museum.
- intermedius—Schoeniparus intermedius Rippon, 1900 . Bull. Brit. Orn. Cl., 11:11. HOLOTYPE, Adult. Reg. no. 1900. 12.20.285. Nanoi, Loi Mai, South Shan States, 27 March 1899. Collected and presented by G. Rippon.
- intermedius—Sphenoeacus intermedius Shelley, 1882 35 Proc. zool. Soc. Lond., 1882:337. HOLOTYPE, Adult. Reg. no. 1895.5.1.968. Kaffraria, South Africa, 25 Dec. 1876. Collected by Barratt. Shelley Collection.

iracunda 267
interposita—Chalcoparia singalensis interposita Robinson and Kloss, 1921
interpositor—Kittacincla malabarica interpositor Robinson, 1922 . 31
J.F.M.S. Mus., 10: 262. SYNTYPE, Adult male. Reg. no. 1936.4 12.1497. Daban, south Annam, 14 March 1918. Collected by R. B. Kloss and presented by the Raffles Museum Singapore. This is the author's selected male. His selected female syntype is also in the collection.
interscapularis—Megalurus interscapularis P. L. Sclater, 1880
Proc. zool. Soc. Lond., 1880: 65, pl. 6. HOLOTYPE, Adult. Reg. no. 1881.3.29.25. New Britain. Collected by and purchased of the Rev. G. Brown, from his fifth collection.
interciacion Duo conductio vin eleministrativi del E. C. C. Delem 1025 50

- intersicolor—Procarduelis nipalensis intensicolor E. C. S. Baker, 1925 59 Bull. Brit. Orn. Cl., 45:92. HOLOTYPE, Adult male. Reg. no. 1922.12.7.455. Mekong-Salwin Divide, north-west Yunnan, 20 Sept. 1921. Collected by G. Forrest. Stephenson Clarke Collection.
- iodura—Oreocincla iodura Gould, 1872 Ann. Mag. nat. Hist., (4) 9:401. HOLOTYPE or SYNTYPE. Adult. Reg. no. 1881.5.1.737. Queensland. Gould Collection.

 Listed as type by Seebohm, *Monograph of the Turdidae*, 1 (1902): 23
- iora—Phoenicomanes iora Sharpe, 1874 Proc. zool. Soc. Lond., 1874: 427. HOLOTYPE, Adult. Reg. no. 1874.6.13.3. Jamaica (error, Malacca, vide Sharpe, Ann. Mag. nat. Hist., (4) 16 (1875): 236). Purchased of P. Boucard.
- iouschistos—Parus iouschistos Blyth, 1845 in Hodgson and Blyth, J. Asiat. Soc. Bengal 13, 1844 (publ. April 1845): 943. Described by Blyth from a specimen in the Museum of the Asiatic Society of Bengal, in Calcutta: now in the collection of the Zoological Survey of India. The specimens referred to as types by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 58, are not valid. Hodgson's name in Gray, Zoological Miscellany, (1844): 83, is a nomen nudum. Hodgson's description of his own specimens in Ann. Mag, nat. Hist., 15 (1845): 326, was not apparently published until May of that year.
- iracunda—Pipra iracunda Salvin and Godman, 1884 . Ibis, (5) 2:447. HOLOTYPE, Adult male. Reg. no. 1888.1.20.36. Roraima, 3,500', British Guiana, 20 June 1883. Collected by H. Whitely. Salvin-Godman Collection.

irbii—Acredula irbii Sharpe and Dresser, 1871
iris—Pitta iris Gould, 1842
irwini—Calornis irwini Hume, 1873
isabellae—Cossypha isabellae G. R. Gray, 1862
isolata—Aethopyga gouldiae isolata E. C. S. Baker, 1925 40 Bull. Brit. Orn. Cl., 46:13. Holotype, Adult male. Reg. no. 1886.12.1.472. (Aimole, on label)., Manipur, India, (22 April 1881, on label). Collected by A. O. Hume. Hume Collection.
isthmicus—Saltator istlunicus P. L. Sclater, 1861
isura—Cysticola isura Gould, 1847
itoculo—Phyllastrephus alfredi itoculo Vincent, 1933 21 Bull. Br. Orn. Cl., 53:134. HOLOTYPE, Adult male in non- breeding condition. Reg. no. 1933.3.1.1. Near Netia, 600', 14°44' S., 40°04' E., Mozambique Province, Portuguese East Africa, 2 June 1932. Collected and presented by J. Vincent.

ituribisciensis—Thryothorus oyapocensis ituribisciensis Brabourne & Chubb, 1912	28
Ann. Mag. nat. Hist., (8) 10: 262. SYNTYPE, (Immature on label). Reg. no. 1922.3.5.3471. Ituribisci River, British Guiana, 1909. Collected by F. V. McConnell. McConnell Collection. There are several other syntypes in the collection.	
jabouillei—Coracina polioptera jabouillei Delacour, 1951	20
jacksoni—Apalis jacksoni Sharpe, 1891	35
jacksoni—Bathmocercus jacksoni Sharpe, 1902	35
jacksoni—Calamocichla jacksoni Neumann, 1901	35
jacksoni—Cryptospiza jacksoni Sharpe, 1902	60
jacksoni—Drepanoplectes jacksoni Sharpe, 1891 Ibis, (6) 3: 246, pl. 5. SYNTYPE, Adult male. Reg. no. 1893.12.1.3. Masai Land (north of Lake Nakuro, on label), Kenya, 22 July 1890. Collected and presented by F. J. Jackson. The female syntype is also in the collection.	61
jacksoni—Dryoscopus jacksoni Sharpe, 1901	23
Jacksoni—Mirafra poecilosterna jacksoni Ogilvie-Grant, 1913 . 1 Ibis, (10) 1:586. HOLOTYPE, Adult. Reg. no. 1912.12.9.1. Kikuyu, 5,500', 8 Aug. 1904 (in text; Mbagathi R., 8 July on label). Collected and presented by F. J. Jackson.	17
jacksoni—Nectarinia jacksoni Neumann, 1899	16

Jacksoni—Parisoma jacksoni Sharpe, 1899
jacksoni—Platystira jacksoni Sharpe, 1891
jacksoni—Ploceus jacksoni Shelley, 1888 61 Ibis, (5) 6: 293, pl. 7. HOLOTYPE, Adult male. Reg. no. 1888.6. 20.1. Kilimanjaro, Tanganyika. Collected for and presented by F. J. Jackson.
jacksoni—Sylviella jacksoni Sharpe, 1897
jacksoni—Turdinus jacksoni Sharpe, 1900
jagoensis—Estrilda jagoensis Boyd Alexander, 1898 60 Ibis, (7) 4:85. Syntype, Adult male. Reg. no. 1911.12.23.3362. Santiago, Cape Verde Islands, 2 Dec. 1897. Collected by Boyd Alexander. Boyd Alexander Bequest. There are several other syntypes in the collection.
jagoensis—Pyrgita jagoensis Gould, 1837
jalla—Pastor jalla Horsfield, 1821
jamaicae—Sycalis jamaicae Sharpe, 1888

jamesi—Cisticola subruficapilla jamesi Lynes, 1930	
jamesii—Phonygama jamesii Sharpe, 1877	
jamesoni—Diaphorophyia jamesoni Sharpe, 1890	
jamesoni—Hyphantornis jamesoni Sharpe, 1890	
jamesoni—Lagonosticta jamesoni Shelley, 1882	60
jamesoni—Pholidornis jamesoni Shelley, 1890	60
janeti—Ammomanes deserti janeti R. Meinertzhagen, 1933 1 Bull. Br. Orn. Cl., 53:151. HOLOTYPE, Adult female. Reg. no. 1939.12.9.269. Oued Tamanrasset, 5,200', Ahaggar Plateau, Sahara, 23 Feb. 1931. Collected by R. Meinertzhagen. Rothschild Bequest.	17
japonica—Loxia curvirostra japonica. See appendix, Corrections and additions, vol. 2	59
japonicus—Parus palustris japonicus Seebohm, 1879 4 Ibis, (4) 3: 32. Syntype, Adult. Reg. no. 1898.9.20.315. Hakodadi, Japan, January. Collected by T. Blakiston. Seebohm Collection. The second syntype is also in the collection.	42

jardineii—Crateropus jardineii Smith, 1836
purchased of G. A. Frank. Listed as type by Sharpe, Cat. Bds. Brit. Mus., loc. cit. Stenhouse, Scottish Nat., 1930: 150, is in error when he states that no original specimen is in existence.
jardinii—Spodiornis jardinii P. L. Sclater, 1866
javana—Hypothymis azurea javana Chasen and Kloss, 1929 35 Bull. Raffles Mus., 2:22. Syntype, Adult female. Reg. no. 1947.60.36. Badjoelmati, East coast of Java, 31 Jan. 1920. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore, as one of the types. This is the author's selected female. His selected male syntype is also in the collection.
javana—Kittacincla malabarica javana Kloss, 1921 31 J.F.M.S. Mus., 10:210. SYNTYPE, Adult male. Reg. no. 1927. 4.18.538. Karangbolang, south coast of mid-Java, 23 Feb. 1920. Collected by C. B. Kloss and presented by H. C. Robinson. The female syntype is also in the collection.
javanica—Mirafra javanica Horsfield, 1821
javanica—Nectarinia javanica Horsfield, 1821
javanica—Sylvia javanica Horsfield, 1821
javanicus—Eurylaimus javanicus Horsfield, 1821

javanicus—Turdus javanicus Horsfield, 1821
javanus—Dicrurus macrocercus javanus Kloss, 1921 64 J. Federated Malay States Mus., 10: 208. Holotype, Adult male. Reg. no. 1947.60.25. Badjoelmati, Besoeki, East Java, 3 Feb. 1920. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore.
javensis—Ceblephyris javensis Horsfield, 1821
javensis—Copsychus saularis javensis Chasen & Kloss, 1930 31 Bull. Raffles Mus., 4:2, 89. HOLOTYPE, Adult male. Reg. no. 1947.60.40. Wynkoops Bay, south-west Java, 18 March 1920. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore.
javensis—Meliphaga javensis Horsfield, 1821
jebelmarrae—Anthus sordidus jebelmarrae Lynes, 1920 19 Bull. Br. Orn. Cl., 41:16. HOLOTYPE, Adult male. Reg. no. 1920.12.22.538. Jebel Marra, Darfur, Sudan, 12 April 1920 (on label; 1912 in text). Collected and presented by H. Lynes.
jebelmarrae—Fringillaria striolata jebelmarrae Lynes, 1920 49 Bull. Br. Orn. Cl., 41: 34. HOLOTYPE, Adult male. Reg. no. 1920.12.22.595. Jebel Marra, Darfur, Sudan, 1 April 1920. Collected and presented by H. Lynes.
jebelmarrae—Lanius excubitor jebelmarrae Lynes, 1923 23 Bull. Br. Orn. Cl., 43:94. HOLOTYPE, Adult male. Reg. no. 1922.12.8.1238. Jebelmarra highlands, 7,900', Darfur, Sudan, 28 Nov. 1921. Collected and presented by the Lynes-Lowe Expedition.
jebelmarrae—Saxicola torquata jebelmarrae Lynes, 1920 31 Bull. Br. Orn. Cl., 41:17. Holotype, Adult male. Reg. no. 1920.12.22.252. Jebel Marra, 8,000', Darfur, Sudan, 6 April 1920. Collected and presented by H. Lynes.
jefferyi—Eudrepanis jefferyi Ogilvic-Grant, 1894

5.13.93. La Trinadad, Benguet District, North-west Luzon, Philippines, 7 Jan. 1894. Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. There are three other syntypes in the collection.
jerdoni—Abrornis jerdoni Brooks, 1871
jerdoni—Accentor jerdoni Brooks, 1872
jerdoni—Drymoica jerdoni Blyth, 1847
jerdoni—Oreicola jerdoni Blyth, 1867
jerdoni—Timalia jerdoni Walden, 1872 32B Ann. Mag. nat. Hist., (4) 10:61. Syntype, Adult. Reg. no. 1888.4.20.927. Khasia Hills, Assam. Collected by A. W. Chennell. Tweeddale Collection.
jerdonii—Merva jerdonii Hodgson, 1847
jessei—Alaemon jessei Finsch, 1870
jessei—Pytelia jessei Shelley, 1903

jewan—Hirundo jewan Sykes, 1832
Sitz. Akad. Wiss. Wien., 70: 124. HOLOTYPE, Adult male. Reg. no. 1895.5.1.2342. Ansus, Jobi Island, New Guinea, (April 1873, on label). Collected by A. B. Meyer. Shelley Collection.
jocosus—Campylorhynchus jocosus P. L. Sclater, 1859 28
Proc. zool. Soc. Lond., 1859: 371. SYNTYPE, Adult male. Reg. no. 1886.9.15.19. Oaxaca, Mexico, March 1858. Collected by A. Boucard for A. Sallé. Sclater Collection. The second syntype is also in the collection. See Sclater, Catalogue of American Birds, (1862): pl. 3.
johannae—Prionochilus johannae Sharpe, 1888
johannis—Warsangli johannis Clarke, 1919
johnsi—Aethopyga sanguinipectus johnsi Robinson and Kloss, 1919 46 Ibis, (11) 1:621, pl. 18. HOLOTYPE, Adult male. Reg. no. 1919. 12.20.507. Dran, 3,000', South Annam, 15 May 1919. Collected and presented by C. B. Kloss.
johnstonei—Euethia johnstonei Lowe, 1906
johnstoni—Merula johnstoni Sharpe, 1906
johnstoni—Musicapa johnstoni Shelley, 1884

johnstoni—Nectarinia jolunstoni Shelley, 1885
johnstoni—Pogonocichla johnstoni Shelley, 1893
johnstoni—Seicercus ruficapilla johnstoni W. L. Sclater, 1927 35 Bull. Brit. Orn. Cl., 48:13. Holotype, Adult male. Reg. no. 1897.11.4.73. Kombi, 7,000′, Masuka Range, north-west of Lake Nyasa, July 1896. Collected by A. Whyte. Presented by H. H. Johnston.
johnstoniae—Ianthia johnstoniae Ogilvie-Grant, 1906 31 Bull. Br. Orn. Cl., 16:118. Syntype, Adult male. Reg. no. 1907.12.12.39. (Racu Racu Mountains, 8,000', central Formosa, Feb. 1906, vide Ogilvie-Grant and La Touche, Ibis, (9) 1 (1907):175, pl. 4.). Collected by and purchased of W. Goodfellow. The female syntype is also in the collection.
johnstoniae—Pericrocotus johnstoniae Ogilvie-Grant, 1905 20 Bull. Br. Orn. Cl., 16:18. SYNTYPE, Adult female. Reg. no. 1905.11.26.84. Mount Apo, 7,000′, south-east Mindanao, Philippines, March 1905. Collected by W. Goodfellow and purchased of Gerrard. The male syntype is also in the collection.
jonesi—Pellorneum ruficeps jonesi E. C. S. Baker, 1920 32B Bull. Br. Orn. Cl., 41:9. HOLOTYPE, Adult male. Reg. no. 1920. 10.28.1. Kalka, 2,000′, Baghat State, north-west Himalayas, 26 Dec. 1919. Collected and presented by A. E. Jones.
josensis—Anthus similis josensis C. H. B. Grant & Mackworth-Praed, 1958
josephae—Vireo josephae P. L. Sclater, 1859
josephinae—Euscarthmus josephinae Chubb, 1914

- kakamegae 277 iosephinae—Synallaxis albigularis josephinae Chubb, 1919 Bull. Br. Orn. Cl., 39:60. HOLOTYPE, Adult male. Reg. no. 1922. 3.5.2403. Mount Roraima, British Guiana, Oct. 1898. Collected by F. V. McConnell. McConnell Collection. iucunda—Pipreola iucunda P. L. Sclater, 1860 Proc. zool. Soc. Lond., 1860: 89, pl. 160. HOLOTYPE, Adult male. Reg. no. 1888.1.13.1632. Cachi-Llacta, near Nanegal, Ecuador, 1859. Collected by L. Fraser. Sclater Collection. iuliae—Nectarophila iuliae Tweeddale, 1877 46 Proc. zool. Soc. Lond., 1877: 547. SYNTYPE, Adult male. Reg. no. 1880.11.18.65. Malanipa, Basilan Straits, Philippines, (vide op. cit., p. 536), 30 Jan. 1875. Collected for J. Murray. From the voyage of H.M.S. Challenger. The two other syntypes are in the collection. juninensis-Muscisaxicola juninensis Taczanowski, 1884 Ornithologie du Perou, 2:214. Adult male. Reg. no. 1888.1.13. 195. This specimen was collected by Jelski with syntypical material at Junin, Central Peru, 1873, but was sent to Sclater in 1874. Taczanowski's description was based on the other specimen at Warsaw. This specimen has no type status but was referred to as an "Authentic specimen" vide Sclater, Cat. Bds. Brit. Mus., 14 (1888): 58, and as "one of the original specimens" vide Hellmayr and Conover, Catalogue of Birds of the Americas, 5 (1927): 23. juninensis—Rhopias spodionota juninensis Chubb, 1918. Bull. Br. Orn. Cl., 38:84. HOLOTYPE, Adult male. Reg. no. 1902.3.13.1234. Perene, 800 m., Junin, Peru, 12 March 1900. Collected by P. O. Simons. Simons Collection.
- juninensis—Upucerthia juninensis Chubb, 1914 Bull. Br. Orn. Cl., 35: 20. HOLOTYPE, Adult. Reg. no. 1914. 12.2.185. Junin, Peru, 20 Feb. 1914. Collected and presented by Lord Brabourne.
- kabalii—Erythropygia leucophrys kabalii C. M. N. White, 1944 . 31 Bull. Br. Orn. Cl., 64:49. HOLOTYPE, Adult male. Reg. no. 1956.16.4. Chikonkwelo stream, Balovale, Northern Rhodesia, 21 Aug. 1943. Collected and presented by C. M. N. White.
- kakamariae—Gymnorhis pyrgita kakamariae Stoneham, 1925. Bull. Br. Orn. Cl., 45:76. SYNTYPE, Adult male. Reg. no. 1929. 8.19.1. Kakamarie, 6-7,000', Karamojo, northern Uganda (north of Kampala), 4 May 1924. Collected and presented by H. F. Stoneham.

The female syntype is in the collection.

kakamegae—Xenocichla kakamegae Sharpe, 1900 Bull. Br. Orn. Cl., 11:29. HOLOTYPE, Adult male. Reg. no. 1900.6.22.3. Kakemega Forest, Nandi, 8 Feb. 1898. Collected and presented by F. J. Jackson.

kalahari—Cisticola kalahari Ogilvie-Grant, 1910 35 Bull. Brit. Orn. Cl., 25: 121. SYNTYPE, Adult female. Reg. no. 1911.12.16.123. Molopo River, 3,000′, Kalahari Desert, 20 April 1909. Collected by R. B. Woosnam, on the Lake Ngami Expedition. The female syntype is also in the collection.
kalahariae—Certhilauda kalahariae Ogilvie-Grant, 1912 17 Ibis, (9) 6: 375. HOLOTYPE, Adult male. Reg. no. 1911.12.16.139. Lehutitu, 3,000', north Kalahari, Ngamiland, 29 May 1909. Collected by R. B. Woosnam and presented by the Ngami Expedition.
kalamantan—Eurylaimus ochromalus kalamantan Robinson & Kloss, 1919
kalulongae—Turdinus kalulongae Sharpe, 1893
kamchatkensis—Nucifraga kamchatkensis Barrett-Hamilton, 1898 71 Bull. Brit. Orn. Cl., 7:46. HOLOTYPE, Adult. Reg. no. 1898.7. 4.59. Kamchatka, (Petropavlowsk, 25 Aug. 1897, on label). Collected and presented by G. E. H. Barrett-Hamilton.
kangrae—Fulvetta vinipectus kangrae Ticehurst & Whistler, 1924 32B Bull. Br. Orn. Cl., 44:71. HOLOTYPE, Adult male. Reg. no. 1924.3.13.1. Bagh Nala, 6,000', Palumpur, Kangra, north-west Himalayas, 27 Dec. 1923. Collected and presented by H. Whistler.
kangrae—Lophophanes dichrous kangrae Whistler, 1932 42 Bull. Brit. Orn. Cl., 53: 20. HOLOTYPE, Adult male. Reg. no. 1932.10.23.1. Koti State, 8,500', near Simla, India, 9 March 1924. Collected by A. E. Jones. Presented by A. Whistler.
kangrae—Phylloscopus pulcher kangrae Ticehurst, 1923 35 Bull. Brit. Orn. Cl., 44: 29. The author selected as a holotype a specimen from the Hume Collection, Reg. no. 1886.7.8.1289., but this cannot now be traced. A paratypical series taken at Simla during the same month by W. Davison is in the collection.
kangrae—Procarduelis nipalensis kangrae Whistler, 1939 59 Bull. Brit. Orn. Cl., 60: 16. HOLOTYPE, Adult male. Reg. no. 1887.6.1.141. Dharmsala, Kangra District, north-west Himalayas, April 1970. Collected by E. W. Brooks. Hume Collection.
kaokoensis—Monticola brevipes kaokoensis Macdonald & Hall, 1857 31 Ann. Transvaal Mus., 23: 24. Syntype, Adult male. Reg. no.

1950.50.343. Kamanjab, 4,000', Kaokoveld, Damaraland, 19 April 1950. Collected by F. O. Cave on the British Museum South West Africa Expedition.

The female syntype is also in the collection.

- karenensis—Aethopyga nipalensis karenensis Ticehurst, 1939 . . . 46
 Ibis, (14) 3:755. HOLOTYPE, Adult male. Reg. no. 1948.80.2580.
 Nattaung, 8,000', Karenni, Burma, 10 April 1939. Collected by P. F. Garthwaite. Presented by P. F. Garthwaite and H. C. Smith.
 Upper mandible of specimen is missing.
- karenni—Alcippe poiocephala karenni Robinson & Kloss, 1922 . 32B J. and Proc. Asiat. Soc. Bengal (new ser.) 18:563. New name for Alcippe magnirostris Walden, 1875, which is preoccupied, vide Moore, Proc. zool. Soc. Lond., 1854:277.
- kasamaensis—Serinus atrogularis kasamaensis Benson, 1955. . . . 59
 Bull. Brit. Orn. Cl., 75: 108. HOLOTYPE, Adult male. Reg. no.
 1955.42.1. Kasama, 4,300', north-eastern Northern Rhodesia,
 2 Sept. 1953. Collected by C. W. Benson. Presented by the
 National Museums, Southern Rhodesia.
- kashmirensis—Dumeticola thoracica kashmirensis Sushkin, 1925. 35

 Proc. Bost. Soc. nat. Hist., 38: 42. Holotype, Adult. Reg. no. 1876.2.12.225. North-western Himalayas. Collected and presented by Capt. Stackhouse Pinwell.
- kashmiriensis—Phylloscopus reguloides kashmiriensis Ticehurst, 1933 35 Bull. Brit. Orn. Cl., 54:19. HOLOTYPE, Adult male. Reg. no. 1886.7.8.975. (Jakko, on label), Simla, Kashmir, 28 Sept. 1877. Collected by W. Davison. Hume Collection.
- kasokae—Prinia subflava kasokae C. M. N. White, 1946 . . . 3

 Ibis, 88:96. Holotype, Adult male. Reg. no. 1956.16.33.

 Mwange Lake, West Balovale, Kuande Lunda Province, Northern Rhodesia, 20 June 1943. Collected and presented by C. M. N. White.
- katanga—Cisticola natalensis katanga Lynes, 1930 35

 Ibis, (12) 6, Supplement: 443. HOLOTYPE, Adult male. Reg. no. 1909.12.31.188. Kambove, 4,500′, Haut Luapula District, Southeast Belgian Congo, 1 March 1907. Collected and presented by S. A. Neave.

Reg.	no.	1909	.12.31	.201.	Kat	tanga,	south	-east	Cong	o, (Lu	fupa
River	, 3	Nov.	1907,	vide 1	bis,	(9) 4	(1910)	: 149)). Co	llected	and
			A. No								
Thi	is sp	ecime	n was	labelle	ed as	the ty	ype bu	t seve	ral oth	ner synt	ypes

were used, one of which is in the collection.

- katangae—Uraeginthus bengalus katangae Vincent, 1934. . . . 60
 Bull. Brit. Orn. Cl., 54: 174. HOLOTYPE, Adult male. Reg. no. 1934.10.19.1. Elizabethville, Belgian Congo, 17 March 1933. Collected by A. W. Vincent.
- kathangorensis—Mirafra hypermetra kathangorensis Cave, 1940. 17

 Bull. Br. Orn. Cl., 60: 96. HOLOTYPE, Adult male. Reg. no. 1939.8.16.4. Near Kathangor, 1,200', Eastern District, Equatoria, Sudan, 13 June 1939. Collected and presented by F. O. Cave.

kathleenae—Batis kathleenae C. M. N. White, 1941 38
Bull. Brit. Orn. Cl., 61:48. HOLOTYPE, Adult female. Reg. no.
1954.58.2. Mwinilunga, Northern Rhodesia, 12 Jan. 1940. Collected by C. M. N. White and received in exchange from him.

- kauriensis—Urocichla kauriensis Harington, 1908 32B

 Ann. Mag. nat. Hist., (8) 2: 246. SYNTYPE, Adult male. Reg. no.
 1908.8.2.21. Watan, Bhamo District, 5,000′, Upper Burma, 8

 April 1908. Collected and presented by H. H. Harington.

 The female syntype is also in the collection.
- kelsalli—Camaroptera brachyura kelsalli W. L. Sclater, 1927. . . 35

 Bull. Brit. Orn. Cl., 48: 16. HOLOTYPE, Adult male. Reg. no. 1914.4.14.57. Tungea, north-north-east of Bo, Sierra Leone, 12

 Sept. 1922. Collected by and purchased of H. J. Kelsall.
- kelsalli—Fraseria ocreata kelsalli Bannerman, 1922 37

 Bull. Brit. Orn. Cl., 42:68. HOLOTYPE, Adult female. Reg. no. 1913.7.6.5. York Pass, Sierra Leone, 1 June 1911. Collected and presented by H. J. Kelsall.

Shelley, *The Birds of Africa*, 5 (1906): 120, makes it clear that this specimen is the holotype but gives the Usambara Mountains as the locality. This might be due to confusion with his other specimen, collected in 1879, which was a "female or young" bird.

- kibaliensis—Trochocercus kibaliensis Boyd Alexander, 1907 . . . 38

 Bull. Brit. Orn. Cl., 19:88. Syntype, Adult male. Reg. no. 1911.
 12.23.1429. Kibali River, Belgian Congo, 6 Aug. 1906. Collected by Boyd Alexander on the Alexander-Gosling Expedition. Boyd Alexander Bequest.

 The famela syntype is also in the collection

The female syntype is also in the collection.

kidepoensis—Mirafra hypermetra kidepoensis Macdonald, 1940 . 17 Bull. Br. Orn. Cl., 60:59. Holotype, Adult male. Reg. no. 1939.10.1.85. Eros, 4,000′, south edge of Didinga Mts., south Sudan, approx. 4°6′ N., 33°44′ E., 17 March 1939. Collected and presented by J. D. Macdonald.

kigomaensis—Symplectes bicolor kigomaensis C. H. B. Grant & Mackworth-Praed, 1956
kigomaensis—Uraeginthus bengalus kigomaensis C. H. B. Grant and Mackworth-Praed, 1956
kikuyuensis—Alethe kikuyuensis Jackson, 1910
kikuyuensis—Xenocichla kikuyuensis Sharpe, 1891 21 Ibis, (6) 3:118. HOLOTYPE, Adult. Reg. no. 1893.12.1.30. Kikuyu, East Africa. Collected and presented by F. J. Jackson.
kikuyuensis—Zosterops kikuyuensis Sharpe, 1891
kilimensis—Alcippe kilimensis Shelley, 1889
kilimensis—Coccopygia kilimensis Sharpe, 1890
kilimensis—Cryptospiza salvadorii kilimensis W. L. Sclater, 1934 . 60 Bull. Brit. Orn. Cl., 55:13. HOLOTYPE, Adult male. Reg. no. 1934.6.18.60., near Ngare, 5,750', Nairobi North River, west side of Kilimanjaro, 7 Feb. 1934. Collected and presented by R. E. Moreau.
kilimensis—Nectarinia kilimensis Shelley, 1884

- kilosa—Turdoides hypoleuca kilosa Vincent, 1935. Bull. Br. Orn. Cl., 55: 176. HOLOTYPE, Adult female. Reg. no. 1935.5.11.1. Kibedya, 3,600', Kilosa, Tanganyika, 22 June 1934. Collected and presented by N. R. Fuggles-Couchman.
- kinabaluensis—Cryptolopha kinabaluensis Sharpe, 1901 1897.7.2.880. Kina Balu, 8,000', North Borneo, 14 Feb. 1888. Collected by J. Whitehead. Seebohm Bequest. This specimen was labelled as the type but there are two other

syntypes in the collection.

- 4526. Jodhpur, Rajputana, India, 29 Oct. 1868. Collected by G. King. Hume Collection.
- kinneari—Actinodura ramsayi kinneari Delacour, 1927. Bull. Br. Orn. Cl., 47:162. HOLOTYPE, Adult male. Reg. no. 1927.6.5.18. Tam Dao, Tonkin, 12 Jan. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indochina.
- kinneari—Ammomanes cinctura kinneari Bates, 1935 Bull. Br. Orn. Cl., 55:140. HOLOTYPE, Adult female. Reg. no. 1903.2.4.19. 50 miles south of Omdurman, Sudan, 9 Oct. 1902. Collected and presented by H. W. Dunn.
- kinneari—Cinnyris habessinicus kinneari Bates, 1935 46 Bull. Brit. Orn. Cl., 55: 120. HOLOTYPE, Adult male. Reg. no. 1935.5.10.41. Near Sail, 4,000', Talf Plateau, Arabia, 14 April 1934 (27 April, on label) Collected and presented by H. St. J. B. Philby.
- kinneari—Dendrocitta rufa kinneari E. C. S. Baker, 1922 Oct. 1875. Collected by R. G. Wardlaw-Ramsay. Tweeddale Collection.
- 1935.5.10.185. Samuda, 4,000', Rakba Plain, Arabia, 25 June 1934. Collected and presented by H. St. J. Philby.
- kinneari—Melidectes belfordi kinneari Mayr, 1936 Bull. Brit. Orn. Cl., 57:42. HOLOTYPE, Adult male. Reg. no. 1916.5.3.654. Camp 9, 5,500', Utakwa River, Dutch New Guinea, 26 Jan. 1913. Collected by C. B. Kloss. Presented by the committee of the British Ornithologists' Union Expedition to New Guinea.

kinneari—Sitta carolinensis kinneari Van Rossem, 1939 43 Proc. Biol. Soc. Washington, 52 : 3. HOLOTYPE, Adult male. Reg. no. 1892.3.1.63. Amula (Tixtla), 6,000′, Guerrero, Mexico, Aug. 1888. Collected by D. W. Smith. Salvin-Godman Collection.
kinnisii—Merula kinnisii Blyth, 1851
kirki—Pytelia kirki Shelley, 1903
kirki—Sigmodius scopifrons kirki W. L. Sclater, 1924 23 Bull. Br. Orn. Cl., 44:92. HOLOTYPE, Adult. Reg. no. 1894.8. 16.236. Lamu, Kenya coast. Collected by Sir J. Kirk. Shelley Collection.
kirki—Zosterops kirki Shelley, 1880
kirkii—Cinnyris kirkii Shelley, 1876
kirkii—Crateropus kirkii Sharpe, 1876
kittlitzi—Fringilla kittlitzi Seebohm, 1890
kleinschmidti—Myzomela kleinschmidti Sharpe, 1884

- lected by T. Kleinschmidt. Purchased of Godeffroy Brothers of Hamburg.
- klossi—Cyornis rubeculoides klossi Robinson, 1921 37

 Bull. Brit. Orn. Cl., 42:12. HOLOTYPE, Adult male. Reg. no. 1947.60.35. Dran, 3,000′, South Annam, 18 May 1918. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore.
- klossi—Pachycephala soror klossi Ogilvie-Grant, 1915 . . . 39

 Ibis, (10) Jubilee Supplement: 88. SYNTYPE, Adult male. Reg.
 no. 1916.5.30.300. Camp 6A, 2,900', Utakwa River, Dutch New
 Guinea. 19 Jan. 1913. Collected by C. B. Kloss. Presented by
 the Committee of the British Ornithologists' Union Dutch New
 Guinea Expedition.

This is the author's selected male syntype. The selected female syntype is also in the collection.

- klossi—Pomatorhinus nuchalis klossi E. C. S. Baker, 1917 32B Bull. Br. Orn. Cl., 38:9. Syntype, Adult male. Reg. no. 1915. 10.14.113. Klong Menao, south-east Siam, 11 Jan. 1915. Collected by C. B. Kloss and presented by Kloss and H. C. Robinson. This is the author's selected male. His selected female syntype is also in the collection.

- koliensis—Serinus capistratus koliensis C. H. B. Grant, 1952. . . 59

 Bull. Brit. Orn. Cl., 72: 1. HOLOTYPE, Adult male. Reg. no. 1923.8.7.2375. Onyulu's, Koli River, Lango, Uganda, 13 March 1910. Collected by T. V. Fox. Stephenson Clarke Collection.
- konyboensis—Sitta europaea konyboensis Kinnear, 1940 . . . 43

 Bull. Brit. Orn. Cl., 60: 57. Holotype, Adult male. Reg. no. 1938.12.13.272. Molo, 11,000', south-east Tibet, 15 April 1938. Collected and presented by F. Ludlow and G. Sherrif. (cont.)

The subspecific name was corrected to kongboensis on p. 74, op. cit.

- kulalensis—Zosterops pallida kulalensis Williams, 1948. 47 Bull. Brit. Orn. Cl., 68: 101. Holotype, Adult male. Reg. no. 1948.1.3. Mount Kulal, Northern Kenya, (20°35′ N., 36°55′E., 6,000′, on label), 8 Oct. 1947. Collected by J. G. Williams. Presented by the Coryndon Museum, Nairobi.
- kumaiensis—Leiothrix lutea kumaiensis Whistler, 1943 32B Bull. Br. Orn. Cl., 63:62. HOLOTYPE, Adult male. Reg. no. 1886.10.1.6902. Dehra Dun, United Provinces, India, 28 Dec. 1870. Collected by G. King. Hume Collection.
- kumbaensis—Dyaphorophyia ansorgei kumbaensis Serle, 1949 . 38 Bull. Brit. Orn. Cl., 69:75. HOLOTYPE, Adult female. Reg. no. 1949.14.9. Kumba, 700', 4°40' N., 9°25' E., British Cameroons, 24 Nov. 1947. Collected and presented by W. Serle.

- kumboensis—Alseonax murinus kumboensis Bannerman, 1921. . . 37 Bull. Brit. Orn. Cl., 42:131. Holotype, Adult male. Reg. no. 1922.11.25.105. Near Kumbo, 6,000', Nigerian Cameroon Highlands, 3 Oct. 1921. Collected by and purchased of G. L. Bates.
- kumboensis—Hirundo rufula kumboensis Bannerman, 1923 . . . 18

 Bull. Br. Orn. Cl., 43: 142. Holotype, Adult male. Reg. no.
 1922.11.25.80. Kumbo, 5,500', Northern Nigeria, 29 Sept. 1921.
 Collected by and purchased of G. L. Bates.
- kungwensis—Alethe poliocephala kungwensis Moreau, 1941 . . . 31 Bull. Br. Orn. Cl., 61:46. HOLOTYPE, Adult male. Reg. no. 1945.34.220. Forest above Ujamba, Kungwe Mountain, 6,900', western Tanganyika, 13 Aug. 1940. Collected by Salimu Asmani for R. E. Moreau and presented by the latter.
- kungwensis—Arizelocichla tephrolaema kungwensis Moreau, 1941. 21 Bull. Br. Orn. Cl., 61:60. HOLOTYPE, Adult "male?, female". Reg. no. 1940.11.1.3. Ujamba Forest, 6,900', Kungwe Mountain, Kigoma District, western Tanganyika, 8 Aug. 1940. Collected by Salimu Asmani and presented by R. E. Moreau.
- kungwensis—Bradypterus alfredi kungwensis Moreau, 1942 . . . 35 Bull. Brit. Orn. Cl., 62: 42. HOLOTYPE, Adult male. Reg. no. 1944.8.1.4. Ujamba, 7,600', Kungwe Mountain, Western Tanganyika, August 1941. Collected for, and presented by R. E. Moreau.
- kungwensis—Cossypha polioptera kungwensis Moreau, 1941 . . . 31 Bull. Br. Orn. Cl., 61:60. HOLOTYPE, Adult male. Reg. no. 1940.11.1.2. Ujamba Forest, 7,900', Kungwe Mountain, Kigoma district, western Tanganyika, 8 Aug. 1940. Collected by Salimu Asmani for R. E. Moreau and presented by the latter.
- kungwensis—Diaphorophyia ansorgei kungwensis Moreau, 1941. 38
 Bull. Brit. Orn. Cl., 61:25. HOLOTYPE, Adult male. Reg. no.
 1945.34.42. Ujamba, 7,000′, Kungwe Mountain, East shore of
 Lake Tanganyika, 4 Aug. 1940. Collected by Salimu Asmani for
 R. E. Moreau and presented by the latter.
- kungwensis—Dryoscopus angolensis kungwensis Moreau, 1941. 23
 Bull. Br. Orn. Cl., 61:45. HOLOTYPE, Adult male. Reg. no. 1945.34.378. Nganja, 3,000′, Kungwe Mountain, western Tanganyika, 27 Aug. 1940. Collected by Salimu Asmani for R. E. Moreau and presented by the latter.
- kungwensis—Phyllastrephus flavostriatus kungwensis Moreau, 1941 21 Bull. Br. Orn. Cl., 62:29. HOLOTYPE, Adult male. Reg. no.

1940.11.1.4. Forest above Ujambe, 6,800', Kungwe Mountain, western Tanganyika, 17 Nov. 1940. Collected by Salimu Asmani and presented by R. E. Moreau.
kupeensis—Chlorophoneus kupeensis Serle, 1951 23 Bull. Br. Orn. Cl., 71:41. HOLOTYPE, Adult female. Reg. no. 1951.34.1. Kupé Mountains, 4°45′ N., 9°40′ E., Kumba Division, British Cameroons, 2 Nov. 1949. Collected by G. Nkwocha for Dr. W. Serle and presented by the latter.
kurdistanica—Sitta neumayer kurdistanica Ticehurst, 1923
kurilensis—Pyrrlula kurilensis Sharpe, 1886
kurrae—Mirafra africana kurrae Lynes, 1923
lactea—Polioptila lactea Sharpe, 1885
ladacensis—Acanthis flavirostris ladacensis R. and A. Meinertzhagen, 1926
laemosticta—Myrmeciza laemosticta Salvin, 1864
laemostictus—Dicrurus laemostictus P. L. Sclater, 1877
laeta—Cryptolopha laeta Sharpe, 1902

Novit. Zool., 2:8. SYNTYPE, Adult male. Reg. no. 1896.10.6.334. Carjabamba, 9,000', northern Peru, Jan. 1894. Collected by O. T. Baron. Salvin-Godman Collection. There are several other syntypes in the collection.
laetior—Melithreptes laetior Gould, 1875
laetior—Oriolus laetior Sharpe, 1897. Bull. Brit. Orn. Cl., 7:17. SYNTYPE, Adult. Reg. no. 1873.9.10.5. Danger River, Gaboon, West Africa. Collected and presented by H. Ansell. There is a second syntype in the collection.
laetissimus—Andropadus laetissimus Sharpe, 1989 21 Bull. Br. Orn. Cl., 10:27. HOLOTYPE, Adult male. Reg. no. 1900.6.22.1. Nandi, Kenya. Collected and presented by F. J. Jackson.
laetissimus—Carpodacus laetissimus Walton, 1905 59 Bull. Brit. Orn. Cl., 15:93. SYNTYPE, Adult female. Reg. no. 1905.12.31.254. Lhasa, Tibet, 21 Sept. 1904. Collected and presented by H. J. Walton. The male syntype is also in the collection.
Proc. zool. Soc. Lond., 1884: 54, pl. 5. HOLOTYPE, Adult. Reg. no. 1884.1.15.8. Ashantee, West Africa (restricted to Kumasi by Hall, Moreau & Galbraith, <i>Ibis</i> , 108 (1966): 170). Collected and presented by Sir Geoffrey Lagden.
Illustrations of the Zoology of South Africa, 2 (Aves): pl. 87, fig. 1. SYNTYPE, Adult male (nearly attained full summer plumage, vide Shelley, Birds of Africa, 3 (1902): 34), (relaxed mount). Reg. no. 1845.7.6.213. Between Berg and Orange Rivers. Collected by and purchased of Sir A. Smith. There is another syntype in the collection; both were listed as types by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 524. A third syntype is at the Academy of Natural Sciences, Philadelphia.
Ahtora—Collurio lahtora Sykes, 1832

- Trans. Linn. Soc. Lond., 15: 221. SYNTYPE, Adult male (relaxed mount). Reg. no. 1863.7.7.22a. Australia (restricted to New South Wales by Gould, Synopsis of the Birds of Australia, (1837). A recent label is inscribed with the locality Port Jackson and collector J. Brown, but no authority can be found for this). Presented as the type by the Linnean Society of London.
- Reg. no. 1846.9.9.120. Bolivia. Collected by and purchased of T. Bridges.
- lanaiensis—Hemignathus lanaiensis Rothschild, 1893 55 Bull. Brit. Orn. Cl., 1:24. SYNTYPE, Adult female. Reg. no. 1939.12.9.55. Lanai Island, Sandwich Islands, 15 Nov. 1892. Collected by H. C. Palmer. Rothschild Bequest. A male syntype is in the American Museum of Natural History,

vide, Hartert, Novit. Zool., 26 (1919): 170.

- landanae—Lagonosticta landanae Sharpe, 1890 . Cat. Bds. Brit. Mus., 13: 283, pl. 12. SYNTYPE, Adult male. Reg. no. 1889.7.20.336. Landana, Angola, Feb. 1883. Collected by L. Petit. Shelley Collection. The female syntype is also in the collection.
- langbianis—Muscicapula melanoleuca langbianis Kloss, 1927 Bull. Brit. Orn. Cl., 47:145. HOLOTYPE, Adult female. Reg. no. 1927.6.1.1. Arbre Broyé, Langbian Massif, South Annam, 14 May 1918. Collected and presented by C. B. Kloss,
- laotianus—Alcippe nipalensis laotianus Delacour, 1926. Bull. Br. Orn. Cl., 47:19. SYNTYPE, Adult male. Reg. no. 1926.9. 8.24. Xieng-Khouang, Laos, 28 Dec. 1925 (not 20 Dec. as in text). Collected by J. Delacour on the Second Franco-British Expedition to Indo-China.

This is the author's selected male. His selected female syntype is

also in the collection.

- laotainus—Pomatorhinus tickelli laotianus Delacour, 1926 Bull. Brit. Orn. Cl., 47:16. SYNTYPE, Adult male. Reg. no. 1926. 9.8.17. Xieng-Khouang, Laos, 3 Jan. 1926. Collected by J. Delacour on the Second Franco-British Expedition to Indo-China. This is the author's selected male. His selected female syntype is also in the collection.
- laotianus—Spizixus canifrons laotianus Delacour, 1926. Bull. Br. Orn. Cl., 47:14. SYNTYPE, Adult male. Reg. no. 1926. 9.8.15. Noug-Het, Laos, 13 Dec. 1925. Collected by J. Delacour on the Second Franco-British Expedition to Indo-China. The female syntype is also in the collection.

- laotianus—Turdinulus epilepidotus laotianus Delacour, 1926 . . . 32B Bull. Br. Orn. Cl., 47:17. HOLOTYPE, Adult male. Reg. no. 1926.9.8.21. Xieng-Khouang, Laos, 8 Jan. 1926. Collected by J. Delacour on the Second Franco-British Expedition to Indo-China.

The specimen is an adult male but has lost the crest. Ticehurst (op. cit.) refers to this as the type.

- Proc. zool. Soc. Lond., 1869: 151. SYNTYPE, Adult male. Reg. no. 1885.6.12.1052. Tinta, 11,000', Peru, 19 June 1868. Collected by H. Whitely. Sclater Collection.

 Listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 292. There are two other syntypes in the collection.

Of two specimens described, this specimen is referred to by its label number as the type. Hellmayr and Conover, *Catalogue of birds of the Americas*, 8 (1935): 125, claim that the type is in the U.S. National Museum but Deignan, *Type specimens of birds in the U.S. National Museum* (1961) does not list it.

Proc. zool. Soc. Lond., 1844: 100. SYNTYPE, Adult female. Reg.

no. 1881.2.18.136. Fernando Po, June. Collected by L. Fraser. Eyton Collection; purchased of E. Gerrard.
Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 108.

One of the immature syntypes is also in the collection.

This is certainly the holotype, though Jackson's type-label is attached to a specimen of *Anthus pyrrhonotus* in the Tervuren Museum, Belgium, *vide* correspondence between B. P. Hall and H. Schouteden, 1959, in the B.M.(N.H.).

- latouchei—Tephrodornis gularis latouchei Kinnear, 1925 . . . 20 Bull. Br. Orn. Cl., 45:105. HOLOTYPE, Adult male. Reg. no. 1905.12.24.368. Ah Ch'ung, Fohkien, south China, 2 March 1900. Collected and presented by C. B. Rickett.

- laurentei—Cryptolopha castaneiceps laurentei La Touche, 1921 . 35

 Bull. Brit. Orn. Cl., 42:53. SYNTYPE, Adult male. Reg. no.
 1923.3.10.7. Mengtsz, South-east Yunnan, 21 March 1921.
 Collected and presented as a syntype by J. D. La Touche.

 There is another syntype at Harvard, vide Bangs, Bull. Mus.

Comp. Zool., Harvard, 70 (1930): 341.

lavaudeni—Erithacus rubecula lavaudeni Bannerman, 1926 31 Bull. Brit. Orn. Cl., 47: 24. HOLOTYPE, Adult male. Reg. no. 1925.12.25.4. Les Sources, 2,200', near Ain Draham, north Tunisia, 10 March 1925. Collected by W. P. Lowe and D. Bannerman and presented by A. S. Vernay.
lavendulae—Cisticola lavendulae Ogilvie-Grant and Reid, 1901 . 35 Ibis, (8) 1:650. SYNTYPE, Adult male. Reg. no. 1902.1.20.115. Aroharlaise, Somaliland, 3 April 1901. Collected and presented by A. E. Pease. This is the male syntype. The described female syntype is also in the collection.
lawrencii—Haemophila lawrencii Salvin and Godman, 1886 49 Biol. CentrAmer., Aves, 1:397. HOLOTYPE, Adult. Reg. no. 1885.12.14.1311. Juchitan, Tehuantepec, Mexico, 13 April 1870. Collected by F. Sumichrast for Boucard. Salvin-Godman Collection.
layardi—Geocichla layardi Walden, 1870
layardi—Merula layardi Seebohm, 1890
layardi—Picolaptes layardi P. L. Sclater, 1873
layardi—Pycnonotus layardi Ayres, 1879
layardi—Saxicola layardi Sharpe, 1876
lazulina—Phyllornis lazulina Swinhoe, 1870

9
leautungensis—Alauda leautungensis Swinhoe, 1861
legalleni—Cutia nipalensis legalleni Robinson & Kloss, 1919 32B Ibis, (11) 1:588, pl. 15. SYNTYPE, Adult male. Reg. no. 1919.12. 20.331. Langbian Peaks, 7,200', south Annam, 18 April 1918. Collected and presented by C. B. Kloss. This is the author's selected male. Their selected female syntype is also in the collection.
leggei—Anthus leggei Ogilvie-Grant, 1906
leggei—Hemipus picatus leggei Whistler, 1939
leggei—Kittacincla malabarica leggei Whistler, 1941 31 Ibis, (14) 5:319. HOLOTYPE, Adult female. Reg. no. 1940.12.3. 284. Uruhaga, 100–200', Ceylon, 23 Oct. 1936. Collected by E. Fernando of the Ceylon Avifaunal Survey.
leggii—Petroica leggii Sharpe, 1879
leighi—Acanthiza leighi Ogilvie-Grant, 1909
lemoneus—Icterus vulgaris lemoneus A. G. Butler, 1907 57 Avicult. Mag., (2) 5:225, pl. Holotype, Adult. Reg. no. 1916.6. 12.1. A caged bird received by Dr. A. G. Butler from Miss J. Gladstone, Dec. 1906, and presented to the collection at its death.
lemprier:—Siphia lemprieri Sharpe, 1884

Reg.

leoninus—Andropadus curvirostris leoninus Bates, 1930 Bull. Br. Orn. Cl., 51:52. HOLOTYPE, Adult male. Reg. no. 1930.12.3.7. Buedu, 1,600', near Kailahun, east Sierra Leone, 12 April 1930. Collected by and purchased of G. L. Bates. leoninus—Macrosphenus leoninus Neumann, 1908.

Bull. Brit. Orn. Cl., 23: 46. HOLOTYPE, Adult. Reg. no. 1904.6.5. 227. Rotifunk, Sierra Leone, West Africa, 1903. Collected and presented by R. Kemp. leoninus—Onychognathus fulgidus leoninus Neumann, 1920 . J. Orn., 1920: 82. HOLOTYPE, Adult male. Reg. no. 1905.1.25. 152. Bo, south-east Sierra Leone, Feb. 1904. Collected by R. Kemp. Purchased through E. Gerrard. leoninus—Zosterops senegalensis leoninus W. L. Sclater, 1927 . Bull. Brit. Orn. Cl., 47:86. HOLOTYPE, Adult male. Reg. no. 1905.1.25.45. Bo, Sierra Leone, April 1904. Collected by and purchased of R. Kemp. leontica—Prinia leontica Bates, 1930. 35 Bull. Brit. Orn. Cl., 51:51. HOLOTYPE, Adult male. Reg. no. 1930.12.3.4. Near Birwa Peak, 4,500', Kono District, Sierra Leone, 19 July 1930. Collected by and purchased of G. L. Bates. lepe—Cisticola erythrops lepe Lynes, 1930 Ibis, (12) 6, Supplement: 376. HOLOTYPE, Adult female. Reg. no. 1928.10.25.7. Lepe Mission, 5,000', Benguela Province, Angola, 29 Jan. 1927. Collected by Admiral H. Lynes and B. B. Osmaston. Presented by Admiral H. Lynes. lepida—Eminia lepida Hartlaub, 1880 35 Proc. zool. Soc. Lond., 1880: 625, pl. 60. Syntype, Adult female. Reg. no. 1895.5.1.1027. Magunga, (head of Albert Nyanza), (26 Nov. 1879, on label). Collected by Emin Bey. Shelley Collection, received by him from Hartlaub of Bremen. See also, Hartlaub, Abhand. Nat. Ver. Bremen, 7 (1881):91. There were five syntypes of which this is one. lepida—Leiothrix lepida McClelland, 1840. in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840): 162. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4479. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.

Proc. zool. Soc. Lond., 1880: 160. SYNTYPE, Adult female.

Southern Palawan, Philippines. Collected by and purchased of

There is a second syntype in the collection.

E. Lempriere as the type.

no. 1889.7.10.684. Sarayacu, east Ecuador. Collected by C. Buckley. Salvin-Godman Collection.

Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 287.

There is a male syntype in the collection.

There are other syntypes in the collection and at the Academy of Natural Sciences of Philadelphia.

Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 295. There are several other syntypes in the collection.

- leucauchen—Thamnophilus leucauchen P. L. Sclater, 1855.

 Proc. zool. Soc. Lond., 1855: 18, pl. 79. SYNTYPE, Adult male.
 Reg. no. 1889.9.20.117. Chamicuros, east Peru (error: Cayenne tradeskin, vide Hellmayr, Novit. Zool., 14 (1907): 60, and Catalogue of Birds of the Americas, 3 (1924): 53, 55). Obtained for Parzudaki. Sclater Collection.
- leucauchen—Turdus leucauchen P. L. Sclater, 1858 31

 Proc. zool. Soc. Lond., 1858 : 447. SYNTYPE, Adult male. Reg.
 no. 1886.8.2.56. Guatemala. Collected by Skinner. Sclater Collection; obtained by Sclater through Leadbetter.
- leucogaster—Euprinodes leucogaster Sharpe, 1904. 35

 Bull. Brit. Orn. Cl., 14:94. HOLOTYPE, Adult male. Reg. no. 1904.6.22.197. Mount Clarence, Fernando Po, 27 Feb. 1904. Collected by E. Seimund. Presented by Mrs P. Sladen, the Duke of Bedford, and W. Rothschild.

- leucogastra—Dendrocitta leucogastra Gould, 1833 71
 Proc. zool. Soc. Lond., 1833 : 57. HOLOTYPE, Adult. Reg. no.
 1876.1.7.21. Eastern Asia (Malabar coast, on label). Purchased of J. Gould.
- leucogastra—Ptilocichla leucogastra Davison, 1892 32B Ibis, (6) 4:100. SYNTYPE, Adult male. Reg. no. 1892.9.1.1. Pahang (ride Sharpe, Hist. Coll. Brit. Mus., 2 (1906):337). Malay Peninsula, 1 Aug. 1891. Collected and presented by W. R. Davison
- **leucogastra**—Setaria albigularis leucogastra Hachisuka, 1926 . . . 32B Bull. Brit. Orn. Cl., 47:54. HOLOTYPE, Adult male. Reg. no. 1878.5.3.15. Paku, Sarawak, Borneo, Dec. 1878. Collected by A. H. Everett and purchased of Gerrard.
- leucogastroides—Munia leucogastroides Moore, 1856 6 in Horsfield and Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 2:510. SYNTYPE, Adult. Reg. no. 1880.1.1.1940. Java. Collected by T. Horsfield. Presented by the Honourable East India Company Museum as the type. The second syntype is also in the collection.

There is another syntype in the collection.

This specimen was labelled 'type', and listed as the type by Gadow, Cat. Bds. Brit. Mus., 8 (1888): 59. There are two other

syntypes in the collection.

- leucogonys—Tyranniscus leucogonys P. L. Sclater and Salvin, 1871.

 Proc. zool. Soc. Lond., 1870 (publ. 1871): 841, pl. 53. SYNTYPE,
 Adult. Reg. no. 1888.1.13.513. Bogota, Colombia. Collected by Bell, of New York. Sclater Collection.

 The second syntype is also in the collection.
- leucolaema—Xenocichla leucolaema Sharpe, 1902 . . . 21 Bull. Br. Orn. Cl., 13:10. HOLOTYPE, Adult male. Reg. no. 1902.12.8.17. Toro Forest, Uganda, 19 March 1902. Collected and presented by F. J. Jackson.
- leucomela—Campephaga leucomela Vigors & Horsfield, 1827. . . 20
 Trans. Linn. Soc. Lond., 15: 215. Holotype, (Immature, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 106; relaxed mount with incomplete tail). Reg. no. 1863.7.7.19. Broad Sound, Queensland, 26 Oct. 1802. Collected by R. Brown and presented by the Linnean Society.
- leucomelanura—Digenea leucomelanura Hodgson, 1845 . . . 37

 Proc. zool. Soc. Lond., 1845 : 26. SYNTYPE, Adult (male). Reg.
 no. 1880.1.1.883. (the Collector's no. is 941, an error for 419).
 Nepal. Presented by the Honourable East India Company Museum.

The specimen is thought to be that referred to by Horsfield and Moore, Catalogue of the birds in the Museum of the Honourable East India Company, 1 (1854): 294.

Proc. zool. Soc. Lond., 1877: 19. SYNTYPE, Adult. Reg. no. 1888.1.13.60. Chihuata, 4,000', near Arequipa, South west Peru. (19 July 1867, on label). Collected by H. Whitely. Sclater Collection.

Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 21. There are several other syntypes in the collection.

leuconotus—Cinclus leuconotus P. L. Sclater, 1857 27

Proc. zool. Soc. Lond., 1857: 274. Syntype, Adult male. Reg. no. 1886.8.2.272. "Bogota", Colombia. Collected for Jamrach; Sclater Collection.

There is also a female syntype in the collection.

leuconotus—Malurus leuconotus Gould, 1865 30
Proc. zool. Soc. Lond., 1865: 198. In a later work (Birds of Australia, Supplement, (1867): pl. 24), the author stated that the specimen belonged to a Mrs Cranford of Devon. There would

appear to be no basis for the statement by Mathews, Austral Avian Record, 1 (1912): 165, that the type was in the British Museum Collection.	
leucophaea—Certhia leucophaea Latham, 1801	
leucophrus—Anabates leucophrus Jardine & Selby, 1830. Illustrations of Ornithology, 2, pt. 6: pl. 93. HOLOTYPE, Adult. Reg. no. 1886.6.24.464. Brazil. Collected by Dr. Such. Jardine Collection.	
leucophrys—Sylviella leucophrys Sharpe, 1891	35
leucoproctum—Trichostoma leucoproctum Tweeddale, 1877. Proc. zool. Soc. Lond., 1877: 366. Holotype, Adult female. Reg. no. 1887.12.30.1376. Muleit Range, Tenasserim, 7 Jan. 1877. Collected by O. Limborg. Tweeddale Collection.	
leucops—Dicrurus leucops Wallace, 1865	64
leucops—Digenea leucops Sharpe, 1888	37
leucopsis—Buarremon leucopsis P. L. Sclater and Salvin, 1878 Proc. zool. Soc. Lond., 1878: 439. SYNTYPE, Adult. Reg. no. 1885.6.8.433. Yauayaca, Ecuador, Dec. 1877. Collected by C. Buckley. Salvin-Godman Collection. The second syntype is also in the collection.	
leucopsis—Motacilla leucopsis Gould, 1837	19
leucopsis—Sitta leucopsis Gould, 1850	43

- Hay. Presented by the Honourable East India Company Museum, which received it from J. Gould in 1853 (N. B. Kinnear, MS. note). The description by Gould, *Proc. zool. Soc. Lond.*, 1849:113, was not published until later in 1850.

- leucoptera—Tanagra episcopus leucoptera P. L. Sclater, 1886 . 57 Cat. Bds. Brit. Mus., 11:154, 155. SYNTYPE, Adult male. Reg. no. 1885.6.12.454. Bogota, Colombia. Collected for Argent. Sclater Collection.

Sclater used a G. R. Gray MS., name, and it appears as a nomen nudum in Sclater's Catalogue of a collection of American birds, (1862): 74. There is another syntype in the collection.

- leucopygia—Crithagra leucopygia Sundevall, 1850 59
 Ofvers K. Sv. Vet.-Akad. Forh. Stockholm, 7: 127. Syntype, Adult.
 Reg. no. 1877.7.11.358. Senaar, Abyssinia, (1837, on label).
 Collected by Prof. Hedenborg. Sharpe Collection; obtained by him from Prof. Hedenborg.

- leucopygius—Basileuterus leucopygius P. L. Sclater and Salvin, 1873 54
 Nomencl. Av. Neotrop.: 156. Syntype, Adult male. Reg. no.
 1884.5.15.460. Costa Rica, 1868. Collected by Carmiol. Sclater
 Collection.

There is an immature syntype in the collection.

39

male. Reg. no. 1873.5.12.1298. Gagie Island (off western New Guinea). 1860. Collected by A. R. Wallace. Wallace Collection. There is another syntype in the collection.	
leucorrhoa—Pipra leucorrhoa P. L. Sclater, 1863	8
leucoschistus—Henicurus leucoschistus Swinhoe, 1870	31
leucosticta—Corythocichla leucosticta Sharpe, 1887 32 Proc. zool. Soc. Lond., 1887 : 438. SYNTYPE, Adult male. Reg. no. 1886.12.22.16. Larut Range, c. 4,400', Perak, Malaya, Sept.—Oct. 1886. Collected and presented by L. Wray. The female syntype is also in the collection.	
leucosticta—Cossypha leucosticta Sharpe, 1883	31
leucostictus—Dysithamnus leucostictus P. L. Sclater, 1858. Proc. zool. Soc. Lond., 1858: 66, pl. 140. SYNTYPE, Adult (female, vide Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 128). Reg. no. 1889.9.20.207. Rio Napo, Ecuador. Collected for Verreaux. Sclater Collection. There are two immature syntypes in the collection.	4
leucothorax—Lanio leucothorax Salvin, 1865 Proc. zool. Soc. Lond., 1864 (publ. 1865): 581. SYNTYPE, Adult. Reg. no. 1885.6.7.514. Tucurriqui, Costa Rica. Collected by E. Arcé. Salvin-Godman Collection. The female syntype is also in the collection.	51
leucotis—Garrulus leucotis Hume, 1874	71
leucotis—Garrulax monileger leucotis E. C. S. Baker, 1917	В

leucorhynchus—Rectes leucorhynchus G. R. Gray, 1862

leucotis—Monarcha leucotis Gould, 1850
leucura—Eopsaltria leucura Gould, 1869
leucura—Monarcha leucura G. R. Gray, 1858
leucura—Montifringilla leucura Bonaparte, 1855
leucura—Niltava leucura Tweeddale, 1877
leucurus—Agriornis leucurus Gould, 1839
levaillantii—Drymoica levaillantii Smith, 1842

- **Ihamarum**—Alauda arvensis Ihamarum R. Meinertzhagen, 1926 . 17

 Bull. Br. Orn. Cl., 46: 100. HOLOTYPE, Adult male. Reg. no. 1887.7.1.3254. Ladak, Kashmir, 1870. Collected by Dr. G. Henderson on the Lahore to Yarkand Expedition. Hume Collection.
- Bull. Br. Orn. Cl., 58:8. HOLOTYPE, Adult male. Reg. no. 1911. 11.18.87. Nanna Kru, Liberia, 8 Jan. 1911. Collected by W. P. Lowe and purchased of O. E. Janson.
- Report of the Expedition for exploring central Africa, 1834, Appendix: 45. SYNTYPE, Adult (relaxed mount). Reg. no. 1838.6.9.39. Near Kurrichane. Collected by A. Smith and purchased of the South African Museum.

Listed as type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 230.

There is also a female syntype in the collection, and a third syntype in the Liverpool Museum.

- limitans—Cyornis banyumas limitans Robinson and Kloss, 1927. 37 Bull. Brit. Orn. Cl., 48:44. HOLOTYPE, Adult male. Reg. no. 1936.4.12.2120. Tamansari, 1,400', East Java, 15 April 1916. Collected by a native collector. Presented by the Raffles Museum, Singapore.

lineata—Acanthiza lineata Gould, 1838
lineaticeps—Anabates lineaticeps P. L. Sclater, 1856
lineatocephalis—Dendrocolaptes lineatocephalis G. R. Gray & Mitchell, 1847
lineolata—Pachycephala lineolata Wallace, 1862 39 Proc. zool. Soc. Lond., 1862 : 341. SYNTYPE, Adult male (= female, vide Sharpe, Cat. Bds. Brit. Mus., 8 (1883) : 217). Reg. no. 1873. 5.12.548. Bouru Island, Moluccas, 1861. Collected by A. R. Wallace. Wallace Collection. See also Hartert, Novit. Zool., 5 (1898) : 132. A syntype from Sula is also in the collection.
lineoventer—Munia lineoventer Hodgson, 1836
lingoo—Orthotomus lingoo Sykes, 1832
Bull. Brit. Orn. Cl., 16: 104. HOLOTYPE, Adult (male, on label). Reg. no. 1906.12.21.298. Mayotte Island, Comoro Group, South Atlantic, 1 March 1906. Collected by M. J. Nicoll on the voyage of R. Y.S. Valhalla. Presented by the Earl of Crawford.
lipiniana—Estrelda lipiniana Smith, 1836
lisettae—Anthreptes hypogrammica lisettae Delacour, 1926 4 Bull. Brit. Orn. Cl., 47:22. Syntype, Adult male. Reg. no.

1926.9.8.27. Col des Nuages, Annam, 4 Dec. 1925. Collected by J. Delacour. From the Franco-British Expedition to French Indo-China.

The female syntype is also in the collection.

- litoralis—Cyornis rufigastra litoralis Stresemann, 1925 37

 Orn. Mber., 33: 50. Holotype, Adult male. Reg. no. 1887.12.30.
 1275. Puerto Princesa, Palawan, Philippines, Dec. 1877. Collected by A. H. Everett. Tweeddale Collection.

- livingstonei—Erythrocercus livingstonei Finsch and Hartlaub, 1870. 38
 Vögel Ost-Afrikas: 303. HOLOTYPE, Adult. Reg. no. 1863.12.8.
 44. Zambesi (Collected by Rev. C. Livingstone, vide British Museum Register). Presented by the Foreign Office through the Earl of Russell.
 Finsch and Hartlaub used a G. R. Gray MS. name.

loanda—Cisticola brachyptera loanda Lynes, 1930 35. Ibis, (12) 6, Supplement: 486, pl. 16. HOLOTYPE, Adult male. Reg. no. 1928.10.25.6. Lepe, 4,900′, Benguela District, Angola, 28 Jan. 1927. Collected by Admiral H. Lynes and B. B. Osmaston. Presented by Admiral H. Lynes.	5
loandae—Eliminia longicauda loandae W. L. Sclater and Mackworth-Praed, 1918	8
Iobito—Cisticola aridula lobito Lynes, 1930	5
locustella—Paludipasser locustella Neave, 1909 60 Bull. Brit. Orn. Cl., 25:25. HOLOTYPE, Adult male. Reg. no. 1909.11.21.338. (Upper Luansenshi River, 4,000', vide Neave, Ibis, (9) 4 (1910):251), North-east of Bangeolo, Northern Rhodesia, 1 June 1908. Collected by and purchased of S. A. Neave.	0
lomaensis—Dyaphorophyia ansorgei lomaensis Serle, 1948 3 Bull. Brit. Orn. Cl., 66: 73. HOLOTYPE, Adult female. Reg. no. 1946.27.1. Bintimani Peak, 3,000', Loma Mountains, Sierra Leone, 15 April 1943. Collected and presented by W. Serle.	8
lombokia—Meliphaga lombokia lombokia Mathews, 1926 4 Bull. Brit. Orn. Cl., 46:60. New name for Ptilotis virescens Wallace, 1864, (q.v.) on the grounds of its preoccupation by Melithreptus virescens Vieillot, 1817.	8
longicauda—Dendrocincla longicauda Pelzeln, 1868 Zur Ornithologie Brasiliens: 42, 60. SYNTYPE, Adult male. Reg. no. 1889.5.20.626. Barra do Rio Negro, Brazil, 4 May 1833. Collected by J. Natterer. P. L. Sclater Collection. Listed as "authentic specimen" by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 165. This specimen conforms to the description; see Hellmayr, Catalogue of Birds of the Americas, 4 (1925): 363, concerning the syntypes in the Vienna Museum.	2
longicauda—Rhipidura longicauda Wallace, 1865	A
longicauda—Suthora longicauda Campbell, 1892	3

- longicaudata—Pnoepvga longicauda Moore, 1854 in Horsfield & Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 1:398. HOLOTYPE, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 263). Reg. no. 1880.1. 1.2582 Khasia Hills, Assam (vide Sharpe, loc, cit.). Collected by Griffiths and presented by the Honourable East India Company Museum. longicaudatus—Ammodramus longicaudatus Gould, 1839 Zoology of the vovage of H.M.S. Beagle, 3, Birds: 90, pl. 29. Syn-Type, Adult (relaxed mount). Reg. no. 1856.3.15.9. Maldonado Uraguay. Collected and presented by C. Darwin. A second syntype from Montevideo is listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 767. longicaudatus—Lanius nigriceps longicaudatus Ogilvie-Grant, 1902 . 23 Novit. Zool., 9:480. SYNTYPE, Adult. Reg. no. 1881.5.1.53. Siam. Collected by Schomburgk. Gould Collection. There are four other syntypes in the collection. Lanius longicandatus Gould, Proc. zool, Soc. Lond., 1859: 151 is a nomen nudum. longipennis—Sporophila longipennis Chubb, 1921. Ann. Mag. nat. Hist., (9) 7:193. HOLOTYPE, Adult male. Reg. no. 1885.12.14.515. Mount Roraima, 3,500', British Guiana, 24 Jan. 1884. Collected by H. Whitely. Salvin-Godman Collection. longipes—Motacilla longipes Gmelin, 1789. Systema Naturae, 1, pt. 2:979. HOLOTYPE. Unsigned finished painting. New Zealand. No. 165 in Forster, Unpublished watercolour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1, (1959): 309. longirostra—Certhia longirostra Latham, 1790 Index Orn., 1:299. This species was referred to under this name by Horsfield, Trans, Linn. Soc. Lond., 13 (1821): 167, and by Raffles, Trans. Linn. Soc. Lond., 13 (1822): 299. Gadow, Cat. Bds. Brit. Mus., 9 (1884): 109 lists type specimens for both Horsfield's and Raffles' descriptions, but since the type was Latham's specimen neither of Gadow's birds has any type status. longirostris—Chrysomitris longirostris Sharpe, 1888 Cat. Bds. Brit. Mus., 12:220. SYNTYPE, Adult male. Reg. no. 1885.12.14.1219. Roraima, British Guiana, 6 Jan. 1882. Col-

lected by H. Whitely. Salvin-Godman Collection.

- longirostris—Dicrurus (Chibia) longirostris Ramsay, 1882 Proc. Linn. Soc. New South Wales, 7:300. SYNTYPE, Adult female. Reg. no. 1884.1.19.8. Wano, San Christoval, Solomon Islands. April 1882. Collected by J. Stephens. Purchased of J. T. Cockerell. Specimen label marked 'Type ♀ 2'. longirostris—Otocoris longirostris Moore, 1855 Proc. zool. Soc. Lond., 1855: 215, pl. 111. SYNTYPE, Adult male. Reg. no. 1881.5.1.2145. Neighbourhood of Agra, Kulu, Upper India. Collected by Major Hav. Gould Collection. There are four other specimens from Hay in the collection, presumably all syntypes. Moore used Gould's MS, name. longirostris—Piezorhynchus alecto longirostris Mathews, 1928 Bull. Brit. Orn. Cl., 48:93. HOLOTYPE, Adult female. Reg. no. 1883.5.30.76. Larat, Timor Laut, 5 Sept. 1882. Collected and presented by H. O. Forbes. longirostris—Ptilotis longirostris Ogilvie-Grant, 1911 48 Bull. Brit. Orn. Cl., 29:27. HOLOTYPE, Adult male. Reg. no. 1873.5.12.1037. Wamma Island, Aru Islands, Jan. 1857. Collected by A. R. Wallace. Wallace Collection. . 32B longirostris—Pyctorhis longirostris Moore, 1854. in Horsfield & Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 1:409. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4741. Nepal. Collected by B. H. Hodgson and presented by the Honourable East India Company Museum. Moore used Hodgson's MS. name. longstaffi—Malacocincla canicapillus longstaffi T. H. Harrisson &

- Bull. Br. Orn. Cl., 19:46. HOLOTYPE, Adult male. Reg. no. 1911.12.23. 1876. Near Libokwa, Welle River, north-east Belgian Congo, 13 March 1906. Collected by J. Lopes on the Alexander-Gosling Expedition. Boyd Alexander Bequest.

A female collected on 12 March, Reg. no. 1911.12.23.1875. has for

many years been labelled as the type.

lopezi—Phlexis lopezi Boyd Alexander, 1903 35 Bull. Brit. Orn. Cl., 13:48. Syntype, Adult female. Reg. no. 1911.12.23.2246. Moka, Fernando Po., 11 Dec. 1902. (14 Dec. 1902, on label.) Collected by Boyd Alexander. Boyd Alexander Bequest.
The above specimen was described and selected but a male specimen is also labelled "Type", and there are two other syntypes in the collection.
lopezi—Pytelia lopezi Boyd Alexander, 1906
loretoyacuensis—Thannophilus loretoyacuensis Bartlett, 1882 . 4 Proc. zool Soc. Lond., 1882: 374. Syntype, Adult male. Reg. no. 1885.5.19.2. Loretoyacu, east Peru, 14 Nov. 1877. Collected by J. Hauxwell and received in exchange from the Maidstone Museum.
loricata—Grallaria loricata P. L. Sclater, 1857 4 Proc. zool. Soc. Lond., 1857 : 129. Sclater referred only to material in the Paris Museum. The two specimens he later listed as types, Cat. Bds. Brit. Mus., 15 (1890) : 327, have no type-status.
loricata—Monarcha loricata Wallace, 1863
louisae—Rhynchostruthus louisae Phillips, 1897 59 Bull. Brit. Orn. Cl., 6:47. Syntype, Adult (female, on label). Reg. no. 1898.7.27.27. Goolis Mountains, Sheikh, Somaliland, 31 Jan. 1897. Collected by and purchased of E. Lort Phillips. This specimen was labelled "Type" by the author. There is another syntype in the collection.
louisiadensis—Cracticus louisiadensis Tristram, 1889 68 Ibis, (6) 1:555. HOLOTYPE, Adult. Reg. no. 1945.53.6. Sudest Island, Louisiade Islands, off New Guinea, Autumn, 1888. Collected by B. H. Thomson. Received in exchange with York Museum.
lovati—Cisticola lovati Ogilvie-Grant, 1900
lovati—Othyphantes lovati Shelley, 1905 61 The Birds of Africa, 4:457. HOLOTYPE, Adult male. Reg. no.

- 1900.1.3.309. Lekamte, south Abyssinia, 16 March 1899. Collected and presented by H. W. Blundell and Lord Lovat.
- lowei—Alethe lowei C. H. B. Grant & Mackworth-Praed, 1941 . 31 Bull. Br. Orn. Cl., 61:61. HOLOTYPE, Adult female. Reg. no. 1932.5.10.269. 8 miles south of Njombe, 6,600′, Tanganyika, 3 Dec. 1931. Collected by H. Lynes on the Lynes-Lowe Expedition.

- luangwae—Passer diffusus luangwae Benson, 1956 61
 Bull. Br. Orn. Cl., 76: 40. HOLOTYPE, Adult male. Reg. no. 1955.41.3. Mupamadzi River, Luangwa Valley, Mpika Distr., Northern Rhodesia, 12°37′ S., 32°07′ E., 21 June 1954. Collected and presented by C. W. Benson.
- luapula—Cisticola galactotes luapula Lynes, 1933 35

 Bull. Brit. Orn. Cl., 53: 169. HOLOTYPE, Adult female. Reg. no. 1933.3.13.2. Lake Bangweolo, 3,800′, North-east Rhodesia, 26

 Nov. 1930. Collected by Admiral H. Lynes and J. Vincent. Presented by Admiral H. Lynes.
- lubomirskii—Pipreola lubomirskii Taczanowski, 1879 7
 Proc. zool. Soc. Lond., 1879 : 236. Syntype, Adult male. Reg. no. 1888.1.13.1631. Tambillo, 5,800', Cajamarca dept. 6° S., Northern Peru, 22 March 1878. Collected by Stolzmann and Jelski. Sclater Collection.
- lucida—Coereba lucida P. L. Sclater and Salvin, 1859 . . . 51

 Ibis, (1) 1:14. Syntype, Adult male. Reg. no. 1854.7.31.156.

 (Guatemala, on label). Collected by G. U. Skinner. Sclater Collection, received by him from J. Gould.

 There are several syntypes in the collection.

- vide J. E. and G. R. Gray, Catalogue of the Mammals and Birds of New Guinea in the British Museum, (1858): 27). Presented by J. MacGillivray.
- lucieni—Nigrita lucieni Sharpe and Bouvier, 1878 60
 Bull. Soc. zool. France, 3:75. HOLOTYPE, Adult. Reg. no. 1878.
 2.23.57. Ungomongo, Interior of Congo, Dec. 1877. Collected by A. Lucan and L. Petit. Purchased of A. Bouvier.
- lucifer—Carpodacus rubicilloides lucifer R. Meinertzhagen, 1926 . 59 Bull. Brit. Orn. Cl., 46:83. HOLOTYPE, Adult male. Reg. no. 1922.12.31.216. Chusha, 13,000', Southern Tibet, 18 June 1921. Collected by A. R. F. Wollaston. Presented by the Royal Geographical Society, from the 1st Everest Expedition.
- ludlowi—Phylloscopus trochiloides ludlowi Whistler, 1931 . . . 35

 Bull. Brit. Orn. Cl., 52:62. Holotype, Adult male. Reg. no. 1931.12.10.1. Maran River, 5,500', (2 stages from Kishtwar), 25

 May 1931. Collected and presented by F. Ludlow.
- ludoviciae—Merula ludoviciae Lort Phillips, 1895
 Bull. Br. Orn. Cl., 4:36. SYNTYPE, Adult male. Reg. no. 1898.7.
 27.102. Darra Ass, Somaliland. Collected and presented by E. Lort Phillips.
 There is a second syntype in the collection.

- lugentoides—Saxicola lugentoides Seebohm, 1881 31
 Cat. Bds. Brit. Mus., 5:371. SYNTYPE, Adult male. Reg. no. 1898.9.1.2274. Sennaar (error, Yemen, vide W. L. Sclater, Bull. Br. Orn. Cl., 49 (1928):17), July 1839. Collected by Botta. Seebohm Bequest.

There are two other syntypes in the Paris Museum.

lugubris—Ochthodiaeta lugubris Berlepsch, 1883
luguieri—Myiagra luguieri Tristram, 1879
lumsdeni—Babax lanceolatus lumsdeni Kinnear, 1938 32E Bull. Br. Orn. Cl., 58:76. HOLOTYPE, Adult male. Reg. no. 1937.1.17.240. Le La, between Chayul Dzong and Charme, 14,000', south Tibet, 7 May 1936. Collected by F. Ludlow and G. Sherriff and presented by the Godman-Thomas Trust.
lunaris—Ianthocincla lunaris McClelland, 1840
lundae—Eremomela grisoflava lundae C. H. B. Grant and Mackworth-Praed, 1941
lunigera—Calliste lunigera P. L. Sclater, 1851 51 in Jardine, Contributions to Ornithology: 65, pl. 70. HOLOTYPE, Adult. Reg. no. 1885.6.12.382. Ecuador. Collected for Verreaux. Sclater Collection, received by him from E. Wilson.
lunulata—Pithys lunulata P. L. Sclater & Salvin, 1873
Iunulatus—Turdus lunulatus Latham, 1801
Vol. 2, no. 41 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews. Birds of Australia, 9 (1922): 316

luptoni—Paradisea apoda luptoni P. R. Lowe, 1923 70 Bull. Brit. Orn. Cl., 43:101. HOLOTYPE, Reg. no. 1923.3.1.1. New Guinea. Presented by H.M. Customs. Taken from a box seized at Newhaven, Sussex.
luridus—Planesticus luridus Bonaparte, 1854
lurio—Cisticola (emini) lurio Vincent, 1933
luscinia—Microcerculus luscinia Salvin, 1866
luteifrons—Hylophilus luteifrons P. L. Sclater, 1881
Proc. zool. Soc. Lond., 1858: 71. HOLOTYPE, Adult. Reg. no. 1888.1.13.690. Upper Napo River, Quixos, eastern Ecuador. Probably collected by Villavicencio for Verreaux. Sclater Collection.
Iuteola—Calliste luteola P. L. Sclater, 1850
Proc. zool. Soc. Lond., 1845: 30. ?SYNTYPE, Adult. Reg. no. 1845.1.9.467. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
lutescens — <i>Mixornis rubricapilla lutecens</i> Delacour, 1926 32B <i>Bull. Br. Orn. Cl.</i> , 47:18. HOLOTYPE, Adult male (in text; female on label). Reg. no. 1924.12.21.1. Bao-Ha, Tonkin, 22 Aug. 1923 (in text; 22 Oct. 1923 on label). Collected by H. Stevens and presented by the Percy Sladen Trust and Godman Exploration Fund.
 Iutleyi—Tangara lutleyi Hellmayr, 1917

Bull. Brit. Orn. Cl., 4:22. SYNTYPE, Adult male. Reg. no. 1897.5.13.175. (Manitoe, Albay District, on label), South-east Luzon, Philippines, (3 Sept. 1844, on label). Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition.

There is no indication of the number of specimens in the original description but the author refers to 'Types' in *Ibis*, (7) 1 (1895): 453. A female specimen no. 1897.5.13.163. would appear to be a syntype, but specimen no. 1896.1.17.9. appears to have been listed as a syntype

in error by Mees, Zool. Verh., Leiden, 35 (1957): 165.

Labelled as type by author. The female syntype is also in the

collection.

This is the described male. The described female syntype is also

in the collection.

- lynesi—Anthus rufulus lynesi Bannerman & Bates, 1926 . . . 19
 Ibis, (12) 2:802. HOLOTYPE, Adult male. Reg. no. 1926.8.8.1.
 East of Bamenda, 5,000′, Cameroon, 15 Feb. 1926 (in text; 1925 on label). Collected by and purchased of G. L. Bates.

lynesi—Batis orientalis lynesi C. H. B. Grant and Mackworth-Praed, 1940
lynesi—Cisticola ayresii lynesi Bates, 1926
lynesi—Galerida cristata lynesi Whistler, 1928
lynesi—Hirundo atrocaerulea lynesi C. H. B. Grant & Mackworth-Praed, 1942
lynesi—Mirafra rufa lynesi C. H. B. Grant & Mackworth-Praed, 1933 17 Bull. Br. Orn. Cl., 53: 246. HOLOTYPE, Adult male. Reg. no. 1933.6.28.1. Delami, Kordofan Province, Sudan, 13 July 1927. Collected by W. Ruttledge and presented by the Sudan Government.
lynesi—Myrmecocichla lynesi Bannerman, 1927
maal—Petrocincla maal Sykes, 1832
mabirae—Stiphrornis mabirae Jackson, 1910
macarthuri—Pogonocichla stellata macarthuri V. G. L. van Someren, 1939
macclounii—Callene macclounii Shelley, 1903

- 4.20.2. Nwenembe, 7,500', Nyasaland, Sept. 1902. Collected by McClounie and presented by A. Sharpe.
- macconnelli—Automolus turdinus macconnelli Chubb, 1919 . . . 3

 Bull. Br. Orn. Cl., 39:60. Holotype, Adult male. Reg. no. 1922.3.5.2427. Ituribisi River, British Guiana, 1909. Collected by F. V. McConnell. McConnell Collection.
- macconnelli—Elaenia flavogaster macconnelli Chubb, 1919 . . . 9

 Ann. Mag. nat. Hist., (9) 4:304. Holotype, Adult. Reg. no.
 1922.3.5.2742. Supenaam River, British Guiana. Collected by
 F. V. McConnell. McConnell Collection.
- macconnelli—Hylopezas macularia macconnelli Chubb, 1918 . 4

 Bull. Br. Orn. Cl., 38:86. Holotype, Adult male. Reg. no. 1922.3.5.2373. Ituribisi River, British Guiana, Aug. 1909. Collected by F. V. McConnell. McConnell Collection.
- macconnelli—Pachyrhamphus macconnelli, Chubb, 1920 . . . 78

 Bull. Brit. Orn. Cl., 40: 73. HOLOTYPE, Adult male. Reg. no. 1922.3.5.3176. Bonasika River, British Guiana, 1911. Collected by F. V. McConnell. McConnell Collection.
- macconnelli—Pipromorpha oleaginea macconnelli Chubb, 1919 . 9

 Ann. Mag. nat. Hist., (9) 4:303. HOLOTYPE, Adult. Reg. no. 1922.3.5.2725. Kamakabra River, British Guiana, 1911. Collected by F. V. McConnell. McConnell Collection.
- macconnelli—Piranga saira macconnelli Chubb, 1921 5.

 Ann. Mag. nat. Hist., (9) 8: 446. Holotype, Adult male. Reg. no. 1922.3.5.4330. Upper Takutu Mountains, British Guiana, 1908. Collected by F. V. McConnell. McConnell Collection.
- macconnelli—Sclerurus maxicanus macconnelli Chubb, 1919 . . . 3

 Bull. Br. Orn. Cl., 39:41. HOLOTYPE, Adult male. Reg. no. 1922.3.5.2397. Ituribisi River, British Guiana, Feb. 1907. Collected by F. V. McConnell. McConnell Collection.

Listed as the type by Hellmayr and Conover. Catalogue of hirds of the Americas, 11 (1938): 585. There are several other syntypes in the collection

- macdonaldi-Mirafra africanoides macdonaldi C. M. N. White, 1953 17 Bull. Br. Orn. Cl., 73:88. HOLOTYPE, Adult male. Reg. no. 1946.5.2826. Yavello, south Abyssinia, 3 July 1941. Collected and presented by C. W. Benson.
- macdonaldi-Sathrocercus cinnamomeus macdonaldi C. H. B. Grant and Mackworth-Praed, 1941 Bull. Brit. Orn. Cl., 61: 26. HOLOTYPE, Adult. Reg. no. 1938.5. 18.43. Gummaro Stream, 3 miles west of Goré, Wallega area, Western Abyssinia, 7 Dec. 1917. Collected by Ödön Kovác. Purchased of Effendi Johannes.
- macgillivray—Zosterops macgillivray Sharpe, 1900 Ibis, (7) 6:345. SYNTYPE, Adult male. Reg. no. 1900.1.10.43. Foreland Bay, Epi, New Hebrides, 23 May 1899. Collected and presented by A. M. Farguhar, R.N.

This specimen was labelled and registered as the type. The other

syntypes are also in the collection.

macgillivrayi—Rhipidura macgillivrayi Sharpe, 1881 Proc. zool. Soc. Lond., 1881: 789, pl. 67. SYNTYPE, Adult male. Reg. no. 1881.5.1.559. Lord Howe Island, 13 Sept. 1853. Collected by J. Macgillivray on the voyage of H.M.S. Herald. Gould Collection.

The female syntype is also in the collection.

- machika—Geocichla machika Forbes, 1883 Proc. zool. Soc. Lond., 1883: 589. HOLOTYPE, Adult male. Reg. no. 1883.5.30.101. Timor Laut, Sept. 1882. Collected by H. O. Forbes and presented by the Committee of the Timor Laut Expedition.
- mackenziana—Cryptolopha mackenziana Sharpe, 1892. Ibis, (6) 4:153. SYNTYPE, Adult male. Reg. no. 1893.12.1.20. Kikuyu, Kenya, 25 Aug. 1889. Collected and presented by F. J. Jackson.

This specimen was presented as the type but a second specimen is referred to in the description.

- mackenziei—Emberiza schoeniculus mackenziei Bird, 1935 49 Bull. Brit. Orn. Cl., 56:54. HOLOTYPE, Adult female. Reg. no. 1947.15.2. South Lochboisdale, South Uist, Outer Hebrides, Scotland, 6 April 1935. Collected and presented by E. G. Bird.
- mackinnoni—Lanius mackinnoni Sharpe, 1891 Ibis, (6) 3:444. HOLOTYPE, Adult male. Reg. no. 1893.12.1.15. (Bugemaia, op. cit.: 596), Kikuyu, Central East Africa, 14 Oct. 1889. Collected and presented by F. J. Jackson.

- macleani—Pseudogerygone macleani Ogilvie-Grant, 1907 Ibis, (9) 1:545. HOLOTYPE, Adult male. Reg. no. 1908.3.30.6. Maungahaumia, 2,000', North Island, New Zealand, 8 July 1906. Collected by and purchased of J. C. Maclean. macleavii—Ptilonorhynchus macleavii Vigors & Horsfield, 1827 Trans. Linn. Soc. Lond., 15:263. HOLOTYPE, Adult male. Reg. no. 1863.7.7.51a. Australia. Collected by Caley. Presented by the Linnean Society. macmillani—Anomalospiza macmillani Bannerman, 1911 Bull. Br. Orn. Cl., 29:37. SYNTYPE, Adult male. Reg. no. 1912. 10.15.1564. Korssa, 5,000', Jimma, south-western Abyssinia, 21 May 1905. Collected by P. C. Zaphiro. Presented by W. N. Macmillan. The female syntype is in the collection. macmillani—Estrilda macmillani Ogilvie-Grant, 1907 . Bull. Brit. Orn. Cl., 19:108. SYNTYPE, Adult male. Reg. no. 1905.12.25.117. Ibago, Bara River, Sudan, 27 March 1904. Collected by P. C. Zaphiro. Presented by W. N. McMillan. The two other syntypes are also in the collection. macmillani—Pentholaea macmillani Sharpe, 1906. Bull. Br. Orn. Cl., 16:126. HOLOTPYE, Adult female. Reg. no. 1912.10.15.830. Winke Goffa, 6,000', South Abyssinia, 27 June 1905. Collected by P. Zaphiro and presented by W. N. McMillan. macphersoni—Apalis macphersoni Vincent, 1934. Bull. Brit. Orn. Cl., 59: 177. HOLOTYPE, Adult male. Reg. no. 1934.3.4.1. Cholo Mountain, 3,500', Nyasaland, 21 July 1933. Collected and presented by D. W. Macpherson. macroptera—Myiagra macroptera Vigors and Horsfield, 1827 . 38 Trans. Linn. Soc. Lond., 15:254. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.48. Australia. Collected by Caley. Presented by the Linnean Society as the type. The specimen now lacks a tail. macropus—Locustella macropus Swinhoe, 1863 . Proc. zool. Soc. Lond., 1863:93. HOLOTYPE, Adult. Reg. no. 1898.9.1.1378. Amoy, South-east China, 31 May 1861. Collected by R. Swinhoe. Seebohm Bequest. macrorhyncha—Pratincola macrorhyncha Stoliczka, 1872 J. Asiat. Soc. Bengal, 41:238. SYNTYPE, Adult female. Reg. no. 1886.4.1.1533. Near Bhuj, Cutch, India, 20 Feb. 1872. Collected by F. Stoliczka. Hume Collection. The second syntype is also in the collection. macrorhynchus—Acrocephalus macrorhynchus Hume, 1871 .
 - macrorhynchus—Acrocephalus macrorhynchus Hume, 1871 . . . 35

 Ibis, (3) 1:31. HOLOTYPE, Adult. Reg. no. 1886.7.8.1742.

 Sutlej Valley, not far from Rampoor, Himalayas, (13 Nov. 1867, on label). Collected by A. O. Hume. Hume Collection.

- macrorhynchus—Bradypterus macrorhynchus Jackson, 1910 . . . 35 Bull. Brit. Orn. Cl., 17:8. HOLOTYPE, Adult female. Reg. no. 1910.10.8.2. Il-polossat, 7,500', Laikipia, Central Kenya, 21 June 1908. Collected and presented by F. J. Jackson.
- maculata—Malacocichla maculata P. L. Sclater, 1858 31

 Proc. zool. Soc. Lond., 1858: 64. HOLOTYPE, Adult. Reg. no. 1858.2.4.33. Rio Napo, Quixos Province, east Ecuador. Purchased of Verreaux.

 Sclater used Verreaux's MS. name.

- maculicaudis—Pyriglena maculicaudis P. L. Sclater, 1858 . . . 4
 Proc. zool. Soc. Lond., 1858: 247. Syntype, Adult male. Reg.
 no. 1889.9.20.437. Trinidad (error, imported from north Colombia,
 vide Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 223).
 Sclater Collection.

Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 268. The other syntype is in the collection.

The description by Sharpe, *Proc. zool. Soc. Lond.*, 1895 (publ. Oct.): 470 indicates that the original description was that of the female, although this was not stated in the original. The unsexed specimen collected at the same date and place, now in the Academy of Natural Sciences of Philadelphia, and listed by De Schauensee, *Proc. Acad. nat. Sci. Philadelphia*, 101 (1957): 243, as a syntype, appears to have no type status.

maculipectus—Garrulax ocellatus maculipectus Hashisuka, 1953 . 32B Auk, 70:92. New name for Ianthocincla ocellata similis Rothschild, 1921 (q.v.) on the grounds of its preoccupation by Trochalopteron simile Hume, 1871.

maculipectus—Rhipidura maculipectus G. R. Gray, 1858 . . . 37A Proc. zool. Soc. Lond., 1858: 176. SYNTYPE, Adult male. Reg. no. 1858.3.10.60. Aru Islands. Collected by A. R. Wallace. Wallace Collection.

The female syntype is also in the collection.

The author stated that his type was at the India Museum. The

wing measurement given in the text is a printer's error.

maculipennis—Coccothraustes maculipennis P. L. Sclater, 1860 . 59

Proc. zool. Soc. Lond., 1860: 251, pl. 163. SYNTYPE, Adult

(female, vide Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 35). Reg. no.

1885.2.10.645. (Vera Cruz, on label), Mexico. Collected by A.

Sallé. Sclater Collection.

Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 35. There is another syntype in the collection.

- maculipennis—Thanmophilus maculipennis P. L. Sclater, 1855

 Edinb. New Philos. Journ. (n.s.), 1:247. As this name was introduced as a 'new name' for Thanmophilus stellatus Spix it stands as an objective synonym of the latter, in spite of Sclater's intention of superceding Spix's loose description. He later listed as type a specimen from his collection, Cat. Bds. Brit. Mus., 15 (1890): 217. None of the specimens referred to by Sclater as Thannophilus maculipennis (loc. cit.) have any type-status.

This specimen has been selected rather than the syntype listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 267, since it

agrees better with the description.

- maculosus—Graucalus maculosus McClelland, 1840 20 in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840) : 159. HOLOTYPE, Adult male (relaxed mount in poor condition). Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.

maderensis—Fringilla maderensis Sharpe, 1888
Cat. Bds. Brit. Mus., 12:175. SYNTYPE, Adult male. Reg. no.
1879.4.5.724. Madeira, 20 June 1871. Collected by F. D.
Godman. Salvin-Godman Collection.
This specimen was labelled as the type by the author. There
are four other syntypes in the collection.
madierensis—Regulus madierensis Harcourt, 1851
Sketch of Madeira: 118. (also quoted in Proc. zool. Soc. Lond.,
1854: 153). SYNTYPE, Adult (relaxed mount). Reg. no. 1856.2.
4.1. Madeira. Collected by E. Vernon Harcourt.
Listed as the type by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 85,
but a second specimen was referred to in the description.
out a second specimen was referred to in the description.
modesi Mouticela selitarius madesi Chesen 1040
madoci—Monticola solitarius madoci Chasen, 1940 31
Bull. Br. Orn. Cl., 60:97. HOLOTYPE, Adult female. Reg. no.
1947.60.43. Batu Caves, near Kuala Lumpur, Selangor, 24 May
1910. Collected by 'C.B.H.H.', and presented by the Raffles
Museum, Singapore.
n · 16 · 11 · 1700
magellanica—Motacilla magellanica Gmelin, 1789 6
Systema Naturae, 1, pt. 2:979. HOLOTYPE, Unsigned, almost
finished painting. Terra del Fuego, 28 Dec. 1774. No. 163 in
Forster, Unpublished water-colour drawings made during
Cook's second voyage. Library of the Zoology Department,
British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus.
(Nat. Hist.), Hist. Ser., 1 (1959): 309.
magellanicus—Troglodytes magellanicus Gould, 1837
Proc. zool. Soc. Lond., 1836 (publ. 1837): 88. SYNTYPE, Adult. Reg.
no. 1885.3.6.408. In Fretu Magellanico. Collected by C. Darwin.
Salvin-Godman Collection; obtained by them from J. Gould.
This specimen bears the date 4 Jan. 1837, possibly when Darwin
handed it to Gould.
magna—Acanthiza magna Gould, 1855
Birds of Australia, Supplement, pt. 2: text to pl. 28. HOLOTYPE, Adult. Reg. no. 1858.12.2.35. Van Dieman's Land. Collected
Adult Reg. no. 1858 12.2.35. Van Dieman's Land. Collected
by R. Gunn. Purchased of J. Gould.
This specimen was labelled "Type and only specimen" by Gould,
but a later specimen was figured, vide Mathews, Birds of Australia,
10 (1922): 46.
10 (1922) . 40.
magna—Cinnyris magna Hodgson, 1837
India Review, 1837: 272. Syntype, Adult male (relaxed mount).
Reg. no. 1843.1.13.221. Nepal. Collected and presented by
B. H. Hodgson. The specimen is foded. There are other syntypes in the col
The specimen is faded. There are other syntypes in the collection
11.1.11.11

magna—Galerida magna Hume, 1871
magna—Sitta magna Wardlaw Ramsay 1876
magnifica—Aethopyga magnifica Sharpe, 1876
magnirostratus—Cinnyris cyanolaemus magnirostratus Bates, 1930. 46 Bull. Brit. Orn. Cl., 51:53. Holotype, Adult male. Reg. no. 1930.12.3.6. Sandaru, Kono District, Sierra Leone, 18 June 1930. Collected by and purchased of G. L. Bates.
magnirostris—Alcippe magnirostris Moore, 1854
magnirostris—Alcippe magnirostris Walden, 1875 32B in Blyth, Birds of Burma: 115 (extra number of J. Asiat. Soc. Bengal, 44, pt. 2 Aug. 1875). SYNTYPE, Adult male. Reg. no. 1888.4.20. Kare-nee (Tonghoo Hills, 25 March 1875, on label). Collected by R. G. Wardlaw Ramsay. Tweeddale Collection. Renamed Alcippe poiocephala karenni by Robinson & Kloss, on the grounds of preoccupation by A. magnirostris Moore, 1854.
magnirostris—Eopsaltria magnirostris Gould, 1869 37 Ann. Mag. nat. Hist., (4) 4:109. HOLOTYPE, Adult. (female, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883):178). Reg. no. 1881.5.1. 1111. Rockingham Bay, Queensland. Collected by E. P. Ramsay (the label bears his initials). Gould Collection. Gould used a Ramsay MS. name.
magnirostris—Geospiza magnirostris Gould, 1837
magnirostris—Myiobius magnirostris

magnirostris—Tyrannula magnirostris Gould, 1839 Zoology of the Voyage of H.M.S. Beagle, 3, Birds: 48, pl. 8. SYNTYPE, Adult (relaxed mount). Reg. no. 1856.3.15.10. Chatham Island, Galapagos Archipelago, October, (the date 4 Jan. 1837) on the label appears to refer to the date when Gould received it). Collected and presented by C. Darwin.

The generic name was changed to Myoibius by G. R. Grav when the text was edited after the publication of the plate. Listed as the

type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 263.

- magnirostris—Zosterops novaeguineae magnirostris Mees, 1955 Zool. Meded. Rijksmus. Nat. Hist. Leiden, 34:153. HOLOTYPE, Adult male, Reg. no. 1936.4.20.94. Awar, North-east New Guinea, Jan. 1936. Collected by A. Chaplin. Presented by Lord Moyne.
- magnum—Malacopteron magnum Eyton, 1839 Proc. zool. Soc. Lond., 1839: 103. HOLOTYPE, Adult male. Reg. no. 1881.2.18.121. Sumatra (on label; paper entitled 'Birds from Malaya'). Evton Collection.
- magrathi—Anorthura magrathi C. H. T. Whitehead, 1907 Bull. Br. Orn. Cl., 21:19. SYNTYPE, Immature female. Reg. no. 1908.11.10.142. Kurram Valley, 12,500', Safed Koh Range, North-West Frontier Province, India, 6 Aug. 1906. Collected and presented by C. H. T. Whitehead.

There is a second syntype in the collection.

- magrathi—Molpastes magrathi C. H. T. Whitehead, 1908 21 Bull. Br. Orn. Cl., 21:48. SYNTYPE, (Adult female on label). Reg. no. 1908.11.12.35. Bannu, 1,000', North-west Provinces, India, 20 Feb. 1907. Collected and presented by H. A. F. Magrath. There are two other syntypes in the collection.
- mahali—Plocepasser mahali Smith, 1836 Report of the Expedition for exploring Central Africa of 1834:51. SYNTYPE, Adult male. Reg. no. 1845.7.6.132. Between the Orange River and Tropic, South Africa. (Restricted to Modder River near the junction with the Riet River, Orange Free State, vide, 3rd Report of the S. A. O. S. List Committee, Ostrich 30 (1959): 116). Collected by and purchased of A. Smith.

This specimen was listed as the type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 246, but there are other syntypes in the collection

and in the Academy of Natural Sciences of Philadelphia.

mahei—Cinnyris mahéi Nicoll, 1906 46 Bull. Brit. Orn. Cl., 16:106. SYNTYPE, Adult male. Reg. no. 1906.12.21.379. Mahé Island, Seychelles, Indian Ocean, 25 March 1906. Collected by M. J. Nicoll on the voyage of R. Y.S. Valhalla. Presented by the Earl of Crawford. There are two other syntypes in the collection.

mahrattensis—Pastor mahrattensis Sykes, 1832 62 Proc. zool. Soc. Lond., 1832:95. Holotype, Adult (relaxed mount). Reg. no. 1880.1.1.4417. The Ghauts, Deccan, India. Collected by W. H. Sykes. Presented by the Honourable East India Company Museum.
mahrattensis—Turdus simillimus mahrattensis Whistler & Kinnear, 1933
Collected by S. B. Fairbank. Hume Collection.
major—Alcippe nipalensis major Delacour, 1926 32B Bull. Br. Orn. Cl., 47:18. Syntype, Adult male. Reg. no. 1926. 9.8.22. Col des Nuages, Huê, Annam, 9 Feb. 1926. Collected by J. Delacour on the Second Franco-British Expedition to Indo-China. This is the author's selected male. His selected female syntype is also in the collection.
major—Ceblepyris major Sharpe, 1870
major—Diglossa major Cabinis, 1849
major—Dumeticola major W. E. Brooks, 1871
major—Glyphorhynchus major P. L. Sclater, 1862
major—Horeites major Horsfield and Moore, 1854 35 A Catalogue of birds in the nuseum of the Honourable East India Company, 1:322. SYNTYPE, Adult, (relaxed mount). Reg. no. 1880.1.1.3057. Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company as the type specimen, but a second syntype was available to the author. Horsfield and Moore used a Hodgson MS. name. This specimen was listed as the type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 145.

- major—Hyloterpe major Bourns and Worcester, 1894 . . . 39

 Minn. Acad. Nat. Sci. Occ. Papers, 1:22. SYNTYPE, Adult male.

 Reg. no. 1896.6.6.346. Cebu, (17 March 1888, on label). Collected

 by D. C. Worcester. Steere Collection.

 There are several other syntypes in the collection.
- major—Orthotomus atrogularis major Chasen and Kloss, 1928. 35 J. Malay Branch R. Asiat. Soc., 6, pt. 3:60. Holotype, Adult male. Reg. no. 1947.60.41. Siantan Island, Anamba Group, South China Seas, 5 Sept. 1925. Collected by F. N. Chasen. Presented by the Raffles Museum, Singapore.
- major—Orthotomus major Blundell and Lovat, 1899 . . . 35 Bull. Brit. Orn. Cl., 10: 20. HOLOTYPE, Adult male. Reg. no. 1900.1.3.111. Getemma, Abyssinia, 14 March 1899. Collected and presented by H. W. Blundell and Lord Lovat.
- major—Rhinocichla mitrata major Robinson & Kloss, 1919 . . . 32B Bull. Br. Orn. Cl., 40:16. Holotype, Adult male. Reg. no. 1936.4.12.28. Gunong Ijan, 4,000′, Larut Hills, Perak, Malaya, 18 April 1915. Collected by H. C. Robinson and C. B. Kloss and presented by the Raffles Museum, Singapore.
- major—Schoeniparus rufigularis major E. C. S. Baker, 1920 . 32B Bull. Br. Orn. Cl., 41:11. HOLOTYPE, Adult male. Reg. no. 1920.10.27.2. Pak Mat, Camp 18, French Laos, 19 Jan. 1920. Collected and presented by E. J. Herbert.
- makalaka—Erythropygia makalaka Neumann, 1920 31 J. Orn., 68:83. The author stated that the holotype, collected by Bradshaw, was in the collection but it cannot be traced either in the collection or in the register of Bradshaw's specimens.

(G. R. Gray).

- makayii—Sylvietta ruficapilla makayii C. M. N. White, 1953. . . 35 Bull. Brit. Orn. Cl., 73:69. HOLOTYPE, Adult male. Reg. no. 1910.5.6.897. Malandje, Northern Angola, 18 Feb. 1909. Collected by and purchased of W. J. Ansorge.
- malabaroides—Chibia malabaroides Hodgson, 1837 . . . 64
 The Indian Review, 1 (1836, publ. 1837): 325. ?Syntype, Adult.
 Reg. no. 1843.1.13.706. Nepal. Collected and presented by
 B. H. Hodgson. (cont.)

Received with his main collection, the specimen has Hodgson's drawing number on the label and is therefore likely to have been before him when the description was made.

- malayana—Chlorura hyperythra malayana Robinson, 1928 . . . 60
 Bull. Brit. Orn. Cl., 48: 72. Holotype, Adult female. Reg. no.
 1905.2.1.267. Telôm (Cameron Highlands), 3,500', PerakPehang Border, Malaya, 22 Jan. 1902. Collected by H. C.
 Robinson. Presented by the Royal Society and the Universities of
 Edinburgh and Liverpool.
- malayana—Musicicapula malayana Ogilvie-Grant, 1906 . . . 37

 Bull. Brit. Orn. Cl., 19:10. SYNTYPE, Adult female. Reg. no. 1906.7.23.269. Gunong Tahan, 400–500′, north of Pahang, Malaya, 12 June 1905. Collected by H. C. Robinson. Presented by the Selangore State Museum.

The described male syntype is also in the collection.

- malayanus—Oriolus cruentus malayanus Robinson and Kloss, 1923. 63

 J. Federated Malay States Museum, 11:56. Holotype, Adult male.
 Reg. no. 1936.4.12.143. Semangko Pass, 2,500–4,000′, Selangor-Pahang Boundary, Malaya, 26 Feb. 1908. Collected by H. C. Robinson and C. B. Kloss. Presented by the Raffles Museum, Singapore.
- malayanus—Phyllergates cucullatus malayanus Chasen, 1938 . 3:
 Orn. Mber., 46: 7. HOLOTYPE, Adult male. Reg. no. 1948.23.2.
 Semangko Pass, 2,500–4,500′, Selangor-Pahang Boundary, Malaya, 24 Feb. 1908. Collected by F. N. Chasen. Presented by the Raffles Museum, Singapore.

The male syntype is also in the collection.

malayensis—Pastor malayensis Eyton, 1839
malayorum—Parus major malayorum Robinson and Kloss, 1918. 42 J. Fed. Malay St. Mus., 8: 226. Syntype, Adult male. Reg. no. 1920.6.29.509. Sungi Kumbang, 4,500′, Korinchi, West Sumatra, 1 April 1914. Collected and presented by H. C. Robinson and C. B. Kloss.
This is the authors' selected male, the selected female syntype is also in the collection.
malcolmi—Timalia malcolmi Sykes, 1832
malcolmsmithi—Cryptolopha malcolmsmithi Robinson and Kloss, 1919
malopensis—Spiloptila malopensis Sharpe, 1903 35 Bull. Brit. Orn. Cl., 13:80. SYNTYPE, Adult male. Reg. no. 1900.2.4.169. Malopo River, Monysui's Country, Bechuanaland, (13 Jan. 1881, on label). Collected by J. S. Jameson. Presented by Mrs Jameson. There is a second syntype in the collection.
manaiensis—Zosterops maderaspatana menaiensis Benson, 1969. 47 Bull. Brit. Orn. Cl., 89: 25. HOLOTYPE, Adult, ? male. Reg. no. 1952.60.28. Menai Island, Cosmoledo Atoll, Western Indian Ocean, 15 Feb. 1952. Collected by D. Vesey Fitzgerald. Presented by J. G. Williams, Coryndon Museum, Nairobi.
mandanus—Telephonus jamesi mandanus Neumann, 1903 23 Orn. Mber., 11:183. Holotype, Adult female. Reg. no. 1887.11. 3.110. Manda Island, East Africa, 30 Nov. 1885. Collected and presented by F. J. Jackson.
mandelli—Dumeticola mandelli Brooks, 1875

mandelli—Minla mandelli Godwin-Austen, 1876 321 Ann. Mag. nat. Hist., (4) 17:33. HOLOTYPE, Adult male. Reg. no. 1895.7.14.1285. Naga Hills forests, about 6,000′, north-east Bengal. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.
mandelli—Montifringilla mandelli Hume, 1876 65 Stray Feathers, 4:488. Syntype, Adult. Reg. no. 1887.6.1.370. Tibet, Sept. 1874. Collected for L. Mandelli. Hume Collection. The other syntype in the collection.
mandelli—Procarduelis mandelli Hume, 1873
mandellii—Certhia mandellii Brooks, 1874
mandellii—Cyornis mandellii Hume, 1874
mandellii—Reguloides mandellii Brooks, 1879 35 Stray Feathers, 8: 389. Syntype, Adult male. Reg. no. 1898.9. 1.897. (Darjeeling, Sikkim, 25 Oct. 1879, on label). Collected by W. E. Brooks. Seebohm Collection. This specimen is no. 1 on the list on page 390 (op. cit.) and was labelled "type" by the author. There are several other syntypes in the collection.
manengubae—Bradypterus mariae manengubae Serle, 1949 35 Bull. Brit. Orn. Cl., 69: 55. HOLOTYPE, Adult female. Reg. no. 1949.14.8. Manengabu Mountain, 6,500′, Cameroons, 5°5′ N., 9°50′ E., 19 March 1948. Collected and presented by W. Serle.
manengubae—Poliolais lopesi manengubae Serle, 1949 35 Bull. Brit. Orn. Cl., 69: 74. HOLOTYPE, Adult female. Reg. no. 1949.14.11. Manenguba Mountain, 6,000′, Kumba Division, Cameroons, 25 March 1948. Collected and presented by W. Serle.
manimbe—Ammodramus manimbe G. R. Gray, 1839

- manipurensis—Proparus manipurensis Ogilvie-Grant, 1906 . . . 32B Bull. Br. Orn. Cl., 16:123. Syntype, Adult. Reg. no. 1895.7. 14.1948. Owenkulno Peak, 8,000′, Manipur Hills. Collected by H. H. Godwin-Austen. Godwin-Austen Collection. There are three other syntypes in the collection.
- manningi—Malaconotus manningi Shelley, 1899 23
 Bull. Br. Orn. Cl., 8:35. HOLOTYPE, Adult male. Reg. no. 1899.3.1.94. Nyasaland (may be presumed to be from Mambwe Dist., Tanganyika, between July-Sept., vide Shelley, Ibis, (7) 5 (1899) : 364, 370). Collected and presented by W. Manning.
- marchanti—Illadopsis cleaveri marchanti Serle, 1956 32B Bull. Br. Orn. Cl., 76:22. HOLOTYPE, Adult male. Reg. no. 1955.59.1. Omanelu, Rivers Province, 100', Nigeria, 5 Aug. 1953. Collected and presented by W. Serle.
- margaritae—Sporaeginthus margaritae Blundell and Lovat, 1899 . 60 Bull. Brit. Orn. Cl., 10:20. HOLOTYPE, Adult male. Reg. no. 1900.1.3.244. Gelongol, Abyssinia, 13 March 1899. Collected and presented by Lord Lovat.
- margaritata—Myrmeciza margaritata P. L. Sclater, 1855 . . . 4

 Proc. zool. Soc. Lond., 1854 (publ. 1855): 253, pl. 71. SYNTYPE,

 Adult male. Reg. no. 1889.9.20.182. Chamicuros, Peru, 1854.

 Collected for Verreaux. Sclater Collection.

 There is a female syntype in the collection.

330 marginalis
marginalis—Amadina marginalis Sharpe, 1890 60 Cat. Bds. Brit. Mus., 13:290. SYNTYPE, Adult male (relaxed mount). Reg. no. 1838.1.27.3. West Africa. Presented by J. Pulham. Listed as the type by Shelley, Birds of Africa, 4 (1905): 125, but a second syntype is also referred to in the original description.
marginata—Campephaga marginata Wallace, 1863 20 Proc. zool Soc. Lond., 1863 : 31. SYNTYPE, Adult male. Reg. no. 1873.5.12.276. Bouru, Moluccas. Collected by A. R. Wallace. Wallace Collection. There are two other syntypes in the collection.
marginata—Heterocnemis marginata P. L. Sclater, 1855 28 Proc. zool. Soc. Lond., 1855: 145. SYNTYPE, Adult. Reg. no. 1886.9.15.83. East Peru. Collected for Verreaux. Sclater Collection. There is another syntype in the collection.
marginata—Mirafra marginata Hawker, 1898
marginatum—Cinclosoma marginatum Sharpe, 1883 32A Cat. Bds. Brit. Mus., 7:336. SYNTYPE, Adult male (in poor condition). Reg. no. 1856.12.24. North-west Australia (restricted to Mount Kenneth, Western Australia, 17 Aug. 1854, by Condon, Rec. S. Aust. Mus., 14 (1962):364). Collected by the Robert Austin Expedition. The second syntype is also in the collection. Sharpe, Cat. Bds. Brit. Mus., 7 (1883):336 incorrectly stated the two specimens to have been collected by J. R. Elsey.
marginatus—Cinnyris marginatus Ogilvie-Grant, 1907 46 Bull. Brit. Orn. Cl., 19:106. HOLOTYPE, Adult male. Reg. no. 1907.6.26.94. Upper Congo, 2,000′, 8 Feb. 1907. Collected by and purchased of D. Carruthers.
marginatus—Tropidorhynchus marginatus G. R. Gray, 1862. 48 Proc. zool. Soc. Lond., 1861 (publ. 1862): 429. SYNTYPE, Adult male. Reg. no. 1873.5.12.1014. Waigiou, off New Guinea. Collected by A. R. Wallace. Wallace Collection. There are two other syntypes in the collection.
mariae—Nessillas mariae Benson, 1960

Bandamale, 500 metres, Moheli, Comoro Islands, 18 Sept. 1958. Collected by C. W. Benson. Presented by the B. O. U. Centenary Expedition.

mariae—Pomatorhinus mariae Walden, 1875
Reg. no. 1888.4.20.724. Tonghoo Hills, Tenasserim, Burma, April 1874. Collected by R. G. W. Ramsay. Tweeddale Collection.
mariae—Urolais mariae Boyd Alexander, 1903
marila—Hydrobata marila Swinhoe, 1859
History, vide Hartert, Novit. zool., 27 (1920): 504. Both specimens were marked as types by Swinhoe.
marinae—Myiagra caledonica marinae Salomonsen, 1934 38 J. Orn., 82:437. HOLOTYPE, Adult male. Reg. no. 1926.12.18. 62. (Hog Harbour, on label), Santo Island, New Hebrides, 19 July 1925. Collected and presented by P. A. Buxton.
marionae—Mirafra assamica marionae E. C. S. Baker, 1915 17 Bull. Br. Orn. Cl., 36: 34. Syntype, Adult male. Reg. no. 1915. 12.18.157. Ayuthia, central Siam, 8 July 1914. Collected by M. Williamson and presented by J. F. Williamson. The female syntype is also in the collection.
mariquensis—Bradornis mariquensis Smith, 1847
mariquensis—Ploceus mariquensis Smith, 1845
maritima—Agriornis maritima G. R. Gray

marmorata—Ptilotis marmorata Sharpe, 1882
marmoratus—Turdinus marmoratus Wardlaw-Ramsay, 1880 32B Proc. zool. Soc. Lond., 1880 : 15. HOLOTYPE, Adult. Reg. no. 1888.4.20.1032. Sumatra. Collected by C. Bock. Tweeddale Collection.
maroantsetrae—Oxylabes zosterops maroantsetrae Salomonsen, 1934 21 Ann. Mag. nat. Hist., (10) 14:69. Holotype, Adult male (bill broken). Reg. no. 1931.8.18.2048. Maroantsetra, (on coast), north-east Madagascar, 15 July 1930. Collected by A. L. Rand and Du Mont on the Archbold-Vernay Expedition to Madagascar.
marrineri—Myiomoira macrocephala marrineri Mathews and Iredale, 1913
marshalli—Serinus marshalli Shelley, 1902
martini—Campophaga martini Jackson, 1912
martinicensis—Thryothorus martinicensis P. L. Sclater, 1866
mashona—Cisticola lais mashona Lynes, 1930
masukuensis—Andropadus masukuensis Shelley, 1897 21 Ibis, (7) 3:534. SYNTYPE, Adult male. Reg. no. 1897.11.4.199. Kombi, 7,000', Masuka Range, north Nyasa, July 1896. Collected by A. Whyte and presented by Sir H. H. Johnston. The two other syntypes are also in the collection.

masukuensis—Parus masukuensis Shellev, 1900 Birds of Africa, 2:238. SYNTYPE, Adult male. Reg. no. 1897.11. 4.121. Masuka Range, Nyasaland, July 1896. Collected by A. Whyte. Presented by Sir H. H. Johnston. The second syntype is also in the collection. matucanensis—Buarremon matucanensis Brabourne and Chubb, 1914 49 Bull. Brit. Orn. Cl., 35:20. HOLOTYPE, Adult female. Reg. no. 1914.12.2.263. Matucana, Peru, 15 June 1914. Collected and presented by Lord Brabourne and C. Chubb. mauritanica—Cotile mauritanica Meade-Waldo, 1901 . 18 Bull. Brit. Orn. Cl., 12:27. SYNTYPE, Adult male. Reg. no. 1904.4.12.20. Wad Moorbei, Rehamra, Morocco, 5 June 1901. Collected and presented by E. G. B. Meade-Waldo. This is the author's selected syntype. There are two other syntypes in the collection. mayornata—Aplonis mayornata Buller, 1888 Birds of New Zealand (2nd edn.), 1:25. HOLOTYPE, Adult (relaxed mount). Reg. no. Old Vel. Cat. no. 12. 192. ?Raiatea, Society Islands. Sharpe Cat. Bds. Brit. Mus., 13 (1890): 135, used the spelling A. inornata, regarding the above name as an error. maxima—Hirundo puella maxima Bannerman, 1922 18 Bull. Br. Orn. Cl., 43:75. HOLOTYPE, Adult male. Reg. no. 1922.11.25.31. Kumbo, 5,500', North Nigerian highlands, 28 Sept. 1921. Collected by and purchased of G. L. Bates. maxima—Merula maxima Seebohm, 1881 31 Cat. Bds. Brit. Mus., 5:405. SYNTYPE, Adult male. Reg. no. 1888.2.20.583. Gulmerg, Kashmir, Aug. 1867. Collected by Dr. Jerdon. Tweeddale Collection. maxima—Mviolestes maxima Layard, 1876 38 Ibis, (3) 6:498. HOLOTYPE, Adult male. (= female, vide Mayr, Amer. Mus. Novit., 628 (1933):16, 19). Reg. no. 1897 6.1.613. Kandavu, Fiji Islands. Collected for E. L. Layard. Seebohm Collection. maxwelli—Melanopteryx maxwelli Boyd Alexander, 1903 61 Bull. Br. Orn. Cl., 13:54. HOLOTYPE, Adult male. Reg. no. 1911.12.23.3664. Ribola, Fernando Po, (31 Oct. vide Ibis, (8) 3 (1903): 355). Collected and bequeathed by Boyd Alexander. mayensis—Icterus chrysater mayensis van Rossem, 1938 57 Bull. Brit. Orn. Cl., 58: 136. HOLOTYPE, Adult male. Reg. no. 1898.12.14.193. (not 1888 as printed, op. cit.). Peto, Yucatan, Mexico. April 1888. Collected by G. F. Gaumer. Salvin-

Godman Collection.

334 mayeri
mayeri—Astrapia mayeri Stonor, 1939
mayi—Karua leucomela mayi Ashby, 1914
mbololo—Seicercus ruficapilla mbololo V. G. L. van Someren, 1939. 35 J. E. Afri. and Uganda nat. Hist. Soc., 14:91. Holotype, Adult male. Reg. no. 1948.11.10. Mount Mbololo, 5,000', Eastern Teita Range, Kenya, Oct. 1938. Collected by V. G. L. van Someren. Presented by the Coryndon Museum, Nairobi.
mbuleunsis—Bessonornis macclounii mbuluensis C. H. B. Grant & Mackworth-Praed, 1936
mbuluensis—Zosterops virens mbuluensis W. L. Sclater and Moreau, 1935
mcchesneyi—Aethocorys personata mcchesneyi Williams, 1957. 17 Bull. Br. Orn. Cl., 77:157. HOLOTYPE, Adult male. Reg. no. 1956.6.6. Marsabit Plateau, 4,000′, 6 miles north of Marsabit boma, North Frontier Province, Kenya, 22 June 1955. Collected and presented by J. G. Williams.
mcclellandi—Pomatorhinus mcclellandi Godwin-Austen, 1870 32E J. Asiat. Soc. Bengal. 39, pt. 2:103. SYNTYPE, Adult. Reg. no. 1895.7.14.2315. Nenglo, Asalu, (under Burrail Hills on label). Khasi Hills, Assam. Collected by H. H. Godwin-Austen.

There are several other syntypes in the collection. Pomatorhimus mclellandi: Jerdon, The Birds of India, 2 (1863): 32 is a nomen nudum.

Godwin-Austen Collection.

mcclellandii—Hypsipetes mcclellandii McClelland, 1840 in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840): 159. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4747. Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.

mearsi—Pomatorhinus mearsi Ogilvie-Grant, 1905
media—Fringillaria media Sharpe, 1904
media—Urobrachya media Sharpe, 1902
media—Zosterops auriventer media Robinson and Kloss, 1923 . 47 Ibis, (11) 5:152. HOLOTYPE, Adult male. Reg. no. 1927.4.18.966. (Bukar, on label), 1,000', Gunong Sidong, Samarahan, South-west Sarawak, Borneo, 28 Oct. 1919. Collected by Ulck for R. C. Robinson. Presented by R. C. Robinson. Labelled as the type by the authors.
mediocris—Cinnyris mediocris Shelley, 1885
Proc. zool. Soc. Lond., 1863: 488. SYNTYPE, Adult (female on label). Reg. no. 1873.5.12.750. East Timor, 1861. Collected by A. R. Wallace. Wallace Collection.
Report on the Zoological Collections of H.M.S. Alert: 14. HOLOTYPE, Adult male. Reg. no. 1882.2.18.59. Australia (Port Molle, Queensland, 9 May 1881, vide Gould, Birds of New Guinea, 2 (1888): pl. 50). Collected by Dr. R. W. Coppinger. Presented by the Lords of the Admiralty.
Bull. Br. Orn. Cl., 67:26. HOLOTYPE, Adult male. Reg. no. 1946.5.11. 10 miles north of Mega, south Abyssinia, 5 Sept. 1941. Collected and presented by C. W. Benson.
Bull. Br. Orn. Cl., 63:10. HOLOTYPE, Adult male. Reg. no. 1946.5.1648. 10 miles north of Mega, 4,000', south Abyssinia, 10 Sept. 1941. Collected and presented by C. W. Benson.

- magaensis—Pseudalaemon fremantlii megaensis Benson, 1946. . . 17

 Bull. Br. Orn. Cl., 67:25. HOLOTYPE, Adult male. Reg. no. 1946.5.10. 15 miles north of Mega, south Abyssinia, 12 Oct. 1941. Collected and presented by C. W. Benson.
- megalolophus—Trochocercus megalolophus Swynnerton, 1907 . 38

 Bull. Brit. Orn. Cl., 19: 109. Syntype, Adult male. Reg. no. 1910.7.1.362. Jihu District, 2,000', Gazaland, (12 August 1906, on label). Collected by and purchased of C. F. M. Swynnerton.

 Labelled as the type by the author, but three other syntypes were available and are in the collection.
- megalornis—Dicrurus megalornis G. R. Gray, 1858 64

 Proc. zool. Soc. Soc. Lond., 1858: 179. HOLOTYPE, Adult. Reg.
 no. 1873.5.12.1952. Ké Island, off New Guinea, 1857. Collected
 by A. R. Wallace. Wallace Collection.
- megarhyncha—Peucaea megarhyncha Salvin and Godman, 1889 . 49

 Ibis, (6) 1:238. HOLOTYPE, Adult female. Reg. no. 1899.2.1.

 3227a. Santa Ana, Sonora, Mexico, 18 April 1888. Collected by W. Lloyd. Salvin-Godman Collection.
- megarhynchus—Ploceus megarhynchus Hume, 1869 61 Ibis, (2) 5:356. SYNTYPE, Adult female. Reg. no. 1887.1.1.1660. Kaladoongi, (below Nynee Tal, vide Hume, Stray Feathers, 6 (1878): 400), India, Dec. 1866. Collected by A. O. Hume. Hume Collection.

The other syntype is in the collection.

- meinertzhageni—Anthus spinoletta meinertzhageni Bird, 1935. 19
 Bull. Br. Orn. Cl., 56:55. HOLOTYPE, Adult male. Reg. no. 1947.15.1. Lochboisdale, South Uist, Outer Hebrides, 5 April 1935. Collected and presented by E. G. Bird.
- meinertzhageni—Certhia familiaris meintzhageni Clancey, 1942 . 44A Bull. Brit. Orn. Cl., 63:42. Holotype, Adult male. Reg. no. 1965.M.49. Caragh Lake, Co. Kerry, south-western Ireland, 6 Jan. 1935. Collected by R. Meinertzhagen. Meinertzhagen Collection.

meinertzhageni—Troglodytes	troglodytes	meinertzh	ageni	Clancey,	,
1942					28
Bull. Br. Orn. Cl., 62:0					
1965.M.21. Ushant, Fini				Collect	ted
by R. Meinertzhagen. M	einertzhagen	Collection			

- melaena—Munia melaena P. L. Sclater, 1880 Proc. zool. Soc. Lond., 1880 : 66, pl. 7. Holotype, Adult. no. 1881.3.29.38. Kabahadai, New Britain, February 60 Collected by the Rev. G. Brown. Purchased of E. Gerrard.
- melaena—Myrmotherula melaena P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1857: 130. SYNTYPE, Adult male. Reg. no. 1889.9.20.285. Bogota, Colombia. Collected for Verreaux. Sclater Collection.

The other syntype is in the collection.

- melampyra—Tchitrea melampyra Verreaux, 1857 in Hartlaub, System der Ornithologie Westafrika's.: 90. A specimen (no. 1875.4.9.48.) was regarded as the type by Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 357, and by Grant and Mackworth-Praed, Bull Brit. Orn. Cl., 77 (1957): 64, but has been shown to have the wrong measurements. It was received with Sharpe's Collection and is labelled "Gaboon, Verreaux". Another specimen (no. 1875.4.9.359) is labelled "Gaboon, Walker" and "Sharpe's Cat. Bds. Afr. 420a". It conforms with the measurements and description. We suspect that the labels were inadvertently changed, but this cannot be proven.
- melanocephala—Anthornis melanocephala G. R. Gray, 1843. in Dieffenbach's Travels in New Zealand, 2:188. HOLOTYPE, Adult male (relaxed mount). Chatham Island, New Zealand. Collected by Dieffenbach. Presented by the New Zealand Company.
- melanocephala—Cisticola melanocephala Anderson, 1871 Proc. zool. Soc. Lond., 1871:212. SYNTYPE, Adult. Reg. no. 1876.4.7.82. Sonda Valley, 1,700', Yunnan, China, July 1868. Collected by and purchased of J. Anderson.

Two specimens are referred to by the author in Anatomical and Zoological researches . . . Western Yunnan in 1868–1875, 2 (1878):

641.

- melanocephala—Guiraca melanocephala Swainson, 1827 Philosophical Magazine (n.s.) 1827:438. A specimen purchased from the Zoological Society Museum was listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 65. The type is, however, at Cambridge, vide van Rossem, Auk, 49 (1932): 489.
- melanocephala—Lonchura melanocephala McClelland, 1840 in Horsfield, Proc. Zool. Soc. Lond., 1839 (Publ. 1840):163. HOLOTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4691.

Assam. Collected by J. McClelland. Presented by the Honour-

able East India Company Museum.

Although labelled "Amadina sinensis" this is clearly the McClelland specimen listed as the type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 336.

- melanocephala—Motacilla melanocephala Lichtenstein, 1823. 19
 Verz. Doubl. Mus. Berlin, 1823: 36. A specimen in the collection
 was listed as type of Budytes melanocephalus of Sykes (Proc. zool.
 Soc. Lond., 1832: 90) by Sharpe, Cat. Bds. Brit. Mus., 10 (1885):
 531 but Lichtenstein's description has priority.
- melanocephala—Muscicapa melanocephala Latham, 1801 . . . 36
 Index Ornithologicus, Suppl.: lii. ?Holotype, Water-colour drawing. New South Wales. No. 187 in Watling, Original water-colour drawings . . . Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 141.

Vol. 3, no. 38 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 10 (1922): 116.

- melanocephalus—Lanius collurioides melanocephalus Delacour, 1926 23

 Bull. Br. Orn. Cl., 47: 13. SYNTYPE, Adult male. Reg. no. 1926.

 9.8.10. Dalat, south Annam, 13 Oct. 1925. Collected by J. Delacour on the Second Franco-British Expedition to French Indochina.

This is the author's selected male. His selected female syntype is also in the collection. (See also appendix, Corrections and Additions, vol. 2.)

- Reg. no. 1889.9.20.45. Huiro, 4,800', Urubamba Valley, south Peru, 29 June 1874. Collected by H. Whitely. Sclater Collection. There is another syntype in the collection.

The other syntypes were in the Senckenberg Museum, Frankfurt-

am-Main.

This specimen was labelled by Shelley as the type and corresponds in its measurements with the original description. Shelley, *Birds of Africa*, 2 (1900): 25, refers to the discovery of the type by Fischer and his illustration (pl. 1, fig. 2, *op. cit.*) appears to be of this specimen, but there is no evidence concerning the way in which it

came into his possession.

- melanogaster—Ploceus melanogaster Shelley, 1887 . . . 61
 Proc. zool. Soc. Lond., 1887 : 126, pl. 14. Holotype, Adult male.
 Reg. no. 1887.3.7.34. Cameroons, 8,000', Sept. 1886. Collected by H. H. Johnston.
- melanogaster—Thryothorus melanogaster Sharpe, 1881 . . . 28
 Cat. Bds. Brit. Mus., 6:230. Syntype, Adult male. Reg. no.
 1885.3.6.336. Bugaba, Chiriqui, Veragua, 1869. Collected by E.
 Arcé. Salvin-Godman Collection.
 There are three other syntypes in the collection.

The female syntype is also in the collection.

- melanolaema—Hypocnemis melanolaema P. L. Sclater, 1855 . . . 4
 Proc. zool. Soc. Lond., 1854 (publ. 1855): 254, pl. 72. SYNTYPE,
 Adult male. Reg. no. 1889.9.20.525. Chamicuros, Peru, 1854.
 Collected for Verreaux. Sclater Collection.
- melanolaema—Pipreola melanolaema P. L. Sclater, 1856 . . . 7

 Ann. Mag. nat. Hist., (2) 17:469. SYNTYPE, Adult male. Reg.
 no. 1888.1.13.1614. Caracas, Venezuela, 1853. (Collected by Parzudaki; on label). Sclater Collection. (cont.)

Hellmayr, Catalogue of birds of the Americas, 6 (1929): 113, states that the type is in Paris, but Sclater referred to specimens in various museums and all would appear to be syntypes.
melanolaemus—Buarremon melanolaemus P. L. Sclater and Salvin, 1879
melanoleuca—Rhodophila melanoleuca Jerdon, 1863 31 The Birds of India, 2:128. Syntype, Adult male. Reg. no. 1864. 10.27.1. Purneah, Ganges River, Upper India. Collected and presented by Dr. T. C. Jerdon. There is also a female syntype in the collection. This species was renamed Oreicola jerdoni by Blyth, Ibis, (2) 3 (1867):14, on the grounds of preoccupation by O. melanoleuca Bonaparte, 1854.
melanoleucos—Microtarsus melanoleucos Eyton, 1839 21 Proc. zool. Soc. Lond., 1839 : 102. HOLOTYPE, Adult. Reg. no. 1881.2.18.101. Malacca. Collected by Evans, who sent it to his brother. Eyton Collection; purchased of E. Garrard.
melanolora—Campephaga melanolora G. R. Gray, 1860 20 Proc. zool. Soc. Lond., 1860: 353. SYNTYPE, Adult male. Reg. no. 1860.2.4.44. Batchian, Moluccas. Collected by A. R. Wallace and purchased of S. Stevens. There are four other syntypes in the collection.
melanonotus—Sarcops melanonotus Ogilvie-Grant, 1906 62 Bull. Brit. Orn. Cl., 16:100. SYNTYPE, Adult male. Reg. no. 1905.11.26.7. Davao, south-east Mindanao, Phillipines, March 1905. Collected by W. Goodfellow. Purchased of E. Gerrard. The female syntype is also in the collection.
melanonotus—Thammophilus melanonotus P. L. Sclater, 1855. Proc. zool. Soc. Lond., 1855: 19, pl. 80. Holotype, Adult male. Reg. no. 1889.9.20.121. Santa Marta, Colombia, 1854. Collected for Verreaux. Sclater Collection.
melanopezus—Anabates melanopezus P. L. Sclater, 1858
melanophrys—Turdus melanophyrs Latham, 1801

British Museum (Natural History). Vide Sharpe, Hist, Coll. Brit.

Mus., 2 (1906): 136.

Vol. 2, no. 48 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 12 (1925): 32.

- melanopleura—Merula melanopleura Sharpe, 1900 in Seebohm, Monograph of the Turdidae, 2, pt. 9:87, pl. 103. HOLOTYPE, Adult male. Reg. no. 1896.6.1.613. Trinidad. Seebohm Bequest.
- melanopogon—Hypocnemis melanopogon P. L. Sclater, 1857. Proc. zool. Soc. Lond., 1857: 130. SYNTYPE, Adult male. Reg. no. 1889.9.20.537. Guiana (vide Sclater, Catalogue of American Birds, 1862: 188 and Hellmayr, Novit. Zool., 14 (1907): 381). Collected by Parzudaki. Sclater Collection. There are two other syntypes in the collection.
- melanops—Artamus melanops Gould, 1865 Proc. zool. Soc. Lond., 1865: 198. HOLOTYPE, Adult. Reg. no. 1881.5.1.4282. St. A'Beckett's Pool, Central Australia, Lat 28°30' (= St. A'Beckett's Pond, 30°10′ S., 138°20′ E., near Etadunna, vide Whittell, Literature of Australian birds (1954): 758), 23 Aug. 1863. Collected by S. White. Gould Collection.
- melanops—Buarremon melanops P. L. Sclater and Salvin, 1876 Proc. zool. Soc. Lond., 1876: 253. SYNTYPE, Adult. Reg. no. 1888.6.8.141. Simacu, Bolivia, 1875. Collected by C. Buckley. Salvin-Godman Collection.

There are several other syntypes in the collection.

melanops—Certhia melanops Latham, 1801. 48 Index Ornithologicus, Suppl.: xxxvi. ?SYNTYPES, Signed watercolour drawings. New South Wales. Nos. 99, 100, 101 in Watling, Original water-colour drawings. . . . Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 127.

Vol. 2, nos. 59, 61 in Lambert drawings, in Library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also

Mathews, Birds of Australia, 11 (1924): 350.

- melanops—Myiadestes melanops Salvin, 1865 Proc. zool. Soc. Lond., 1864 (publ. 1865): 580, pl. 35. SYNTYPE, Adult. Reg. no. 1885.3.2.514. Tucurrique, Costa Rica, 1864. Collected by E. Arcé. Salvin-Godman Collection. Listed as type by Hellmayr, Catalogue of Birds of the Americas, 7 (1934): 443.
- melanops—Zosterops melanops G. R. Gray, 1859 Catalogue of birds of the tropical islands of the Pacific Ocean, : 15. HOLOTYPE, Adult. Reg. no. 1854.5.31.12. Loyalty Islands. Presented by Sir George Grey.

- melanoptera—Monarcha melanoptera G. R. Gray, 1858 . . . 38

 Proc. zool. Soc. Lond., 1858: 178. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1858.10.11.11. Louisade Archipelago (Round Island, vide British Museum Register). Collected and presented by J. MacGillivray, from the voyage of H.M.S. Rattlesnake.
- melanoptera—Prionops melanoptera Sharpe, 1901. 23

 Bull. Br. Orn. Cl., 11:46. SYNTYPE, Adult female. Reg. no. 1898.12.11.56. Fer-Libah, Somaliland, 20 Oct. 1896. Collected by A. E. Atkinson and presented by Lord Delamere.

 The male syntype is also in the collection.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 3 (1877): 61. A female specimen in the Academy of Natural Sciences of Philadelphia was listed as the type by Stone and Mathews, Austral. Avian Record 1 (1913): 180, but not included by De Schauensee, Proc. Acad. nat. Sci. Philad., 109 (1957): 123–246.

melanopygius—Lanio melanopygius Salvin and Godman, 1883. 51

Biol. Centr.-Amer., Aves., 1 (Dec. 1883): 305. Syntype Adult
male. Reg. no. 1885.6.7.519. Bugaba, Panama. Collected by E.
Arcé. Salvin-Godman Collection.

Salvin and Godman used a Ridgway MS. name, Ridgway publishing his description in April 1884. The female and immature syntypes from the type locality are in the collection.

- melanorhyncha—Lusciniola melanorhyncha Rickett, 1898 . . . 35

 Bull. Brit. Orn. Cl., 8:10. HOLOTYPE, Adult male. Reg. no. 1900.3.10.13. Kuatun, north-west Fohkien, China, 4 May 1897. Collected and presented by C. B. Rickett.
- melanorhynchus—Drymoica melanorhynchus Jardine and Fraser, 1852 35 in Jardine, Contributions to Ornithology: 60. SYNTYPE, Adult male. Reg. no. 1886.6.24.218. Abomey, West Africa. Collected by L. Fraser. Jardine Collection.

The tail is incomplete. This specimen is labelled as the authors' specimen but there are other syntypes in the collection.

melanosticta—Hypocnemis melanosticta P. L. Sclater, 1855 . . . 4
Proc. zool. Soc. Lond., 1854 (publ. 1855): 254, pl. 73. SYNTYPE,
Immature male (vide Sclater, Cat. Bds. Brit. Mus., 15 (1890): 289).

- Reg. no. 1889.9.20.523. Chamicuros, Peru. Collected for Verreaux. Sclater Collection.

 There is a female syntype in the collection.
- melanosticta—Pithys melanosticta P. L. Sclater & Salvin, 1880 . 4
 Proc. zool. Soc. Lond., 1880: 160. HOLOTYPE, Adult (female, vide Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 304).
 Reg. no. 1889.7.10.747. Sarayacu, Ecuador. Collected by C. Buckley. Salvin-Godman Collection.
- melanostoma—Harporhynchus melanostoma Salvin, 1885 . . . 29

 Ibis, (5) 3:187. SYNTYPE, Adult. Reg. no. 1885.3.2.624.

 Cozumel Island, off Yucatan, April 1884. Collected by E. C. J. Devis. Salvin-Godman Collection.

 The second syntype is also in the collection.
- melanothorax—Thammophilus melanothorax P. L. Sclater, 1857. 4

 Proc. zool. Soc. Lond., 1857: 133. HOLOTYPE, Adult (female, vide Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 59). Reg. no. 1846.7.16.177. South America. Presented by Edward Wilson.
- melanotis—Calliste melanotis P. L. Sclater, 1876 51

 Ibis, (3) 6: 408, pl. 12. HOLOTYPE, Adult. Reg. no. 1885.6.12.375.

 Napo River, western Ecuador. Collected for Verreaux, Sclater Collection.

Renamed Tangara lutleyi Hellmayr, Verh. Orn. Ges. Bayern., 13 (1917): 198, on the grounds of preoccupation by Agelaia melanotis Swainson. 1837.

- melanotis—Campephaga melanotis G. R. Gray, 1860 . . . 20

 Proc. zool. Soc. Lond., 1860 : 353. SYNTYPE, Adult male. Reg.
 no. 1873.5.12.300. Gilolo, 1858. Collected by A. R. Wallace.
 Wallace Collection.

The second syntype is also in the collection.

- melanotis—Granatellus venustus melanotis van Rossem, 1940 . . . 54

 Proc. Biol. Soc. Wash., 53:14. HOLOTYPE, Adult female. Reg.
 no. 1892.4.20.1589. Tepic, Mexico. 1 May 1889. Collected by
 W. B. Richardson. Salvin-Godman Collection.

344 meranous
melanotis—Orpheus melanotis Gould, 1837
See Swarth, Occ. Papers California Acad. Sci., 18 (1931): 119 for a note on this specimen. The second syntype is in the collection. melanotis—Pteruthius melanotis Hodgson, 1847 32B in Blyth, J. Asiat. Soc. Bengal, 16: 448. Gadow, Cat. Bds. Brit. Mus., 8 (1883): 118, claimed that the types were in the Brit. Mus. collection, received from the Honourable East India Company Museum. These specimens were not in the possession of the latter museum until 1853, however, and it is therefore unlikely that they are the syntypes (contra Diegnan, in Peters, Checklist of Birds of the World, 10 (1964): 338).
melanotis—Ptilonorhynchus melanotis G. R. Gray, 1858 69 Proc. zool. Soc. Lond., 1858 : 181. Holotype, Adult male (relaxed mount). Reg. no. 1858.3.10.47. Aru Island. Collected by A. R. Wallace. Wallace Collection.
melanoxantha—Phainoptila melanoxantha Salvin, 1877 25 Proc. zool. Soc. Lond., 1877 : 367. Holotype, Adult male. Reg. no. 1885.3.20.13. Costa Rica (San Francisco, Costa Rica, on label), Feb. 1877. Collected by H. Rogers. Salvin-Godman Collection.
melanozantha—Pachyglossa melanozantha Blyth, 1843 45 J. Asiat. Soc. Bengal, 12:1010. Holotype, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 10 (1885):72), (relaxed mount). Reg. no. 1845.1.9.285. Nepal. Collected and presented by B. H. Hodgson. Blyth used a Hodgson MS. name. Sclater, Ibis, (3) 4 (1874); 3 stated that Hodgson had loaned the specimen to Blyth and that it had subsequently come to the British Museum and had been mounted.
melanozanthos—Coccothraustes melanozanthos Hodgson, 1836 . 59 Asiat. Res., 19:150. Syntype, Adult male. Reg. no. 1845.1.9. 297. Nepal, northern region. Collected by and purchased of B. H. Hodgson. This specimen has an original label bearing the no. 326 (the plate number). There are several other syntypes in the collection.
melanthes—Lanius melanthes Swinhoe, 1867
melanura—Cinclosoma melanura Hodgson, 1836

- melanura—Enicornis melanura G. R. Grav and Mitchell, 1846 The Genera of Birds, 1:133, pl. 41. HOLOTYPE, Adult (relaxed mount). Reg. no. 1845.11.8.64. No locality (Chile, vide Sclater, Cat. Bds. Brit. Mus., 15 (1890): 27). Purchased of Cuming.
- melanura—Euphonia melanura P. L. Sclater, 1851 in Jardine, Contributions to Ornithology: 86. SYNTYPE, Adult male. Reg. no. 1885.6.12.152. Barra do Rio Negro (= Manáos), Brazil. Collected by A. R. Wallace. Sclater Collection. Listed as the type by Sclater, Catalogue of a collection of American

birds (1862): 59.

- melanura—Hypocnemis melanura P. L. Sclater and Salvin, 1866 Proc. zool. Soc. Lond., 1866: 186. SYNTYPE, Adult male. Reg. no. 1889.9.20.530. Cashaboya, Ucayali River, east Peru, 10 Aug. 1865. Collected by E. Bartlett. Sclater Collection. The female syntype is also in the collection.
- melanura—Mviagra melanura G. R. Grav. 1859. Catalogue of the birds of the tropical islands of the Pacific Ocean: 18. SYNTYPE, Adult male. Reg. no. 1859.6.14.9. Erromango, New Hebrides. Collected by J. Macgillivray. Purchased of Stevens. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 380. The female syntype from Aneiteum is also in the Collection.
- melanura—Rhipidura melanura J. E. Grav. 1843 in, Dieffenbach, Travels in New Zealand, 2:190. Specimens collected by Capt. Stokes of H.M.S. Acheron and listed as types by Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 316, were probably collected later than the date of the description.
- melanurus—Thamnophilus melanurus P. L. Sclater, 1855 Edinb. New philos. J., (n.s.) 1:233. SYNTYPE, First year male (vide Hellmayr, Catalogue of Birds of the Americas, 3 (1924):50). Reg. no. 1855.11.5.17. Ucavali River, Peru. Collected by J. Hauxwell and purchased of J. Gould. Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890: 186.

Sclater used Gould's MS. name.

melas—Oryzoborus melas P. L. Sclater and Salvin, 1868 Proc. zool. Soc. Lond., 1867 (publ. 1868): 978. SYNTYPE, Adult male. Reg. no. 1885.2.10.722. Pebas, Upper Amazons, Peru, (2 April 1867, on label). Collected by J. Hauxwell. Sclater Collection.

Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 80. There are two other syntypes in the collection.

melaschistos—Volvocivora melaschistos Hodgson, 1837. 20 Indian Rev., 1 1836 (publ. 1837): 328. SYNTYPE, Adult male. Reg. no. 1843.1.13.729. Nepal. Collected and presented by B. H. Hodgson.

The female syntype is also in the collection.

melindae—Anthus melindae Shelley, 1900
mellivora—Certhia mellivora Latham, 1801
melophilus—Erithacus rubecula melophilus Hartert, 1901
menachensis—Poliospiza menachensis Ogilvie-Grant, 1913 59 Bull. Brit. Orn. Cl., 31:90. SYNTYPE, Adult male. Reg. no. 1913.7.18.11. Menacha, 7,000′, Yemen, 31 Jan. 1913. Collected by and purchased of G. W. Bury. The female syntype is also in the collection.
menachensis—Turdus menachensis Ogilvie-Grant, 1913 3 Bull. Br. Orn. Cl., 31:86. SYNTYPE, Adult male. Reg. no. 1913. 7.18.1. Menacha, Yemen mountains, south Arabia, 28 Dec. 1912. Collected by and purchased of G. W. Bury. The female syntype is also in the collection.
mendozae—Pseudochloris mendozae Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:778. HOLOTYPE, Adult male. Reg. no. 1885.2.10.754. Mendoza, Argentina. Collected by Burmeister. Sclater Collection.
meneliki—Oriolus meneliki Blundell and Lovat, 1899 66 Bull. Brit. Orn. Cl., 10: 19. Holotype, Adult. Reg. no. 1900.1.3. 121. Burka, south Abyssinia, 6 Jan. 1899. Collected and presented by H. W. Blundell and Lord Lovat.
mennelli—Poliospiza mennelli Chubb, 1908
mentalis—Catharus mentalis P. L. Sclater & Salvin, 1876 3 Proc. zool. Soc. Lond., 1876: 352. SYNTYPE, Adult. Reg. no. 1885.3.2.35. Suape, near Tilotilo, Yungas Province, Bolivia, 1876. Collected by C. Buckley. Salvin-Godman Collection. The second syntype is also in the collection.

meridae 347 mentalis—Drymoica mentalis Fraser, 1843. Proc. zool. Soc. Lond., 1843: 16. HOLOTYPE, Adult. Reg. no. 1847.1.18.29. Accra. West Africa. Collected by and purchased of L. Fraser. mentalis—Graucalus mentalis Vigors & Horsfield, 1827 Trans. Linn. Soc. Lond., 15: 217. HOLOTYPE (Adult, probably male. by plumage, vide I. C. J. Galbraith MS, note: Sharpe, Cat. Bds. Brit. Mus., 6 (1879): 38, and others, listed it as a juvenile). Reg. no. 1863.7.7.20. "South Coast of Australia, 1803" (error, 1802, vide Brown's MS.). Collected by R. Brown and presented by the Linnean Society. Brown's main collecting localities on the south coast were at the head of Spencer's Gulf, South Australia, and Port Phillip Bay, Victoria. As the species to which this specimen belongs (i.e. Coracina papuensis) occurs only marginally in South Australia, it is

more likely that the holotype came from Port Phillip Bay.

- mentalis—Pachycephala mentalis Wallace, 1863 Proc. zool. Soc. Lond., 1863: 30. SYNTYPE, Adult male. Reg. no. 1873.5.12.513. Batchian Island, south-west Pacific, 1858. Collected by A. R. Wallace. Wallace Collection. Two males are listed as types by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 190.
- mentalis—Pipra mentalis P. L. Sclater, 1856 Proc. zool. Soc. Lond., 1856: 299, pl. 121. Syntype, Adult male (relaxed mount). Reg. no. 1857.7.30.6. Cordova, Vera Cruz, South Mexico. Collected by and purchased of A. Sallé. The female syntype is also in the collection.
- mentalis—Rhodospingus mentalis Sharpe, 1888 Cat. Bds. Brit. Mus., 12:809, pl. 15. HOLOTYPE, Adult male. Reg. no. 1885.12.14.715. Puna Island, Gulf of Guayaquil, Ecuador. Collected by Villagomez for C. Buckley. Salvin-Godman Collection.
- mentalis—Sigmodius mentalis Sharpe, 1884. J. Linn. Soc. Lond., 17: 425. Syntype, Adult. Reg. no. 1884.5.1. 22. Sassa, Belgian Congo, Oct.. Collected by and purchased of F. Bohndorff.
- mentawi—Oriolus xanthonotus mentawi Chasen and Kloss, 1926 . 63 Ibis, (12) 2:295. HOLOTYPE, Adult female. Reg. no. 1947.60.31. Siberut Island, west Sumatra, 13 Sept. 1924. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore.
- meridae—Buarremon meridae P. L. Sclater and Salvin, 1870 Proc. zool. Soc., Lond., 1870: 785. HOLOTYPE, Adult male. Rcg. no. 1885.6.12.934. Merida, Venezuela, July 1869. Collected by A. Goering. Sclater Collection.

546 meridana
meridana—Cyanocitta armillata var. meridana P. L. Sclater and Salvin, 1876
meridanus—Basileuterus meridanus Sharpe, 1885 5 Cat. Bds. Brit. Mus., 10:387. HOLOTYPE, Adult male. Reg. no. 1885.3.8.670. Merida, Venezuela, July 1869. Collected by A. Goering. Salvin-Godman Collection.
meridionale—Trochalopterum meridionale Blanford, 1880 32 J. Asiat. Soc. Bengal, 49, pt. 2:142. SYNTYPE, Adult male, Reg. no. 1880.8.19.10. Travancore – Tinnevelly boundary, 4,400', south India, 3 April 1880. Collected by and purchased of F. W. Bourdillon. The two other syntypes are also in the collection.
meridionalis—Certhia discolor meridonalis Robinson and Kloss, 1919 44. Ibis, (11) 1:609, pl. 16. SYNTYPE, Adult male. Reg. no. 1919.12. 20.436. Langbian Peaks, 6-7,000', South Annam, 8 April 1918 (apparently a misprint, the correct date is 18 April). Collected and presented by C. B. Kloss. The female syntype is also in the collection.
meridionalis—Culicicapa ceylonensis meridionalis E. C. S. Baker, 1923 37, Bull. Brit. Orn. Cl., 44:12. HOLOTYPE, Adult female. Reg. no. 1916.12.27.928. Keo, Tung Song, Siam, 22 Sept. 1915. Collected by C. Chunggat for E. G. Herbert. Presented by E. G. Herbert.
meridionalis—Dicrurus leucogenys meridionalis Hachisuka, 1926. 6 Bull. Brit. Orn. Cl., 47:56. HOLOTYPE, Adult male. Reg. no. 1909.8.30.36. Seven Finger Range, Central Hainan, 16 Nov. 1906. Collected and presented by R. Douglas.
meridionalis—Garrulax pectoralis meridionalis Robinson & Kloss, 1919
meridionalis—Loxia curvirostra meridionalis Robinson and Kloss, 1919

by C. B. Kloss.

This is the authors' selected male syntype. The female syntype is also in the collection.

meridionalis—Parus meridionalis P. L. Sclater, 1857 42 Proc. zool. Soc. Lond., 1856 (publ. 1857): 293. Holotype, Adult male. Reg. no. 1857.7.30.24. (Collector's no. 167). El. Jacale, Vera Cruz, south Mexico, May 1855. Collected by and purchased of A. Sallé. (See also appendix, Corrections and Additions, vol. 2.)	2
meridionalis—Rectes meridionalis Sharpe, 1888 39 Ibis, (5) 6:437. SYNTYPE, Adult male. Reg. no. 1880.9.13.43. Astralobe Mountains, south-east New Guinea. Collected by C. Hunstein. Purchased of E. Gerrard. The second syntype is also in the collection.	9
meridionalis—Sturnella meridionalis P. L. Sclater, 1861 57 Ibis, (1) 3:179. SYNTYPE, Adult. Reg. no. 1885.7.12.210. Bogota, Colombia. Collected for Bell. Sclater Collection. Listed as the type by Hellmayr and Conover, Catalogue of birds of the Americas, 10 (1927):215.	7
merrotsyi—Amytornis merrotsyi Mellor, 1913	6
meruensis—Zosterops virens meruensis W. L. Sclater and Moreau, 1935	7
meruloides—Mimeta meruloides Vigors and Hoisfield, 1827 63. Trans. Linn. Soc. Lond., 15:327. HOLOTYPE, Immature (vide Sharpe, Cat. Bds. Brit., Mus., 3 (1877):2,3). Reg. no. 1863.7.7.70. New South Wales. Collected by Caley. Presented by the Linnean Society.	3
mesochrysus—Basileuterus mesochrysus P. L. Sclater, 1860 54 Proc. zool. Soc. Lond., 1860: 251. HOLOTYPE. Adult. Reg. no. 1884.5.15.434. Bogota, Colombia. (Collected by Mather, on label). Sclater Collection.	4
mesoleuca—Dendrophila mesoleuca Ogilvie-Grant, 1894 4. Bull. Brit. Orn. Cl., 3:49. Holotype, Adult male. Reg. no. 1897.5.13.92. La Trinidad, Benguet District, nort-west Luzon, (4 Jan. 1894, on label). Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition.	3

mesoleucus—Basileuterus mesoleucus P. L. Sclater, 1865 . . . 54 Proc. zool. Soc. Lond., 1865: 286, pl. 9. HOLOTYPE, Adult. Reg.

no. 1884.5.15.462. Demerara, Guiana, 1864. (Collected by Wilkens, on label). Sclater Collection, received by him from Hartlaub of Bremen.

mesoleucus—Contopus mesoleucus P. L. Sclater, 1859 9

Proc. zool. Soc. Lond., 1859: 43. SYNTYPE, Adult. Reg. no. 1888.1.13.994. Orizaba, Mexico. Collected by Botteri. Sclater Collection.

Listed as the type by Sclater, Catalogue of a collection of American birds (1862): 230. The species had been described previously, unknown to Sclater, as Muscicapa mesoleuca Lichtenstein, Preis-Verzeichniss Mexicanischer Vögel. . . . 1830: 2, (reprinted in J. Orn., 1863: 56), vide Salvin and Godman, Biol. Centr.-Amer., Birds, 2 (1889): 81.

- mesopotamia—Pycnonotus leucotis mesopotamia Ticehurst, 1918 . 21 J. Bombay nat. Hist. Soc., 26: 279. HOLOTYPE, Adult male. Reg. no. 1920.10.23.2. Basra, Mesopotamia, 20 Nov. 1917. Collected and presented by C. B. Ticehurst.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 66.

- mexicana—Certhiola mexicana P. L. Sclater, 1856 54

 Proc. zool. Soc. Lond., 1858: 286. HOLOTYPE, Adult. Reg. no. 1857.7.30.17. Cordova, Vera Cruz. Collected by and purchased of A. Sallé.
- mexicana—Sturnella mexicana P. L. Sclater, 1861 57

 Ibis, (1) 3:179. SYNTYPE, Adult. Reg. no. 1885.7.12.206.

 Jalapa, Vera Cruz, Mexico. Collected by De Oca. Sclater Collection.

Listed as the type by Hellmayr and Conover, Catalogue of Birds of the Americas, 10 (1927): 213.

- Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 214. There are two other syntypes in the collection. mexicanus—Sclerurus mexicanus P. L. Sclater, 1857 3 Proc. zool. Soc. Lond., 1856 (publ. 1857): 290. SYNTYPE, Adult. Reg. no. 1889.5.20.417. Cordova, Vera Cruz, southern Mexico. Collected by A. Sallé. Sclater Collection. There are three other syntypes in the collection. mexicanus—Xenops mexicanus P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1856 (publ. 1857): 289. SYNTYPE, Adult female. Reg. no. 1857.7.30.14. Cordova, Vera Cruz, southern Mexico. Collected by and purchased of A. Sallé. microptera—Mirafra microptera Hume, 1873 Stray Feathers, 1:483. SYNTYPE, Adult female. Reg. no. 1887.7. 1. 2668. Palow, Thayetmyo district, Burma, 26 May 1872. Collected by E. W. Oates. Hume Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 615. There are several other syntypes in the collection. micropterus—Agathopus micropterus P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858:69. SYNTYPE, Adult. Reg. no. 1889.9.20.700. Rio Napo, Ecuador, Probably collected by Villavicencio. Sclater Collection. There is another syntype in the collection. micropterus—Agriornis micropterus Gould, 1839 Zoology of the voyage of H.M.S. Beagle, 3, Birds: 57, pl. 12. SYNTYPE, (female in worn breeding garb vide Hellmayr, Catalogue of birds of the Americas, 5 (1927): 3) (relaxed mount). Reg. no. 1855.12.19.253. Port Desire, Patagonia, January. Collected by C. Darwin. Purchased of the Zoological Society of London as the type, but two specimens are referred to in the text. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 4 (1888): 15. microrhyncha—Lalage karu microrhyncha Ogilvie-Grant, 1915 Ibis, (10) 2, Jubilee Supplement: 118. HOLOTYPE, Adult male. Reg. no. 1911.12.20.1595. Mouth of Mimika River, Dutch New Guinea, 4 Dec. 1910. Collected by C. H. B. Grant and presented by the Committee of the British Ornithologists' Union Expedition to Dutch New Guinea microrhynchus—Cassiculus microrhynchus P. L. Sclater and Salvin, Proc. zool. Soc. Lond., 1864: 353. SYNTYPE, Adult male. Reg. no. 1885.7.12.61. Panama Railroad, Panama, 1863. Collected by J. McCleannan. Sclater Collection. There are three other syntypes in the collection. microrhynchus—Cinnyris microrhynchus Shelley, 1876. 46
- microrhynchus—Cinnyris microrhynchus Shelley, 1876 . . . 46
 Monograph of the Nectariniidae : 219, pl. 67. SYNTYPE, Immature

male. Reg. no. 1878.10.26.132. Mombasa. Collected by the

Rev. T. Wakefield. Sharpe Collection.

The two other immature syntypes are also in the collection. Gadow, Cat. Bds. Brit. Mus., 9 (1884): 48, listed an adult male as the type in error.

There are several other syntypes in the collection.

- milanjensis—Xenocichla milanjensis Shelley, 1894 . . . 21 Ibis, (6) 6:9, pl. 1. HOLOTYPE, Adult (female on label; feathers of neck missing). Reg. no. 1893.6.1.89. Milanji Hills, Nyasaland, Sept. 1892. Collected by A. Whyte and presented by Sir H. H. Johnston.
- milleti—Garrulax milleti Robinson & Kloss, 1919 32B Ibis, (11) 1:574, pl. 12. Syntype, Adult male. Reg. no. 1919.12. 20.266. Dalat, 5,000′, south Annam, 2 May 1918. Collected and presented by C. B. Kloss.

This is the author's selected male. Their selected female syntype

is also in the collection.

- mimika—Quoyornis leucurus mimika Mathews, 1931 37

 Bull. Brit. Orn. Cl., 52:25. Holotype, Adult male. Reg. no.
 1911.12.20.1181. Mouth of Mimika River, Dutch New Guinea,
 22 March 1911. Collected by C. H. B. Grant. Presented by the
 subscribers to the British Ornithologists' Union Expedition to Dutch
 New Guinea.
- mimika—Setosura maculipectus mimika Mathews, 1928 . . . 37A Bull. Brit. Orn. Cl., 48:91. Holotype, Adult male. Reg. no. 1911.12.20.1189. Mimika River, Dutch New Guinea, 18 May, 1911. Collected by C. H. B. Grant. Presented by the subscribers to the British Ornithologists' Union Expedition to Dutch New Guinea.
- mimikae—Myiagra mimikae Ogilvie-Grant, 1911 38

 Bull. Brit. Orn. Cl., 29:26. SYNTYPE, Adult male. Reg. no.
 1911.12.20.1343. Mouth of the Mimika River, Dutch New
 Guinea, 22 March 1911. Collected by C. H. B. Grant. Presented
 by the subscribers to the British Ornithologists' Union Expedition
 to Dutch New Guinea.

This is the described male. The described female syntype is also

in the collection.

mimikae—Pseudogerygone conspicillata mimikae Ogilvie-Grant, 1915 36

Ibis, (10) 2, Supplement: 168. SYNTYPE, Adult male. Reg. no.
1911.12.20.1116. Mouth of Mimika River, Dutch New Guinea,
20 May 1911. Collected by C. H. B. Grant. Presented by the
subscribers to the British Ornithologists' Union Expedition to
Dutch New Guinea.

This is the described male, the described female syntype is also in the collection

The female syntype is also in the collection.

- mindanensis—Artamides mindanensis Steere, 1890 20
 List of the Birds & Mammals collected by the Steere Expedition to the
 Philippines: 14 (reprinted in Ibis, (6) 3 (1891): 309). SYNTYPE,
 Adult male. Reg. no. 1896.6.6.49. Ayala, Mindanao, 1 Oct.
 1887. Collected by and purchased of J. B. Steere.
 The immature syntype is also in the collection.
- mindanensis—Macronus mindanensis Steere, 1890 32E List of the Birds and Mammals collected by the Steere Expedition to the Philippines: 17 (reprinted in Ibis, (6) 3 (1891): 312). SYNTYPE, Adult male. Reg. no. 1896.6.6.381. (Ayala, 17 Oct. on label), Mindanao, Philippines. Collected and purchased by J. B. Steere.

There is also a female syntype in the collection.

mindanensis—Mirafra javanica mindanensis Hachisuka, 1931 . 17
Oiseau, (n.s.) 1:471. Syntype, Adult male. Reg. no. 1932.6.2.14.
Gogoag, Mindanao, Philippines, 1930. Collected by M. Nakamura and presented by M. Hachisuka.

There are three other syntypes in the collection.

- mindanensis—Volvocivora mindanensis Tweeddale, 1878 . . . 20

 Proc. zool. Soc. Lond., 1878: 947. Syntype, Adult male. Reg.
 no. 1887.12.30.135. Zamboagna, Mindanao, Philippines, June
 1878. Collected by A. H. Everett. Tweeddale Collection.
 The adult and immature female syntypes are also in the collection.

the Philippines	s: 14 (reprinted in <i>Ibis</i>	s, (6) 3 (1891) : 309).	SYNTYPE,
Adult male.	Reg. no. 1896.6.6.47.	Calaran, Mindoro,	Philippines,
1 June 1888.	Collected by and pur	chased of J. B. Steer	e.
There is an	other syntype in the co	ollection.	

- mindorensis—Muscicapula hyperythra mindorensis Hachisuka, 1935 37

 Birds of the Philippine Islands, 2:299. HOLOTYPE, Adult female.

 Reg. no. 1897.5.13.362. Mount Dulangan, 4,500', Mindoro,
 Philippines, 26 Jan. 1896. Collected by J. Whitehead. Presented
 by the subscribers to the Whitehead Expedition.
- mindorensis—Turdus mindorensis Ogilvie-Grant, 1896 3:

 Ibis, (7) 2:465. Syntype, Adult male. Reg. no. 1897.5.13.376.

 Mount Dulangan, Mindoro, 27 Dec. 1895. Collected by J.

 Whitehead and presented by the subscribers to his expedition.

 This is the author's selected male. His selected female syntype is also in the collection.
- minima—Geokichla interpres minima Hachisuka, 1934. . . . 31
 Tori, 8:221. HOLOTYPE, Adult male. Reg. no. 1896.6.6.267.
 Malaunavi, Basilan, Philippines. Collected by and purchased of J. B. Steere.

This is the author's selected male syntype. The selected female and several other syntypes are in the collection.

- minima—Platystira minima J. and E. Verreaux, 1855 38

 Rev. Zool., 7:219. SYNTYPE, Adult male. Reg. no. 1854.4.25.2.

 Gaboon, West Africa. Purchased of Verreaux.

 Hartlaub, J. Orn., 1861:169, confirms that of the two syntypes there was an adult in the British Museum Collection.

3.85. Manda Island, British East Africa. Collected and presented by F. J. Jackson. The female syntype is also in the collection. minimus—Anthus minimus Vigors and Horsfield, 1827. Trans. Linn. Soc. Lond., 15:230. SYNTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.32.b. Australia. Presented as the type by the Linnean Society of London. This specimen has a broken bill. There is another syntype in the collection Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 305). Reg. no. 1881.5.1. 339. (Somerset, on label), Cape York, Australia. (Collected by Cockerell, vide Sharpe, op. cit.). Gould collection. This specimen was labelled by Gould as minimus, and listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 305, but a number of specimens were referred to in the description. minor—Acrocephalus boeticatus minor Lynes, 1921 Bull. Brit. Orn. Cl., 43:96. HOLOTYPE, Adult male. Reg. no. 1922.12.18.1014. Zalingei, 3,000', West Central Darfur, Sudan, 2 Sept. 1921. Collected and presented by Admiral H. Lynes. From the Lynes-Lowe Expedition. minor—Actinodura ramsavi minor Kinnear, 1925. Bull. Br. Orn. Cl., 45:74. HOLOTYPE, Adult male. Reg. no. 1924.12.21.179. Ngoi-Tio, Tonkin, 2 June 1924. Collected by H. Stevens and presented by the Percy Sladen Trust and Salvin-Godman Exploration Fund. minor—Anthreptes seimundi minor Bates, 1926 46 Bull. Brit. Orn. Cl., 46:107. HOLOTYPE, Adult male. Reg. no. 1926.8.8.9. Sanaga River, (1,700', on label), north of Yaunde, Cameroons, 23 Jan. 1924. Collected by and purchased of G. L. Bates. (See also appendix, Corrections and Additions, vol. 2.) minor—Apalis jacksoni minor Ogilvie-Grant, 1917 Ibis, (10) 5:76. SYNTYPE, Adult male. Reg. no. 1920.6.26.475. Bitye, 2,000', Ja River, Cameroons, 8 Dec. 1912. Collected and presented by G. L. Bates. This is the described male. The female syntype is also in the collection. minor—Campophaga minor Davison, 1892. Ibis, (6) 4:99. HOLOTYPE, Adult male. Reg. no. 1892.9.1.4. Qualla Tahau, east coast of Malaya, 20 July 1891. Collected and presented by W. Davison.

minor—Cnipodectes minor P. L. Sclater, 1884

Proc. zool. Soc. Lond., 1883 (publ. 1884): 654. HOLOTYPE,

(Adult male, on label). Reg. no. 1888.1.13.817. Chamicuros, east Peru. (28 Jan. 1867, on label). Collected by E. Bartlett. Sclater Collection.
minor—Lamprolia minor Layard, 1876
minor—Nilaus minor Sharpe, 1895
minor—Pellorneum minor Hume, 1873
minor—Percnostola minor Pelzeln, 1869
minor—Pseudolalage minor Steere, 1890
minor—Pyrenestes minor Shelley, 1894
minor—Rhabdornis minor Ogilvie-Grant, 1896
minor—Sturnus minor Hume, 1873

28. Sindh, India, (near Gunga Dhun, Larkhana, 11 Jan. 1872, on label). Collected for A. O. Hume. Hume Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 39.
minor—Trichastoma minor Hume, 1874
minula—Sylvia minula Hume, 1873
minuta—Hirundo minuta Hodgson, 1844
minuta—Locustella minuta Swinhoe, 1863
minutus—Arundinax minutus Swinhoe, 1860
Proc. zool. Soc. Lond., 1858: 292. SYNTYPE, Adult. Reg. no. 1853.1.26.5. Ashantee, West Africa. Collected by and purchased of L. Fraser. The other syntype is also in the collection.
nirabilis—Dicrurus mirabilis Walden and E. L. Layard, 1872. 64 Ibis, (2) 3: 103, pl. 5. HOLOTYPE, Adult. Reg. no. 1887.11.20.637. Negros, Philippines, March 1871. Collected by E. L. Layard. Tweeddale Collection.
Ibis, (7) 3:380. SYNTYPE, Adult female. Reg. no. 1898.10.10.3. Mt. Scratchley, 12,200', New Guinea, 1 Oct. 1896. Collected by A. Giulianetti for Sir W. Macgregor (vide op. cit., : 371). Presented by C. W. De Vis as one of the types.

miserabilis—Perissotriccus ecaudatus miserabilis Chubb, 1919. Ann. Mag. nat. Hist., (9) 4:301. Holotype, Adult. Reg. no. 1922.3.5.2695. Bonasika River, British Guiana, 1911. Collected by F. V. McConnell. McConnell Collection.
mixta—Pachyprora mixta Shelley, 1889
mjæbergi—Dendrobiastes hyperythrus mjæbergi Hartert, 1925. 3 Sarawak Mus. J., 8:3. Holotype, Adult female. Reg. no. 1925. 9.10.1. Mount Poi, 5,300', West of Kuching, Sarawak, Borneo, 30 Sept. 1923 (Oct., on label). Collected by E. Mjöberg. Presented by the Sarawak Museum. The male syntype is also in the collection.
mocuba—Cisticola chiniana mocuba Vincent, 1933
modesta—Anthreptes modesta Eyton, 1839
modesta—Bessonornis modesta Shelley, 1897
modesta—Bradyornis modesta Shelley, 1873
modesta—Ethyrura modesta, Sharpe, 1862
modesta—Glyciphila modesta G. R. Gray, 1858

	Brit. Mus., 9 (1884): 216, also lists a specimen under G . incand as the type of G . modesta but this is an error and the latter has no type status.	
me	odesta—Grallaria modesta P. L. Sclater, 1877	4
	odesta—Myiagra modesta G. R. Gray, 1859	38
	odesta—Progne modesta Gould, 1839	
	odestum—Dicaeum modestum Tweeddale, 1878	
	odestus—Prionochilus modestus Hume, 1875	
	odestus—Synallaxis modestus Eyton, 1851	
	There is another syntype in the collection. See Hellmayr, Catalogue of Birds of the Americas, 4 (1925): 139.	
1	odestus—Turdus modestus Eyton, 1839	
mo	odestus—Vireo modestus P. L. Sclater, 1860	56
	Proc. zool. Soc. Lond., 1860: 462. SYNTYPE, Adult. Reg. no. 1886.9.15.276. Jamaica. Collected by P. H. Gosse. (Collected by Ward, on label). Sclater Collection. The second syntype is also in the collection.	
	,	62
	Bull. Br. Orn. Cl., 53:7. HOLOTYPE, Adult male. Reg. no. 1938. 3.4.358. Kulikoro, French Sudan (between 12° and 15° North, on label), 9 July 1928. Collected by and purchased of G. L. Bates.	

modulator—Orpheus modulator Gould, 1836
moeritica—Galerida cristata moeritica Nicoll & Bonhote, 1909 . 17 Bull. Br. Orn. Cl., 23:101. Syntype, Adult male. Reg. no. 1909.6.10.1. South shore of Lake Birket-el-Kerun, Fayum, Egypt, 15 March 1909. Collected and presented by M. J. Nicoll and J. L. Bonhote. This is the author's selected male. His selected female syntype is also in the collection.
moesta—Lalage moesta P. L. Sclater, 1883
moesta—Synallaxis moesta P. L. Sclater, 1856
moheliensis—Coracina cinerea moheliensis Benson, 1960 20 Ibis, 103b: 70. Holotype, Adult male. Reg. no. 1959.5.7. Bandamale, 700 metres, Moheli, Comoro Islands, 25 Sept. 1958. Collected by C. W. Benson and presented by the British Ornithologists' Union Centenary Expedition.
moheliensis—Microscelis crassirostris moheliensis Benson, 1960 . 21 Ibis, 103b: 68. Holotype, Adult male. Reg. no. 1959.5.8. Bandamale, 500 metres, Moheli, Comoro Islands, 21 Sept. 1958. Collected by C. W. Benson and presented by the British Ornithologists' Union Centenary Expedition.
moheliensis—Nesillas typica moheliensis Benson, 1960 35 Ibis, 103b:81. Holotype, Adult male. Reg. no. 1959.5.11. Bandamale, 500 metres, Moheli, Comoro Islands, 20 Sept. 1958. Collected by C. W. Benson and presented by the British Ornithologists' Union Centenary Expedition.
moheliensis—Turdus bewsheri meheliensis Benson, 1960 31 Ibis, 103b: 76. Holotype, Adult male. Reg. no. 1959.5.9. Nioumachoua, Moheli, Comoro Islands, 13 Sept. 1958. Collected by C. W. Benson and presented by the British Ornithologists' Union Centenary Expedition.

Bull. Br. Orn. Cl., 10: 62. HOLOTYPE, Adult male. Reg. no. 1889. 5.14.57. Chile, 1875. Collected by Leybold. Salvin-Godman Collection.
mollissima—Chamaeza mollissima P. L. Sclater, 1855 4 Proc. zool. Soc. Lond., 1855 : 89, pl. 95. HOLOTYPE, Adult. Reg. no. 1854.1.25.67. Santa Fé de Bogota, Colombia. Purchased of S. Stevens.
moloneyanus—Turdinus moloneyanus Sharpe, 1892 32B Proc. zool. Soc. Lond., 1892: 228, pl. 20. HOLOTYPE, Adult. Reg. no. 1891.5.25.1. Gold Coast. Collected and presented by Sir A. Moloney.
mombasae—Chlorocichla mombasae Shelley, 1896 21 The Birds of Africa, 1:64. HOLOTYPE, Adult male. Reg. no. 1895.5.1.734. Mombasa, 15 July 1877. Collected by Dr. Fischer. Shelley Collection.
monacha—Pachycephala? monacha G. R. Gray, 1858 39 Proc. zool. Soc. Lond., 1858: 179. HOLOTYPE, Adult male. Reg. no. 1858.3.10.69. Aru Islands. Collected by A. R. Wallace. Wallace Collection.
monachus—Garrulax monachus Swinhoe, 1870
monapo—Erythrocercus livingstonei monapo Vincent, 1933 38 Bull. Brit. Orn. Cl., 53: 137. HOLOTYPE, Adult male. Reg. no. 1933.3.1.15. Near Iamorrimo, 400′, Mozambique Province. 14° 55′ S., 40°25′ E., Portuguese East Africa, 25 May 1932. Collected and presented by J. Vincent.
monarcha—Alcippe monarcha G. R. Gray, 1858 36 Proc. zool. Soc. Lond., 1858: 175. HOLOTYPE, Adult female. Reg. no. 1858.3.10.81. Aru Island. Collected by A. R. Wallace. Purchased of S. Stevens.
monarcha—Sitagra monarcha Sharpe, 1890 61 Cat. Bds. Brit. Mus., 13: 426. Syntype, Adult male. Reg. no. 1889.7.25.51. Fantee, West Africa, 1872. Collected by H. J. Ussher. Shelley Collection. There are other syntypes in the collection.
mongalla—Cisticola ruficeps mongalla Lynes, 1930 35 Ibis, (12) 6, Supplement; 541, pl. 18. HOLOTYPE, Adult male. Reg. no. 1928.8.19.12. Malek, Mongalla Province, Sudan, 27 Dec. 1925. Collected by R. Harwood. Presented by Sir J. Harrington.

- mongolicus—Carnodacus mongolicus Swinhoe, 1870 Proc. zool. Soc. Lond., 1870: 447. HOLOTYPE, Adult. Reg. no. 1898,11.1.489. Keming, near Kalgan, Northern China, 27 Sept. 1868. Collected by R. Swinhoe. Seebohm Collection. monilegera—Cinclosoma monilegera Hodgson, 1836 . 32B Asiat. Res., 19:147. SYNTYPE, Adult (relaxed mount). Reg. no. 1843.1.13.561. Nepal. Collected and presented by B. H. Hodgson. moniliger—Formicarius moniliger P. L. Sclater, 1856 Proc. zool. Soc. Lond., 1856: 294. SYNTYPE, Adult male. Reg. no. 1857.7.30.22. Cordova, Vera Cruz, Mexico. Collected by and purchased of A. Sallé. Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 304. monotonus—Graucalus monotonus Tristram, 1879 Ibis, (4) 3:441. SYNTYPE, Adult male. Reg. no. 1880.2.10.2. Makira Harbour, San Christoval, Solomon Islands, 25 Aug. 1878. Collected by Lt. G. E. Richards, R.N., and purchased of H. B. Tristram. There are other syntypes in the Liverpool Museum. montana—Brachvotervx montana Horsfield, 1821 Trans. Linn. Soc. Lond., 13: 157. SYNTYPE, Adult (relaxed mount). Reg. no. Old Vellum Cat. 13.285a. (Forests of Mount Prahu, 7.000', Java, vide Horsfield, Zoological Researches in Java, 1824.) Collected by T. Horsfield and presented by the Honourable East India Company Museum. montana—Rhipidura montana Mathews, 1928 . 37A Bull. Brit. Orn. Cl., 48:92. HOLOTYPE, Adult. Reg. no. 1898.5. 31.13. Mount Albert Edward, South-east New Guinea. Purchased of Messrs Blyth and Co. montana—Saxicola montana Gould, 1865 . 31 Birds of Asia, 4 (pt. 17): pl. 30. SYNTYPE, Adult. Reg. no. 1898.9.1.2341. Tibet, Collected by W. E. Hay. Seebohm Collection. (See also appendix, Corrections and Additions, vol. 2.) montana—Sylvia montana Horsfield, 1821. 35 Trans. Linn. Soc. Lond., 13:156. HOLOTYPE, Adult. Reg. no. 1860.4.16.394. Java, between 1811 and 1818. Collected by T. Horsfield. Presented by the Honourable East India Company Museum. The type was overlooked by Seebohm, Cat. Bds. Brit. Mus., 5 (1881), but was referred to by Hartert, Novit. Zool., 3 (1896): 538. under the name of Cettia montana. montanella—Linota montanella Henderson and Hume, 1873.
- montanella—Linota montanella Henderson and Hume, 1873. . . 59

 Lahore to Yarkand, : 261 (note), pl. 26. SYNTYPE, Adult male.

 Reg. no. 1887.6.1.330. Arpalak River, Yarkand, (near Sanju, Western Kun Lun, Chinese Turkestan, vide Vaurie, Birds of the

Palearctic Fauna, Passerifomes (1959): 613). Collected by G. Henderson on the Expedition of 1870 under T. D. Forsyth. Hume Collection.

There are two other syntypes in the collection.

- montanorum—Poliospiza gularis montanorum Bannerman, 1923 . 59 Bull. Brit. Orn. Cl., 44: 27. HOLOTYPE, Adult male. Reg. no. 1923.10.26.357. Banso Mountains, 6,000', north of Kumbo, Cameroons, 11 Sept. 1922. Collected by and purchased of G. L. Rates
- montanus—Pomatorhinus montanus Horsfield, 1821 . 32B Trans. Linn. Soc. Lond., 13:165. SYNTYPE, Adult male. Reg. no. 1881.1.1.627. Java. Collected by T. Horsfield and presented by the Honourable East India Company Museum. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 410.

A second, a relaxed mount, since destroyed, was almost certainly another syntype.

montebelli—Anthus australis montebelli Montague, 1913 19 Aust. Av. Rec., 1:181. SYNTYPE, Adult male. Reg. no. 1913.5. 20.14. Hermite Island, Monte Bello Group, West Australia, 9 July 1912. Collected by P. D. Montague and presented by the Royal Society.

There are two other syntypes in the collection.

- 23 Reg. no. 1873.12.10.92. Loando, Rio Dande, Angola, 28 April 1869. Collected by and purchased of J. J. Monteiro.
- chased of J. J. Monteiro.
- Listed as type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 170.
- monticola—Ixos monticola McClelland, 1840 in Horsfield, Proc. zool. Soc. Lond., 1839 (publ. 1840): 160. HOLOTYPE, Adult. Reg. no. 1880.1.1.1906. Kossia mountains, Assam. Collected by J. McClelland and presented by the Honourable East India Company Museum.
- monticola—Newtonia brunneicauda monticola Salomonsen, 1934 . 37 Novit. Zool., 39: 207. HOLOTYPE, Adult male. Reg. no. 1931.8. 18.2849. Manjakatompo, Ankaratra Mountains, Central Madagascar, 12 May 1929. Collected by J. Delacour. Archbold Vernay Expedition, Franco-Anglo-American Expedition.

monticola—Sturnella magna monticola Chubb, 1921. Ann. Mag. nat. Hist., (9) 8:444. HOLOTYPE, Adult mano. 1922.3.5.4555. Mt. Roraima, British Guiana, Collected by F. V. McConnell. McConnell Collection.	57 ale. Reg. Oct. 1898.
monticolus—Cichlops monticolus Hodgson, 1844. in J. E. Gray, Zoological Miscellany, 1844: 83 (nomen Three specimens in the collection were listed as types b Cat. Bds. Brit. Mus., 10 (1885): 567.	nudum).
montium—Myiomela montium Swinhoe, 1864	
montivaga—Calendula magnirostris montivaga Vincent, 1948 Bull. Br. Orn. Cl., 68: 146. HOLOTYPE, Adult male. 1948.39.45. Sanqubetu Valley, 8,000', at junction Lekhalabaletsi and Sanqubetu, 29°16.5' S., 29°21' E., Ba 17 Oct. 1946. Collected and presented by J. Vincent.	Reg. no. of rivers
montpellieri—Pericrocotus montpelleiri La Touche, 1922 Bull. Br. Orn. Cl., 42: 125. HOLOTYPE, Adult male (in adult male by plumage). Reg. no. 1906.12.17.97. Yang (Big Bend), 9-10,000′, Chukung to Yangpi, west Yunnan, 1906. Collected and presented by Col. G. Rippon.	text; sub- gtze River
moori—Alethe moori Boyd Alexander, 1903. Bull. Br. Orn. Cl., 13:37. HOLOTYPE, Adult male. 1936.2.21.47. Near Bakaki, 6,000', Fernando Po, 17 N Collected by Boyd Alexander and presented by his nepher	ov. 1902.
moptana—Saxicola torquata moptana Bates, 1932. Bull. Br. Orn. Cl., 53:8. HOLOTYPE, Adult male. Reg. 8.6.4. Mopti, French Sudan, 13 Jan. 1932. Collected purchased of G. L. Bates.	31 no. 1932. d by and
moreaui—Apalis moreaui W. L. Sclater, 1931 Bull. Brit. Orn. Cl., 51:109. HOLOTYPE, Adult male. 1931.5.7.1. Amani Forest, 3,000', Usambara District, Ta: 14 Dec. 1930. Collected and presented by R. E. Moreau	nganyika,
moreaui—Bradypterus brachypterus moreaui W. L. Sclater, 19 Bull. Brit. Orn. Cl., 52:57. HOLOTYPE, Adult male. 1931.5.7.7. Amani, 3,000', Usambara District, Tangar Nov. 1930. Collected and presented by R. E. Moreau.	Reg. no.
moreaui—Cinnyris moreaui W. L. Sclater, 1933 . Ibis, (13) 3: 214. HOLOTYPE, Adult male. Reg. no. 1931 Maskati, Nguru Mountains, 6,000', Tanganyika, 1931. Collected by R. E. Moreau. Purchased by th Godman Fund.	46 .12.18.42. 14 Sept. e Salvin-

- moreani—Malimbus nitens moreaui C. M. N. White, 1957 Bull. Br. Orn. Cl., 77: 29. HOLOTYPE, Adult male. Reg. no. 1903. 10.26.45. Efulen, Cameroons, 16 Dec. 1902. Collected and presented by G. L. Bates.
- male. Reg. no. 1873.5.12.827. Morty Island, South Pacific. Collected by A. R. Wallace. Wallace Collection.
- morotensis—Piezorhynchus morotensis Sharpe, 1879 38 Cat. Bds. Brit. Mus., 4:423. HOLOTYPE, Adult female. Reg. no. 1873.5.12.327. Morty Island, Moluccas, 1861. Collected by A. R. Wallace Wallace Collection
- morrelli—Pellorneum capistratum morrelli Chasen & Kloss, 1929 . 32B J. Orn., 1929 (2):118. HOLOTYPE, Adult male. Reg. no. 1947. 60.50. Kadat, North Borneo, 19 Sept. 1927. Collected by F. N. Chasen and C. B. Kloss and presented by the Raffles Museum, Singapore.
- morrisonia—Alcippe morrisonia Swinhoe, 1863 . Ibis, (1) 5:296. SYNTYPE, Adult female. Reg. no. 1898.10.3.809. North Formosa, China, April 1862. Collected by R. Swinhoe. Seebohm Bequest. The juvenile, and other adult syntypes are also in the collection.
- morrisoniana—Actinodura morrisoniana Ogilvie-Grant, 1906 Bull. Br. Orn. Cl., 16:119. SYNTYPE, Adult male. Reg. no. 1907.12.12.71. Mount Morrison, 8,000', central Formosa, Jan. 1906. Collected by and purchased of W. Goodfellow. There is also a female syntype in the collection.
- morrisoniana—Suthora morrisoniana Ogilvie-Grant, 1906 Bull. Br. Orn. Cl., 16:119. SYNTYPE, Adult male. Reg. no. 1907.12.12.81. Mount Morrison, 9,000', central Formosa, Jan. 1906. Collected by and purchased of W. Goodfellow.
- morrisonianum—Trochalopterum morrisonianum Ogilvie-Grant, 1906 32B Bull. Br. Orn. Cl., 16:120. SYNTYPE, Adult male. Reg. no. 1907.12.12.47. Mount Morrison, 8,000', central Formosa, Jan. 1906. Collected by and purchased of W. Goodfellow. There is also a female syntype in the collection.
- motacilloides—Pericrocotus motacilloides Swinhoe, 1860 20 Ibis, (1) 2:58. SYNTYPE, Adult male. Reg. no. 1881.5.1.594. Amoy, south China, 23 April 1859. Collected by R. Swinhoe. Gould Collection.
- motacilloides—Rhipidura motacilloides Vigors and Horsfield, 1827 . 37A Trans. Linn. Soc. Lond., 15:248. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.45. George River, Australia. Collected by R. Brown. Presented by the Linnean Society as the type. The tip of the bill is broken.

The type locality, originally stated to be sixty miles south of the Orange River, was restricted to near Hopetown, Cape Province, by Macdonald, Contributions to the Ornithology of Western South Africa, (1957): 156-7; but subsequently revised to Kurumen by Clancey, Bull. Br. Orn. Cl., 84 (1964): 110. The female syntype is in the collection.

The specimen appears to be the type but has lost its feet; the label, which had become detached, was found loose in the same drawer.

- moultoni—Chlorocharis emiliae moultoni Chasen and Kloss, 1927. 47 Bull. Brit. Orn. Cl., 48:47. Holotype, Adult male. Reg. no. 1935.12.27.1. Mount Poi, 5,000', south-western Sarawak, 6 Oct. 1923. (Nov. on label). Collected by E. Mjöberg. Presented by the Sarawak Museum.
- moultoni—Ophrydornis albogularis moultoni Robinson & Kloss, 1919 32B Bull. Br. Orn. Cl., 40:17. HOLOTYPE, Adult female. Reg. no. 1927.4.18.995. Betong Saribas, south-west Sarawak, Borneo, 8 Aug. 1916. Collected by natives and presented by H. C. Robinson.
- mpangae—Cinnamopteryx mpangae Ogilvie-Grant, 1907 . . . 61

 Bull. Br. Orn. Cl., 21:15. HOLOTYPE, Adult male. Reg. no. 1906.12.23.124. Mpanga forest, 5,000′, Fort Portal, Uganda, 20 Sept. 1906. Collected by R. B. Woosman. Presented by the subscribers to the Ruwenzori Expedition.
- muelleri—Cisticola muelleri Boyd Alexander, 1899 35
 Bull. Brit. Orn. Cl., 8:49. HOLOTYPE, Adult female. Reg. no.

- 11.12.23.2526. Zambesi River, (Mesanangue, 20 Jan. 1899, on label). Collected by W. Boyd Alexander. Boyd Alexander Bequest.
- muelleri—Cryptolopha muelleri Robinson and Kloss, 1916. J. Straits Branch, Roy. Asiat. Soc., 73: 278. Holotype, Adult male. Reg. no. 1920.6.29.252. Barong Bharu, 4,000', Barison Range, Korinchi, West Sumatra, 8 June 1914. Collected and presented by H. C. Robinson and C. B. Kloss.
- mufumbiri—Laniarius mufumbiri Ogilvie-Grant, 1911 . Bull. Br. Orn. Cl., 29: 30. HOLOTYPE, Adult. Reg. no. 1911.11. 28.15. Mufumbiro Volcanoes (Vichumbi, south end of Lake Edward, East Africa, 4 Dec. 1910, vide Ogilvie-Grant, Ibis, (9) 6 (1912): 332-4). Collected and presented by Mrs M. Roby.
- muhuluensis—Apalis melanocephala muhuluensis C. H. B. Grant and 35 Tanganyika, 30 Jan. 1946. Collected by and purchased of R. E. Moreau.
- multicolor—Laniarius multicolor G. R. Gray & Mitchell, 1849. The Genera of Birds, 1:299, pl. 72. HOLOTYPE, (Adult male, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883): 159). Reg. no. Old Vellum Cat. 10.15. (West Africa on label.) Received from Capt. Sabine.
- multipunctata—Nucifraga multipunctata Gould, 1849. Proc. zool. Soc. Lond., 1849: 3. The specimen described was in the museum of the Philosophical Society of Yorkshire. The specimen listed by Sharpe, Cat. Bds. Brit. Mus., 3 (1877): 56 as the type cannot be accepted.
- multostriata—Myrmotherula multostriata P. L. Sclater, 1858. Proc. zool. Soc. Lond., 1858: 234, pl. 141. SYNTYPE, Adult male. Reg. no. 1855.12.17.48. Ucayali River, Upper Amazon. Collected by Hauxwell and purchased of J. Gould. The female and immature syntypes are in the collection.
- munipurensis—Cisticola munipurensis Godwin-Austen, 1874. Proc. zool. Soc. Lond., 1874: 47. SYNTYPE, Adult. Reg. no. 1895.7.14.2014. Dunsiri Valley, Naga Hills, Munipur. Collected and presented by H. H. Godwin-Austen. There are two other syntypes in the collection.
- Ibis, (3) 5:250. SYNTYPE, Adult. Reg. no. 1895.7.14.1515. Karakhul, Munipur. Collected by W. Robert. Godwin-Austen Collection.

There are three other syntypes in the collection.

- muraria—Prunella rubeculoides muraria R. Meinertzhagen, 1926. 30 Bull. Brit. Orn. Cl., 46:99. HOLOTYPE, Adult female. Reg. no. 1897.12.10.1123. Astor, Kashmir, 21 Nov. 1876. Collected and presented by J. Biddulph.
- murielae—Saxicola dacotiae murielae Bannerman, 1913 . . . 31 Bull. Br. Orn. Cl., 33: 37 (figured Ibis, (10) 2 (1914): pl. 5). Syntype, Adult male. Reg. no. 1913.10.22.105. Allegranza Island, Canary Islands, 12 June 1913. Collected by and purchased of D. A. Bannerman

This is the author's selected male. His selected female and ten other syntypes are also in the collection.

Listed as the type by Sharpe, *Cat. Bds. Brit. Mus.*, 12 (1888): 453. There are five other syntypes in the collection.

- murinus—Thanmophilus murinus P. L. Sclater & Salvin, 1867.

 Proc. zool. Soc. Lond., 1867: 756. SYNTYPE, Adult male. Reg. no. 1889.9.20.63. Barra do Rio Negro (= Manaos, vide Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 88), Brazil, 15 Oct. 1830. Collected by J. Natterer. Sclater Collection.

 There is a female syntype in the collection.
- muscipetoides—Chaptia muscipetoides Hodgson, 1837 . . . 64

 The Indian Review, 1 (1836, publ. 1837): 325. ?SYNTYPE, Adult.

 Reg. no. 1843.1.13.711. Nepal. Collected and presented by B. H. Hodgson.

Received with his main collection, the specimen has Hodgson's drawing number on the label and is therefore likely to have been before him when the description was made.

musicapinus—Hylophilus musicapinus P. L. Sclater and Salvin, 1873 56
Nomencl. Av. Neotrop.: 156. HOLOTYPE, Adult. Reg. no. 1886.
9.15.293. St Louis d'Oyapok, French Guiana, (3 March 1867, on label). Collected by Jelski (vide Sclater, Ibis, (4) 5 (1881): 290).
Sclater Collection; obtained by him through Verreaux.

- musicus—Pomatorhinus musicus Swinhoe, 1859 32B J. North China Branch, Asiat. Soc., 2:228. SYNTYPE, Adult. Reg. no. 1898.10.3.588. Formosa, 21 June 1858. Collected by R. Swinhoe. Seebohm Collection.

 There are several other syntypes in the collection.
- mussooriensis—Aethopyga siparaja mussooriensis E. C. S. Baker, 1925 46
 Bull. Brit. Orn. Cl., 46: 12. HOLOTYPE, Adult male. Reg. no.
 1886.12.1.480. Mussoorie, North-west India, 1872. Collected by
 A. O. Hume. Hume Collection.
- mysolensis—Calornis mysolensis G. R. Gray, 1861 62

 Proc. zool. Soc. Lond., 1861: 431. HOLOTYPE, Adult (male, vide Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 150. Reg. no. 1873.5.12. 1902. Mysol, 1860. Collected by A. R. Wallace. Wallace Collection.

There are several other syntypes in the collection.

Listed as type by Sclater, *Catalogue of American Birds*, (1862): 22. The second syntype is in the collection.

- nagaensis—Turdimus nagaensis Godwin-Austen, 1877 32B Ann. Mag. nat. Hist., (4) 20: 519. HOLOTYPE, Adult female. Reg. no. 1895.7.14.1255. East Naga Hills, north-east Assam, 17 April 1876. Collected by A. W. Chennell and presented by H. H. Godwin-Austen.
- namaqua—Cisticola subruficapilla namaqua Lynes, 1930 . . . 35

 Ibis, (12) 6, Supplement: 216. HOLOTYPE, Adult female. Reg.
 no. 1905.12.29.2123. Klipfontein, Little Namaqualand, 11 April
 1903. Collected by C. H. B. Grant. Presented by C. D. Rudd.
- namaquensis—Bradornis infuscata namaquensis Macdonald, 1957 37 Contribution to the ornithology of Western South Africa: 119. HOLOTYPE, Adult female. Reg. no. 1887.12.1.1332. (not 1889, as stated). Aamhoup, Great Namaqualand, 1 July 1861. Collected by C. J. Andersson. Tweeddale Bequest.
- namaquensis—Cossypha caffra namaquensis W. L. Sclater, 1911 . 31 Ibis, (9) 5:415. HOLOTYPE, Adult female. Reg. no. 1905.12.29. 867. Klipfontein, Little Namaqualand, 19 June 1903. Collected by C. H. B. Grant and presented by C. D. Rudd
- namaquensis—Karrucincla schegelii namaquensis W. L. Sclater, 1928 31 Bull. Br. Orn. Cl., 49:15. HOLOTYPE, Adult male. Reg. no. 1866.7.19.5. Great Namaqualand, 27 June 1861. Collected by C. J. Andersson and purchased of Stevens.

This name is based on the specimens described as *Emarginata* cinerea (Vieillot) by Sharpe, *Ibis*, (8) 4 (1904): 325 and Reichenow, *Die Vögel Afrikas*, 3 (1905): 710.

- namwera—Cinnyris chalybeus namwera Vincent, 1933 46
 Bull. Brit. Orn. Cl., 53:146. HOLOTYPE, Adult male. Reg. no.
 1933.3.1.8. Mangoche Mountains, 3,700', near Namwera's, Nyasaland, 6 May 1932. Collected by and purchased of J. Vincent.

nana—Acanthiza nana, Vigors and Horsfield, 1827
Trans. Linn. Soc. Lond., 15:226. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.25. Australia. Collected by G. Caley. Presented as the type by the Linnean Society of London. The specimen has now lost a wing.
nana—Seisura nana Gould, 1870
nandensis—Dryoscopus nandensis Sharpe, 1900 23 Bull. Br. Orn. Cl., 11:28. HOLOTYPE, Adult male. Reg. no. 1900.6.22.2. Nandi, 6,500', Equatorial Africa, 29 June 1892. Collected and presented by F. J. Jackson.
nandensis—Sycobrotus nandensis Jackson, 1899 61 Ibis, (7) 5:615. HOLOTYPE, Adult female. Reg. no. 1899.8.13.4. Nandi, 6,000', British East Africa, 10 April 1898. Collected and presented by F. J. Jackson.
nanus—Pyrocephalus nanus Gould, 1839
napensis—Cercomacra napensis P. L. Sclater, 1868 4 Proc. zool. Soc. Lond., 1868: 572. SYNTYPE, Adult male. Reg. no. 1889.9.20.405. Rio Napo, eastern Ecuador. Collected for Verreaux. Sclater Collection. Listed as type by Hellmayr, Novit. Zool., 12 (1905): 287. There is another syntype in the collection.
natalensis—Camaroptera natalensis Hartlaub, 1863 35 in Gurney, Ibis, (1) 5:326, pl. 8. SYNTYPE, Adult (female, vide Lynes, Ibis, (12) 6 (1930) supplement: 651). Reg. no. 1890.12.20.45. Natal. Collected by T. Ayres. Presented by J. H. Gurney Inr.
natalensis—Cossypha natalensis Smith, 1840

TYPE, "Possibly immature male" (relaxed mount). Reg. no. 1843. 2.28.36. Port Natal. Collected by A. Smith and purchased

of the South African Society.

natalensis—Nectarinia natalensis Jardine, 1843
natalensis—Sphenoeacus natalensis Shelley, 1882 35 Proc. zool. Soc. Lond., 1882: 337. SYNTYPE, Adult male. Reg. no. 1895.5.1.971. Newcastle, Natal. Collected by A. E. Butler. Shelley Collection.
natalis—Zosterops natalis Lister, 1889
nationi—Buarremon nationi P. L. Sclater, 1881
nationi—Myiobius nationi P. L. Sclater 1866
natronensis—Prinia gracilis natronensis Nicoll, 1917
nattereri—Anthus nattereri P. L. Sclater, 1878
nattereri—Pipra nattereri P. L. Sclater, 1865

- natunensis—Anthreptes macularia natunensis Chasen, 1935 . . . 46
 Bull. Raffles Mus., 10:43. Holotype, Adult male. Reg. no.
 60.64. Bunguran, North Natuna Islands, Malaya, 28 Aug. 1928.
 Collected by F. N. Chasen. Presented by the Raffles Museum, Singapore.
- natunensis—Charitocirus maculatus natunensis Chasen, 1935. . . . 45
 Bull. Raffles Mus., 9: 95. Holotype, Adult male. Reg. no. 1947.
 60.64. Bunguran, North Natuna Islands, China Seas, 28 Aug.
 1928. Collected by F. N. Chasen. Presented by the Raffles Museum, Singapore.
- natunensis—Chloropsis cochinchinensis natunensis Chasen, 1938 . 22 Orn. Mber., 46:7. HOLOTYPE, Adult male. Reg. no. 1948.23.1. Bunguran Island, North Natuna Islands, 11 Sept. 1928. Collected by F. N. Chasen and presented by the Raffles Museum, Singapore.
- nayaritensis—Aimophila acuminata nayaritensis van Rossem, 1938. 49
 Bull. Brit. Orn. Cl., 58: 128. HOLOTYPE, Adult male. Reg. no. 1899.2.1.3675. Tepic, Nayarit, Mexico, 8 June 1889. Collected by W. B. Richardson. Salvin-Godman Collection.
- nayaritensis—Icterus graduacauda nayaritensis van Rossem, 1938. 57

 Bull. Brit. Orn. Cl., 58:137. HOLOTYPE, Adult male. Reg. no. 1898.12.14.1177. Tepic, Nayarit, Mexico, 29 May 1889. Collected by W. B. Richardson. Salvin-Godman Collection.
- nebularum—Saxicola torquata nebularum Bates, 1930 31 Bull. Br. Orn. Cl., 51:51. HOLOTYPE, Adult male. Reg. no. 1930.12.3.5. Near Birwa Peak, 4,500′, Timgi Hills, Kono District, Sierra Leone, 13 July 1930. Collected by and purchased of G. L. Bates.
- neergaardi—Cinnyris neergaardi C. H. B. Grant, 1908 46
 Bull. Brit. Orn. Cl., 21:93. HOLOTYPE, Adult male. Reg. no. 1905.12.29.1642. Coguno, Inhambane District, Portuguese East Africa, 5 Sept. 1906. Collected by C. H. B. Grant. Presented by C. D. Rudd.

neglecta—Calamocichla neglecta Boyd Alexander, 1908 35 Bull. Brit. Orn. Cl., 23:33. Syntype, Adult male. Reg. no. 1911.12.23.2857. Lake Chad, French Equatorial Africa, 5 Dec. 1904. Collected by W. Boyd Alexander. Boyd Alexander Bequest. The female syntype is also in the collection.
neglecta—Calornis neglecta Walden, 1872
neglecta—Chlorodyta neglecta Boyd Alexander, 1899 35 Bull. Brit. Orn. Cl., 10:17. SYNTYPE, Adult (male, on label). Reg. no. 1911.12.23.2196. Zambesi River, (near Zumba, 12 Nov. 1898, on label). Collected by W. Boyd Alexander. Boyd Alexander Bequest. This specimen is labelled as the type, but other syntypes are implied by the description, and one of these is also in the collection.
neglecta—Gerygone neglecta Wallace, 1865
neglecta—Pachycephala neglecta Layard, 1879
neglecta—Sitta neglecta Walden, 1870
neglecta—Volvocivora neglecta Hume, 1877
neglecta—Zosterops neglecta Seebohm, 1893

neglectus—Calamonastes simplex neglectus Benson, 1936 Bull. Brit. Orn. Cl., 56:71. HOLOTYPE, Adult male. Reg. no. 1935.9.11.4. 5 miles East of Fort Hill, 4,300', north-west Nyasaland, 17 Aug. 1935. Collected and presented by C. W. Benson. neglectus—Pericrocotus neglectus Hume, 1877 Stray Feathers, 5: 171, 189. SYNTYPE, Adult male. Reg. no. 1886.4.1.708. Meetan, central Tenasserim, 25 Feb. 1877. Collected by W. Davison. Hume Collection. There is also a female syntype in the collection. (Bhawulpore, Punjab, 30 Jan. 1868, on label), Punjab and Doab, India. Collected by C. H. T. Marshall. Hume Collection. This is the only specimen in the collection obtained before the date of description, but Hume refers to specimens of both sexes. neilgherriensis—Hypsipetes neilgherriensis Jerdon, 1839 Madras J. Lit. Sci., 25: 245. SYNTYPE, Adult. Reg. no. 1845.1. 10.14. Summit of Nilgiri Hills (in text & Madras on label). Collected and presented by T. C. Jerdon. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 40. nelsoni-Geothlypis nelsoni Richmond, 1900 Auk, 17: 179. New name for G. cucullata Salvin and Godman. 1889 (a.v.) on the grounds of preoccupation by Sylvia cucullata Latham 1790. nemoricola—Fringalanda nemoricola Hodgson, 1836 . Asiat. Res., 19:158. SYNTYPE, Adult. Reg. no. 1843.1.13.877. Nepal. Collected and presented by B. H. Hodgson. There are other syntypes in the collection. nesa—Pyrrhula pyrrhula nesa Mathews and Iredale, 1917 Austral. Avian Record, 3:122. There is no evidence that actual specimens were examined when the original description was prepared. Meinertzhagen, Bull. Brit. Orn. Cl., 68 (1947): 32, claims that Iredale briefly examined his specimens and therefore designates as lectotypes a pair (reg. nos. 1965.M.55 and 56) from Tring. Hertfordshire collected on 10 Dec. 1909. Since it cannot be satisfactorily established that these birds were original syntypes these have not been accepted as lectotypes. neseri—Oenanthe pileata neseri Macdonald, 1952 Ostrich, 23:161. HOLOTYPE, Adult male. Reg. no. 1938.10.17.8. Erongo Mountains, Omaruru District, South West Africa, 16 June 1938. Collected by and purchased of W. Hoesch. nesophilus—Cimyris nesophilus Shelley, 1892 . . . Bull. Brit. Orn. Cl., 1:5. SYNTYPE, Adult female. Reg. no.

- 1895.9.9.32. Angasija, Great Comora Island. Collected by J. Kirk. Shelley Collection.

 A second syntype is also in the collection.
- neumanni—Amydrus neumanni Boyd Alexander, 1908 62

 Bull. Brit. orn. Cl., 23:41. HOLOTYPE, Adult male. Reg. no. 1911.12.23.3731. Petti, on Kaduna River, northern Nigeria, 27

 Aug. 1904. Collected and bequeathed by Boyd Alexander.
- neumanni—Hypochera neumanni Boyd Alexander, 1908 . . . 61

 Bull. Brit. Orn. Cl., 25:33. Holotype, Adult male. Reg. no. 1911.12.23.3306. Yo, Lake Chad, 19 Nov. 1904. Collected by Boyd Alexander. Boyd Alexander Bequest.
- neumanni—Othyphantes baglafecht neumanni Bannerman, 1923. 61
 Bull. Br. Orn. Cl., 43:165. Holotype, Adult male. Reg. no. 1922.11.25.347. Banso Mountains, 6,000′, north of Kumbo, Cameroons, 23 Sept. 1921. Collected by and purchased of G. L. Bates.

This specimen is the one figured. There is also a male syntype in the collection. Two more syntypes are in the University of Cambridge Museum of Zoology, and seven in the Leiden Museum.

- newtoni—Riparia paludicola newtoni Bannerman, 1937 18

 Bull. Br. Orn. Cl., 58:31. HOLOTYPE, Adult male. Reg. no. 1926.8.8.195. Bambulue, 4,800′, near Bamenda, Cameroons, 1 Feb. 1925. Collected by and purchased of G. L. Bates.
- ngamii—Pycnonotus tricolor ngamii Ogilvie-Grant, 1912 . . . 21 Ibis, (9) 6: 391. HOLOTYPE, Adult female. Reg. no. 1911.12.16.51. Lake Ngami, 2,700′, Bechuanaland, 2 July 1909. Collected by G. Legge on the Woosnam-Legge Expedition.
- nicaraguensis—Quiscalus nicaraguensis Salvin and Godman, 1891 57

 Ibis, (6) 3:612. Syntype, Adult male. Reg. no. 1898.12.14.2334.

 Momotombo, Lake Managua, Nicaragua, 21 May 1891. Collected by W. B. Richardson. Salvin—Godman Collection.

 Listed as the type by Hellmayr and Conover, Catalogue of a collection of American birds 10 (1937):94. There are several other syntypes in the collection.

nicholsoni—Anthus nicholsoni Sharpe, 1884
nicobarica—Aethopyga nicobarica Hume, 1873 4 Stray Feathers, 1:412. SYNTYPE, Adult male. Reg. no. 1886. 12.1.267. Kondul, Nocobar Islands, Indian Ocean, (14 March 1873, collected by W. Davison, on label). Hume Collection. This specimen was marked as the type in the British Museum Register. There are seven other syntypes in the collection.
nicobarica—Terpsiphone nicobarica Oates, 1890 3 Fauna of British India, 2: 48. SYNTYPE, Adult male. Reg. no. 1887.12.30.1034. (Trinkut, Nicobar Islands, Jan. 1874, on label). Collected by R. I. Wimberley. Tweeddale Collection. The specimen is labelled "type" in an early hand, and from the preface to the first volume it is clear that the author was using the British Museum Collection.
nicobariensis—Dissemurus paradiseus nicobariensis E. C. S. Baker, 1918
nicolli—Ploceus (Symplectes) nicolli W. L. Sclater, 1931 6 Bull. Br. Orn. Cl., 52:26. HOLOTYPE, Adult male. Reg. no. 1931.5.7.6. Amani, 3,000', Tanganyika, 19 June 1931. Collected and presented by R. E. Moreau.
nigellicauda—Psaropholus ardens var nigellicauda Swinhoe, 1870 6 Ibis, (3) 1:342. Syntype, Adult male. Reg. no. 1897.6.1.482. Tainingsze Central Hainan 20 Feb. 1868 Collected by R

Swinhoe. Seebohm Collection.

The second syntype is also in the collection.

cription is of an adult and according to Stonehouse, Scottish Naturalist, (1930): 150, the type is in the Edinburgh Museum. The specimen listed by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 510, is immature and cannot be accepted as the type.

niger—Copsychus niger R. G. W. Ramsay, 1886. Proc. zool. Soc. Lond., 1886: 123. SYNTYPE, Adult male. Reg. no. 1888.4.20.97. Elopura, north-east Borneo, April 1884. Collected by H. Pryer. Tweeddale Collection.

niger—Xenops niger Pelzeln, 1859
nigeriae—Anthreptes collaris nigeriae C. M. N. White, 1950 46 Bull. Brit. Orn. Cl., 70: 43. HOLOTYPE, Adult female. Reg. no. 1949.33.71. Owerri, south-east Nigeria, 25 Jan. 1949. Collected and presented by S. Marchant.
nigeriae—Emberiza affinis nigeriae Bannerman and G. L. Bates, 1926
nigeriae—Erythrocercus maccalli nigeriae Bannerman, 1920 38 Bull. Brit. Orn. Cl., 41:5. HOLOTYPE, Adult male. Reg. no. 1920.3.10.2. Iju Waterworks, near Lagos, Southern Nigeria, 9 Feb. 1920. Collected by W. P. Lowe on a voyage of H.M.S. Dwarf. Presented by Capt. Dane, R.N.
nigeriae—Hypochera nigeriae Boyd Alexander, 1908 61 Bull. Brit. Orn. Cl., 23:15. Holotype, Adult male. Reg. no. 1911.12.23.3302. Kiri, Gongola River, Northern Nigeria, 22 July 1904. Collected by Boyd Alexander on the Alexander-Gosling Expedition. Boyd Alexander Bequest.
nigeriae—Malimbus rubricollis nigeriae Bannerman, 1921 61 Bull. Br. Orn. Cl., 41:77. HOLOTYPE, Adult male. Reg. no. 1920.6.8.202. Iju Waterworks, near Lagos, Nigeria, 31 Dec. 1919. Collected by W. P. Lowe during a voyage of H.M.S. Dwarf, and presented by Capt. Dane RN.
nigeriae—Sylviella flaviventris nigeriae Bannerman, 1920 35 Bull. Brit. Orn. Cl., 41: 4. HOLOTYPE, Adult ?male. Reg. no. 1920.3.10.3. Iju Waterworks, near Lagos, South Nigeria, 29 Dec. 1929. Collected by W. P. Lowe on the voyage of H.M.S. Dwarf, and presented by him.
nigerrima—Hypochera nigerrima Sharpe, 1871
nigerrima—Hypsipetes nigerrima Gould, 1862

- lected by W. Osburn. Sclater Collection.

 Purchased as the type, but Sclater, *Ibis*, (1) 1 (1859): 457, refers to several specimens which are all syntypes. nigerrimus—Myrmelastes nigerrimus P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858: 275. HOLOTYPE, Adult male. no. 1858.4.27.49. Rio Javarri, Upper Amazon, Brazil. Reg. lected by H. W. Bates and purchased of S. Stevens. nigra—Cittocincla nigra Sharpe, 1877. . 31 gra—Cittocincla nigra Sharpe, 18//.

 Trans. Linn. Soc. Lond. (Zool.), (2) 1:335, pl. 52. SYNTYPE,

 Adult male. Reg. no. 1876.8.28.14. Puerta Princesa, Palawan, Philippines, July 1874. Collected by and purchased of J. B. Steere. nigrescens—Chamaeospiza nigrescens Salvin and Godman, 1889 Ibis, (6) 1:381. SYNTYPE, Adult male. Reg. no. 1899.2.1.3854. Patzcuaro, Michoacan, Mexico, Jan. 1888. Collected by F. D. Godman. Salvin-Godman Collection. The female syntype is also in the collection. nigrescens—Chlorophoneus nigrescens W. L. Sclater, 1931 Bull. Br. Orn. Cl., 51:110. HOLOTYPE, Adult male. Reg. no. 1931.5.7.4. Amani Forest, 3,000', Usambara District, Tanganyika, 19 March 1931. Collected and presented by R. E. Moreau. Reg. no. 1882.1.20.799. Kyeikpadein, near Pegu, Tenasserim, 27 April 1876. Collected by and purchased of E. W. Oates. nigrescens—Euprinodes nigrescens Jackson, 1906. grescens—Euprinodes nigrescens Jackson, 1906.

 Bull. Brit. Orn. Cl., 16:90. HOLOTYPE, Adult. Reg. no. 1906.
 6.3.3. Ruwenzori, (probably Ankole, vide Ogilvie-Grant, Trans. zool. Soc. Lond., 19 (1910): 253), 8 April 1902. Collected and presented by F. J. Jackson. nigrescens—Hypsipetes psaroides nigrescens E. C. S. Baker, 1917 . 21 Bull. Br. Orn. Cl., 38: 15. SYNTYPE, Adult male. Reg. no. 1909. 4.7.17. Kalaikhin, Pondaung, Upper Chindwin, Burma, 9 Feb. 1905. Collected by Mears and presented by C. H. Hopwood. This is the author's selected male. His selected female syntype is also in the collection.
- nigrescens—Myiochanes nigrescens P. L. Sclater and Salvin, 1880 . 9

 Proc. zool. Soc. Lond., 1880: 157. SYNTYPE, Adult. Reg. no.
 1888.1.1.1492. Sarayacu, Ecuador. Collected by C. Buckley.
 Salvin-Godman Collection.

There are several other syntypes in the collection.

mgneam corcomacia mgneam in 21 someth, 1000	4
Proc. zool. Soc. Lond., 1858: 245. SYNTYPE, Adult male. Reg.	
no. 1889.9.20.425. Santa Marta, Colombia. Collected for	
Verreaux. Sclater Collection.	
Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 267.	
nigricapillus—Lanius collurioides nigricapillus. See appendix, Correc-	
tions and Additions, vol. 2	3
nigricapillus—Thryothorus nigricapillus P. L. Sclater, 1860 2	R
Proc. zool. Soc. Lond., 1860: 84. SYNTYPE, Adult (female, vide	
Sclater, Catalogue of American Birds, (1862): 21). Reg. no. 1886.	
9.15.121. Nanegal, 4,000', west slope of Pichincha, Ecuador,	
Feb. 1859. Collected by L. Fraser. Sclater Collection.	
Listed as type by Sclater, 1862, <i>loc. cit</i> . The two other syntypes	
are also in the collection.	
nigricauda—Laniarius quadricolor nigricauda Clarke, 1913 2.	3
Bull. Br. Orn. Cl., 31:32. HOLOTYPE, Adult male. Reg. no.	J
1903.8.1.34. Takaungu, coast of Kenya, 15 March 1901. Col-	
lected and presented by A. B. Percival.	
	4
	4
Biol. CentrAmer., 2:230. HOLOTYPE, Adult male. Reg. no.	
1889.7.10.634. Intac, Ecuador, Dec. 1877. Collected by C.	
Buckley. Salvin-Godman Collection.	
nigriceps—Campylorhynchus nigriceps P. L. Sclater, 1860	8
Proc. zool. Soc. Lond., 1860: 461. SYNTYPE, Immature (juvenile,	
vide Hellmayr, Catalogue of Birds of the Americas, 7 (1934): 138).	
Reg. no. 1886.9.15.24. Jalapa, Vera Cruz, Mexico. Collected by	
de Oca. Sclater Collection.	
The second syntype cannot be traced in the collection.	
nigriceps—Cyanocorax nigriceps P. L. Sclater and Salvin, 1876 . 7	1
Proc. zool. Soc. Lond., 1876: 354. SYNTYPE, Adult. Reg. no.	
1887.3.9.35. Tilotilo, Province of Yungas, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.	
C. Buckley. Salvin-Godman Collection.	
A second syntype is also in the collection.	
nigriceps—Dryodromas nigriceps Shelley, 1873	5
Ibis, (3) 3:139. HOLOTYPE, Adult. Reg. no. 1895.5.1.1088.	
Abouri, Aguapim, (Gold Coast), West Africa, 22 Feb. 1872. Col-	
lected by G. E. Shelley. Shelley Collection.	
inglife po 1/2 / tell ellino illo. Itelpo 2 i 21 22 i 22 i 22 i 22 i 23 i 23 i 24 i 25	9
Proc. zool. Soc. Lond., 1860:68. SYNTYPE, Adult male. Reg.	
no. 1888.1.13.1122. Pallantanga, Ecuador, (Oct. 1858, on label).	
Collected by L. Fraser. Sclater Collection.	
Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 258.	
There is another syntype in the collection.	
nigriceps—Orthotomus nigriceps Tweeddale, 1878	5
Proc. zool. Soc. Lond., 1877 (publ. 1878): 828, pl. 85. HOLOTYPE,	
Adult male, (female, on label). Butuan, Mindanao, Philippines,	
May 1877. Collected by A. Everett. Tweeddale Collection.	

- nigriceps—Sibia nigriceps Hodgson, 1839 . . 32B J. Asiat. Soc. Bengal, 8:38. SYNTYPE, Adult. Reg. no. 1843. 1.13.616. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection. nigriceps—Stachyris nigriceps Blyth, 1844 . J. Asiat. Soc. Bengal, 13: 378. Two specimens in the collection were listed as types by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 533. but the material used by Blyth is presumably still at the Calcutta Museum. Stachyris nigriceps J. E. Gray, Zoological Miscellany, 1844: 82, is a nomen nudum. Both authors used Hodgson's MS. nigriceps—Thanmophilus nigriceps P. L. Sclater, 1868 Proc. zool. Soc. Lond., 1868: 571. HOLOTYPE, Adult. Reg. no. 1889.9.20.58. Bogota, Colombia. Obtained by Sclater from O. Salvin. Sclater Collection. nigriceps—Todirostrum nigriceps P. L. Sclater 1855 Proc. zool. Soc. Lond., 1855: 66, pl. 84. HOLOTYPE, Adult. Reg. no. 1888.1.13.243. Santa Marta, Colombia. Collected for Verreaux. Sclater Collection. nigriceps—Xenocichla nigriceps Shelley, 1889 Proc. zool. Soc. Lond., 1889: 362. SYNTYPE, Adult male. Reg. no. 1889.3.21.180. Kilimanjaro, 6,000', East Africa, 9 Aug. 1888. Collected and presented by H. C. V. Hunter. The female syntype is also in the collection. nigricinereus—Parus nigricinereus Jackson, 1898 Bull. Brit. Orn. Cl., 8:22. SYNTYPE, Adult male. Reg. no. 1898. 13.3. (Collector's no. 598). Nandi, Kenya. 10 April 1898. Collected and presented by F. J. Jackson. This specimen was presented as the type but the author refers to three other specimens in *Ibis*, (7) 5 (1899): 638. These are not in the collection. nigricotis—Motacilla nigricotis Shelley, 1900 Birds of Africa, 2:266, pl. 12. HOLOTYPE, Adult male. Reg. no.
- 1895.5.1.2028. North of Limpopo, Zambesi. Collected by Bradshaw. Shelley Collection.
- nigrifrons—Enicurus nigrifrons G. R. Gray, 1859. 31 Proc. zool. Soc. Lond., 1859: 102. SYNTYPE, Adult. Reg. no. 1859.3.4.315. Nepal. Collected and presented by B. H. Hodgson. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7(1883): 323. Gray used Hodgson's MS, name.
- nigrifrons—Formicarius nigrifrons Gould, 1855 Ann. Mag. nat. Hist., (2) 15:344. HOLOTYPE, Adult female. Reg. no. 1854.12.20.16. Chamicuros, River Ucayali, east Peru. lected by Hauxwell and purchased of J. Gould. Col-

- nigrifrons—Leptopogon nigrifrons Salvin and Godman, 1884 . . . 9

 Ibis, (5) 2: 446. HOLOTYPE, Adult male. Reg. no. 1888.1.1.565.

 Roraima, 5,000', British Guiana, 30 Aug. 1883. Collected by H. Whitely. Salvin-Godman Collection.

This is the described male syntype. The two female syntypes are also in the collection.

nigrimenta—Yuhina (Polyodon) nigrimenta Hodgson, 1845. . . 32B J. Asiat. Soc. Bengal, 14:562. HOLOTYPE or SYNTYPE, Adult (relaxed mount). Reg. no. 1843.1.13.362. Nepal. Collected and presented by B. H. Hodgson.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 634.

nigrimentalis—Stoparola nigrimentalis Ogilvie-Grant, 1894 . . . 37

Bull. Brit. Orn. Cl., 3:50. Syntype, Adult male. Reg. no. 1897.

5.13.71. North-west Luzon, Philippines, (La Trinidad, Benguet District, 13 Jan. 1894, on label). Collected and presented by J. Whitehead.

Listed as the type by Hachisuka, *Birds of the Philippines*, 2 (1935): 344. The female syntype is also in the collection.

nigrimentum—Monarcha nigrimentum G. R. Gray, 1860 . . . 38 Proc. zool. Soc. Lond., 1860: 352. HOLOTYPE, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 418). Reg. no. 1873. 5.12.254. Amboyna, Moluccas, 1857. Collected by A. R. Wallace. Wallace Collection.

Of three specimens which might have been available this is the only one which has the black colour under the chin on which the description was based.

nigripennis—Oriolus (Baruffus) nigripennis J. and E. Verreaux, 1855 63 J. Orn. 1855: 105. SYNTYPE, Adult. Reg. no. 1854.5.23.23. Gaboon River, West Africa. Collected by and purchased of J. and E. Verreaux.

There is another syntype in the collection.

- nigrita—Hirundo nigrita G. R. Gray & Mitchell, 1845 18

 The Genera of Birds, 1:58, pl. 20. Holotype, Adult. Reg. no. 1843.3.31.2. River Niger, West Africa. Collected by T. H. Thomson or L. Fraser on the 1841–42 expedition of H.M.S. Wilberforce and presented by Mrs Heywood and Capt. Allen.

- nigrita—Ochthoeca nigrita P. L. Sclater and Salvin, 1871 . . . 9

 Proc. zool. Soc. Lond., 1870 (publ. 1871): 787. HOLOTYPE, Adult male. Reg. no. 1888.1.13.78. Merida, Venezuela, July 1869. Collected by A. Goering. Sclater Collection.

- nigroaurita—Leioptila pulchella nigroaurita Kinnear, 1944 . . . 32B Ibis, 86:83. HOLOTYPE, Adult male. Reg. no. 1938.12.13.241. Lhalung, 7,000', Pachakshiri, south-east Tibet, 29 April 1938. Collected and presented by F. Ludlow and G. Sherriff.
- nigrocapitata—Brachypteryx nigrocapitata Eyton, 1839 32B Proc. zool. Soc. Lond., 1839 : 103. HOLOTYPE, Adult. Reg. no. 1881.2.18.29. Malacca. Collected by Evans. Eyton Collection.
- nigrocapitatus—Mixornis nigrocapitatus Steere, 1890 . . . 32B List of the Birds and Mammals collected by the Steere Expedition to the Philippines: 17 (reprinted in Ibis, (6) 3 (1891): 312). SYNTYPE, Adult male. Reg. no. 1896.6.6.389. Catbalogan, 14 April (on label), Samar, Philippines. Collected by J. B. Steere. Steere Collection.

The female syntype from Leyte is also in the collection.

nigrocinereus—Thamnophilus nigrocinereus P. L. Sclater, 1855. 4 Proc. zool. Soc. Lond., 1855: 19, pl. 81. SYNTYPE, Adult male.

84 nigrogularis
Reg. no. 1889.9.20.53. Rio Tocantins, Brazil, 1854. Collected by A. R. Wallace. Sclater Collection. Listed as type by Sclater, <i>Cat. Bds. Brit. Mus.</i> , 15 (1890): 194. There are two other syntypes in the collection.
igrogularis—Brachypteryx nigrogularis Eyton, 1845 32B Ann. Mag. nat. Hist., (1) 16:228. Syntype, Adult. Reg. no. 1846.3.4.26. Malacca. Collected and presented by A. Charlton.
igrogularis—Lalage nigrogularis Layard, 1875
igrogularis—Ruticilla nigrogularis Moore, 1854 31 in Horsfield and Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 1:307. SYNTYPE, Adult male (relaxed mount). Reg. no. 1879.12.28.703. Nepal. Collected by B. H. Hodgson and presented by the Honourable East India Company Museum. Listed as type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881):352. There is also a female syntype in the collection.
igrogularis—Siphia nigrogularis Everett, 1891
igrogularis—Spermophila nigrogularis Gould, 1839 59 Zoology of the voyage of H.M.S. Beagle, 3, Birds: 88. SYNTYPE, Adult male. Reg. no. 1855.12.19.200. Monte Video, November. Collected by C. Darwin. Purchased of the Zoological Society of London. The female syntype was listed by Sharpe, Cat. Bds. Brit. Mus., 12: 127 but cannot now be traced.
12:127, but cannot now be traced. sigrolineata—Grallaria nigrolineata P. L. Sclater, 1890 4 Cat. Bds. Brit. Mus., 15:321. SYNTYPE, Adult male. Reg. no. 1889.9.20.634. Merida, Venezuela, 6 Nov. 1886. Collected by L. Brieño. Obtained by Sclater from Dr. E. Rey of Leipzig. Sclater Collection. There is another syntype in the collection. Sclater used Berlepsch's MS. name.
nigrolutea—Iora nigrolutea G. F. L. Marshall, 1876

lected by G. F. L. Marshall. Hume Collection.

There are several other syntypes in the collection. nigronotata—Urobrachya nigronotata Sharpe, 1897 Bull. Br. Orn. Cl., 7:7. HOLOTYPE, Adult male. Reg. no. 1897. 10.31.2. Witu, Kenya. Collected and presented by F. J. Jackson.

9.1.426. Saharunpore, North-west Province, 9 Aug. 1873. Col-

nigrorufa—Saxicola nigrorufa Jerdon, 1839
nigrorufo—Sericornis nigro rufo Salvadori, 1894
nigrorum—Alsoenax cinercus nigrorum. See appendix, Corrections and Additions, vol. 2
nigrorum—Cittocincla nigrorum Ogilvie-Grant, 1896 31 Ibis, (7) 2:547. Syntype, Adult male. Reg. no. 1897.5.13.432. Canloan Volcano, Negros, Philippines, 1 May 1896. Collected by J. Whitehead and presented by the subscribers to his expedition. There is also a female syntype in the collection.
nigrorum—Turdus nigrorum Ogilvie-Grant, 1896
nigrorum—Zosterops nigrorum Tweeddale, 1878
nigrostellatus—Pomatorhimus nigrostellatus Swinhoe, 1870 32B Ibis, (2) 6:250. Syntype, Adult. Reg. no. 1898.10.3.599. Central Hainan, China, Feb. 1868. Collected by R. Swinhoe. Seebohm Bequest.
nilghiriensis—Anthus nilghiriensis Sharpe, 1885
Cat. Bds. Brit. Mus., 10:550. HOLOTYPE, Adult (claws damaged). Reg. no. 1846.4.30.13. Nilghiri Hills, India. Collected and presented by T. C. Jerdon.
nilgiriensis—Saxicola caprata nilgirensis Whistler, 1940 31 Bull. Br. Orn. Cl., 60:90. HOLOTYPE, Adult male. Reg. no.
1886.4.1.1510. Ootacamund, Nilgiri Hills, India, 11 Jan. 1881. Collected by W. Davison. Hume Collection.
nilgiriensis—Zosterops palpebrosa nilgiriensis Ticchurst, 1927 47
Bull. Brit. Orn. Cl., 47:89. HOLOTYPE, Adult male. Reg. no. 1886.12.1.1690. Coonoor, Nilgiri Hills, South India, 28 Jan. 1881. Collected by W. Davison. Hume Collection.

Joo Miloteu
nilotica—Calamacichla ansorgei nilotica Neumann, 1905 35 Novit. Zool., 15: 246. SYNTYPE, Adult male. Reg. no. 1887.9. 28.220. Wadelai, on the Bahr-el-Jebel, Equador, Africa, 8 Aug. 1885. Collected and presented by Emin Pasha. The three other syntypes are also in the collection.
niloticus—Passer domesticus niloticus Nicoll & Bonhote, 1909 . 61 Bull. Br. Orn. Cl., 23:101. Syntype, Adult male. Reg. no. 1909.6.10.3. ("Birket-el-Karun" on label), Fayum, Egypt, 13 March 1913. Collected and presented by M. J. Nicoll and J. L. Bonhote. The female syntype is also in the collection.
nipalensis—Brachypteryx nipalensis Moore, 1854 31 in Horsfield and Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 1:397. Syntype, Adult male. Reg. no. 1880.1.1.2381. Nepal. Collected by B. H. Hodgson and presented by the Honourable East India Company Museum. There is also a female syntype in the collection.
nipalensis—Carduelis nipalensis Hodgson, 1836
nipalensis—Certhia nipalensis Blyth, 1845
nipalensis—Cinclosoma? Sibia? nipalensis Hodgson, 1836 32B Asiat. Res., 19:145. Syntype, Adult. Reg. no. 1843.1.13.583. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
nipalensis—Cinnyris nipalensis Hodgson, 1837
ninglansis Delicher ninglensis Moore 1854

There are three other syntypes in the collection. Moore used Hodgson's MS. name.

nipalensis—Drymoica nipalensis Moore, 1854 39	5
in Horsfield and Moore, A Catalogue of birds in the museum of the Honourable East India Company, 1:329. SYNTYPE, Adult (relaxed mount), Reg. no. 1880.1.1.3038. Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company	
Museum. Moore used a Hodgson MS. name. A second syntype was available.	
Asiat. Res., 19: 157. ?SYNTYPE, Adult male. Reg. no. 1880.1.1. 3907. Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company Museum. This specimen, which has the original data label, and is marked "Icon. Ined., pl. no. 341", appears to be an original syntype.	9
Inipalensis—Graucalus nipalensis Hodgson, 1837)
nipalensis—Hirundo nipalensis Hodgson, 1836	3
nipalensis—Paludicola nipalensis Hodgson, 1837	2
nipalensis—Parus nipalensis Hodgson, 1838	2
nipalensis—Pyrrhula nipalensis Hodgson, 1836	•
in Horsfield and Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 1:302. SYNTYPE, Adult male. Reg. no. 1880.1.1.1460. Nepal. Collected by B. H. Hodgson and presented by the Honourable East India Company Museum. There are several other syntypes in the collection.	l
nipalensis—Sitta nipalensis Hodgson, 1836	3

Sclater Collection.

nitidissimus—Tachyphonus nitidissimus Salvin and Godman, 1870 51 Proc. zool. Soc. Lond., 1870: 188. SYNTYPE, Adult male. Reg. no. 1885.6.7.648. Bugaba, Chiriqui, Veragua, 1869. Collected by E. Arcé. Salvin-Godman Collection. There are several other syntypes in the collection.
nitidum—Dicaeum nitidum Tristram, 1889
nitidus—Carpodectes nitidus Salvin, 1865
nitidus—Orthotomus nitidus Hume, 1874
niveiceps—Hypsipetes niveiceps Swinhoe, 1864
niveiceps—Planesticus niveiceps. See appendix, Corrections and Additions, vol. 2
Proc. zool. Soc. Lond., 1860: 281. HOLOTYPE, Adult male. Reg. no. 1888.1.13.1170. Babahoyo, Ecuador, July 1859. Collected by L. Fraser, Sclater Collection.
Proc. zool. Soc. Lond., 1863: 187, pl. 24. SYNTYPE, Adult. Reg. no. 1885.3.8.260. San Geronimo, Vera Paz, Guatamala, Jan. 1862. Collected by O. Salvin. Salvin-Godman Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 647. There are two other syntypes in the collection.
niveni—Tephrocorys cinerea niveni Macdonald, 1952 17
Ann. Transvaal Mus., 22: 32. HOLOTYPE, Adult male. Reg. no. 1933.7.14.233. Gezabuzo, near Pietermaritzburg, Natal, 23 June 1929. Collected by and purchased of J. Vincent.
niveogularis—Acanthiparus niveogularis Gould, 1855 40
Birds of Asia, 2, pt. 7: pl. 67. SYNTYPE, Adult. Reg. no. 1858. 12.2.37. North India. Collected for J. Gould. Gould Collection. Listed as the type by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 59. There is a second syntype in the collection. This species was also described by Moore, Proc. zool. Soc. Lond., 1854: 140, under the

name of <i>Orites niveogularis</i> but according to Baker, <i>Fauna of British India</i> , <i>2nd Edn.</i> 7 (1930): 17, Gould's work was published on 1 April 1855 but Moore's description was not published until 11 April.
nivriventer—Poliositta azurea nivriventer Robinson and Kloss, 1919 43 Bull. Brit. Orn. Cl., 90:17. SYNTYPE, Adult male. Reg. no. 1919.11.5.19. Tjibodas, 4-6,000', Gedeh, West Java, (11 Feb. 1916, on label). Collected by H. C. Robinson. Presented by the Federated Malay States Museum, with a female specimen, as a type. The female syntype is also in the collection.
njombe—Alethe macclouniei njombe Benson, 1936 31 Bull. Br. Orn. Cl., 56: 100. Holotype, Adult male. Reg. no. 1932.5.10.207. Njombe, 6,600′, 9.1°S., 34.75°E., 19 Dec. 1931. Collected and presented by Admiral H. Lynes, from the Lynes-Lowe Expedition.
njombe—Cisticola aberrans njombe Lynes, 1933
nobilior—Sturnus nobilior Hume, 1879
nobilis—Chamaeza nobilis Gould, 1855
nobilis—Corvus nobilis Gould, 1837
nobilis—Myiodynastes nobilis P. L. Sclater, 1859
norrisae—Sylvia norrisae Nicoll, 1917
notabilis—Glyciphilia notabilis Sharpe, 1899

. 37A

Sept. 1899, on label). Collected and presented by Capt. A. M. Farguhar from the voyage of H.M.S. Wallaroo. This specimen was labelled type by the author. There is another male syntype in the collection. notabilis—Tanagra notabilis P. L. Sclater, 1855 . Proc. zool. Soc. Lond., 1855: 84, pl. 91. HOLOTYPE, Adult. Reg. no. 1855.6.12.445. Vicinity of Quito, Ecuador, 1854. Collected by W. Jameson. Sclater Collection, (received by him from W. Jardine, vide Sclater, Catalogue of a collection of American Birds. (1862):74). See also Hellmayr and Conover, Catalogue of birds of the Americas, 9 (1936): 204, footnote. no. 1881.5.1.5558. Cape York District of Queensland, Australia. Gould Collection There are two other syntypes in the collection. notatus—Cyornis tricolor natatus Whistler, 1930 Bull. Brit. Orn. Cl., 50:70. HOLOTYPE, Adult male. Reg. no. 1925.12.23.836. Gund, Kashmir, 26 May 1896. Collected by J. Davidson Davidson Bequest. notatus—Picolaptes notatus Eyton, 1852 in Jardine, Contributions to Ornithology, 1852:26. HOLOTYPE, Adult. Reg. no. 1881.2.18.196. No locality. Eyton Collection: purchased of E. Gerrard. notosticta—Peucaea notosticta P. L. Sclater, 1868. Proc. zool. Soc. Lond., 1868: 322. HOLOTYPE, Adult. Reg. no. 1885.12.14.993. South Mexico. Collected by A. Boucard. Salvin-Godman Collection. nova—Rhipidura rufidorsa nova Mathews, 1928 Bull. Brit. Orn. Cl., 48:92. HOLOTYPE, Adult male. Reg. no. 1911.12.20.1319. Pariman, Mimika River, Dutch New Guinea, 7 Nov. 1910. Collected by C. H. B. Grant. Presented by the subscribers to the British Ornithologists' Union Expedition to Dutch New Guinea. novaeguineensis-Piezorhynchus alecto novaeguineensis Mathews, Bull. Brit. Orn. Cl., 48:93. HOLOTYPE, Adult female. Reg. no. 1911.12.20.1448. (Wakatimi, on label), Mimika River, Dutch New Guinea, 12 March 1911. Collected by C. H. B. Grant. Presented by the subscribers to the British Ornithologists' Union Expedition to Dutch New Guinea.

novaeguineensis-Setosura threnothorax novaeguineensis Mathews,

- Bull. Brit. Orn. Cl., 48:92. HOLOTYPE, Adult male. Reg. no. 1911.12.20.1292. 40 miles up Mimika River, Dutch New Guinea, 12 Feb. 1910. Collected by G. C. Shortridge. Presented by the subscribers to the British Ornithologists' Union Expedition to Dutch New Guinea.
- novaeseelandiae—Alauda novaeseelandiae Gmelin, 1789. 19

 Systema Naturae, 1, pt. 2:799. Holotype, Unsigned finished painting. Queen Charlotte Sound, South Island, New Zealand 7 Nov. 1774. No. 143 in Forster, Unpublished water-colour... drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 305.
- novaezeelandiae—Parus novaezeelandiae Gmelin, 1789 3

 Systema Naturae, 1, pt. 2:1013. Holotype, unsigned finished painting. Dusky Sound, South Island New Zealand, 20 April 1789. No. 166 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959):310.
- nuchalis—Garrulax nuchalis Godwin-Austen, 1876 32B Ann. Mag. nat. Hist., (4) 18:411. HOLOTYPE, Adult female. Reg. no. 1895.7.14.1920. Lhota-Naga Hills, south Assam, 21 Jan. 1876. Collected by M. T. Ogle for A. W. Chennell. Godwin-Austen Collection.

- nuchalis—Pomatorhinus nuchalis Tweeddale, 1877. 32B

 Ann. Mag. nat. Hist., (4) 20:535. SYNTYPE, Adult male. Reg.
 no. 1888.4.20.69. Karen-nee, 3,000′, north-east Tenasserim, 5

 March 1874. Collected by R. G. W. Ramsay. Tweeddale

 Collection.
 See Ramsay, Ibis, (4) 2 (1878):134.
- nuchalis—Sycobius nuchalis Elliot, 1859 61

 Ibis, (1) 1: 393. HOLOTYPE, Adult (female vide Sharpe, Cat. Bds.

 Brit. Mus., 13 (1890): 478). Reg. no. 1859.6.28.5. Gaboon,

 French West Africa. Collected for and purchased of Verreaux.

Collected by Admiral H. Lynes. Presented by A. Chapman and Admiral H. Lynes. The described female syntype is also in the collection.
nyasa—Cisticola erythrops nyasa Lynes, 1930 35 Ibis, (12) 6, Supplement: 374. HOLOTYPE, Adult female. Reg. no. 1920.10.25.10. Chirimo, Ruo District, Nyasaland, 9 April 1918. Collected and presented by R. C. Wood.
nyasae—Amydrus nyasae Shelley, 1898
nyasae—Anomalospiza imberbis nyasae Benson, 1938 61 Bull. Br. Orn. Cl., 58:112. HOLOTYPE, Adult male. Reg. no. 1938.4.13.2. Dowa, 4,000′, central Nyasaland, 19 Sept. 1937. Collected and presented by C. W. Benson.
nyasae—Erythrocercus nyasae Ogilvie-Grant, 1912
nyasae—Hyphantornis nyasae Shelley, 1844 61 Ibis, (6) 6: 20. Syntype, Adult. Reg. no. 1893.6.1.226. Milanji Plains, Nyasaland, Aug. 1892. Collected by A. Whyte. Presented by Sir H. H. Johnston. There are two other syntypes in the collection.
nyasae—Pseudoalcippe pyrrhoptera nyasae Benson, 1939 32B Bull. Br. Orn. Cl., 59:43. HOLOTYPE, Adult male. Reg. no. 1939.1.12.2. Nyakhowa Mountain, near Livingstonia, Nyasaland, 13 Oct. 1938. Collected and presented by C. W. Benson.
nyassae—Anthreptes longmari nyassae Newmann, 1906 46 Orn. Mber., 14:7. HOLOTYPE, Adult male. Reg. no. 1897.11.4. 260. Lahengula, Zomba, Nyasaland, 27 July 1897. Collected by A. Whyte. Presented by Sir H. H. Johnston.
nyassae—Bradypterus nyassae Shelley, 1893
nyika—Cisticola aberrans nyika Lyncs, 1930

nuerensis—Calamocichla leptorhyncha nuerensis Lynes, 1914. . . 35
Bull. Brit. Orn. Cl., 33:130. Syntype, Adult male. Reg. no.
1919.12.17.453. Lake No, Nuer, Upper White Nile, 25 Feb. 1914.

nyikae—Mirafra africana nyikae Benson, 1939
nyikae—Othyphantes stuhlmanni nyikae Benson, 1938 61 Bull. Br. Orn. Cl., 58:112. HOLOTYPE, Adult male. Reg. no. 1938.4.13.1. Nyika Plateau, 8,000', North Nyasa, 12 Nov. 1937. Collected and presented by C. W. Benson.
nyikensis—Muscicapa nyikensis Shelley, 1899 37 Bull. Brit. Orn. Cl., 8:35. Syntype, Adult female. Reg. no. 1897.11.4.80. Nyika Plateau, 6,000–7,000′, Nyasaland, (June 1896, on label). Collected by A. Whyte. Presented by Sir H. H. Johnston. There is another syntype in the collection.
nyikensis—Nectarinia johnstoni nyikensis. See appendix, Corrections and Additions, vol. 2
nyong—Acrocephalus baeticatus nyong Bannerman, 1936 35 Bull. Brit. Orn. Cl., 57:9. HOLOTYPE, Adult male. Reg. no. 1920.6.26.437. Akonolinga, Nyong River, South Cameroons, 4
Jan. 1913. Collected and presented by G. L. Bates. oatesi—Garrulus oatesi Sharpe, 1896
oatesi—Hydrornis oatesi Hume, 1873
oatesi—Siva cyanuroptera oatesi Harington, 1913
oatesi—Urocichla oatesi Rippon, 1904
oatesii—Bradyornis oatesii Sharpe, 1881

purchased of A. E. Hamerton. obscura—Anthocichla phayrei obscura Delacour, 1927. 12 Bull. Brit. Orn. Cl., 47:155. HOLOTYPE, Adult female. Reg. no. 1927.6.5.5. Lake Babé, north-east Tonkin, French Indo-China, 12 Dec. 1926 (20 Dec., on label). Collected by J. Delacour, W. P. Lowe, and F. Jabouille, on the 3rd Anglo-French Expedition to French Indo-China. Bull. Br. Orn. Cl., 47:161. HOLOTYPE, Adult male. Reg. no. 1927.6.5.16. Tam Dao, central Tonkin, 2 Dec. 1926. Collected by J. Delacour, P. Jabouille & W. P. Lowe on the 3rd Anglo-French Expedition to Indo-China. obscura—Muscicapa obscura Horsfield, 1821 Trans. Linn. Soc. Lond., 13:146. SYNTYPE, Adult. Reg. no. 1860. 4.16.120. Java. Collected by T. Horsfield and presented by the Honourable East India Company Museum. There are several other syntypes in the collection. obscura—Siphia obscura Sharpe, 1881 Proc. zool. Soc. Lond., 1881: 789. HOLOTYPE, Adult. Reg. no. 1881.5.1.1346. "Borneo" (= South-east Brazil, vide Hellmayr and Conover, Catalogue of birds of the Americas, 8 (1935): 521). Gould Collection. obscura—Stachyridopsis rufifrons obscura E. C. S. Baker, 1917 . 32B Bull. Br. Orn. Cl., 38: 10. SYNTYPE, Adult male. Reg. no. 1916. 12.27.865. Klong Bang Lai, Siam, 20 Jan. 1916. Collected and presented by E. G. Herbert. This is the author's selected male. His selected female syntype is also in the collection. obscura—Suva obscura Hume, 1874 . Stray Feathers, 2:507. HOLOTYPE, Immature female. Reg. no. 1877.6.5.2. Dunna, Cashmere, August 1874. Collected by and purchased by J. Biddulph. Old Vellum Cat. 14.359a. Straits of Magellan, 1827. Collected and presented by Capt. P. P. King. There is another syntype in the collection. obscurior—Alcippe obscurior Ogilvie-Grant, 1906 . 32B Bull. Br. Orn. Cl., 16:121. SYNTYPE, Adult male. Reg. no. 1907.12.12.63. Mount Morrison, 9,000', central Formosa, Feb. 1906. Collected by and purchased of W. Goodfellow.

There is also a female syntype in the collection.

obscurior—Cinnyris obscurior Ogilvie-Grant, 1894
5.13.46. (La Trinadad, on label) Benguet District, north-west Luzon, Philippines, 25 Feb. 1894. Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. A female syntype is also in the collection.
obscurior—Yuhina occipitalis obscurior Rothschild, 1921
obscurum—Dicaeum obscurum Ogilvie-Grant, 1894
obscurus—Nectarinia obscurus Jardine, 1843
obscurus—Pomatorhinus obscurus Hume, 1873
obscurus—Pyrocephalus obscurus Gould, 1839
obsoleta—Grallaria nuchalis obsoleta Chubb, 1916. Bull. Br. Orn. Cl., 36: 47. HOLOTYPE, Adult male. Reg. no. 1916. 8.24.46. West side of Pichincha Mt., 12,000'. Ecuador, Nov. 1914. Collected by W. Goodfellow and presented by E. J. Brook. Chubb had an example of either sex but only described the male.
occidens—Cisticola eximia occidens Lynes, 1930 35 Ibis, (12) 6, Supplement: 178, pl. 7. HOLOTYPE, Adult male. Reg. no. 1912.12.26.63. Rirn, northern Nigeria, 4 Jan. 1912. Collected and presented by Capt. J. J. Brocklebank.

- occidentalis—Accentor modularis occidentalis Hartert, 1910 . . . 30
 Brit. Bds., 3: 313. HOLOTYPE, Adult male. Reg. no. 1936.10.15.
 10. Tring, Hertfordshire, 10 April 1893. Presented by the American Museum of Natural History, from the Rothschild Collection.
- occidentalis—Catharus occidentalis P. L. Sclater, 1859 31

 Proc. zool. Soc. Lond., 1859: 323. SYNTYPE, Adult (female on label). Reg. no. 1886.8.2.6. Totontepec, Mexico. Collected by Boucard. Sclater Collection.

 Listed as type by Sclater, Catalogue of a collection of American Birds, 1826: 1.
- occidentalis—Certhionyx occidentalis Ogilvie-Grant, 1909 . . . 48

 Ibis, (9) 3:658. HOLOTYPE, Adult male. Reg. no. 1905.12.26.131.

 Gascoyne River, Carnarvon, Western Australia, 19 August 1906.

 Collected by G. C. Shortridge. Presented by W. E. Balston.
- occidentalis—Chlorocichla occidentalis Sharpe, 1881 . . . 21
 Cat. Bds. Brit. Mus., 6:113, pl. 8. Syntype, Adult female. Reg.
 no. 1876.5.23.629. Ovaquenyama, Damaraland, 28 May 1867.
 Collected by C. J. Andersson. Sharpe collection.
 There are two other syntypes in the collection.
- occidentalis—Conopophaga aurita occidentalis Chubb, 1917. . . . 5

 Bull. Br. Orn. Cl., 38: 34. SYNTYPE, Adult male. Reg. no. 1859.

 7.6.62. Rio Napo, Ecuador. Probably collected by Villavicencio; purchased of J. Gould, who obtained it from Verreaux.

 The female syntype is in the collection.
- occidentalis—Dendrocitta formosae occidentalis Ticehurst, 1925 . 71

 Bull. Br. Orn. Cl., 46: 22. HOLOTYPE, Adult male. Reg. no. 1886.3.1.702. Simla, north-west Himalayas, 12 Dec. 1880. Collected by A. O. Hume. Hume Collection.
- occidentalis—Muscivora occidentalis P. L. Sclater, 1860. . . 9

 Proc. zool. Soc. Lond., 1860: 282. Syntype, Adult male. Reg. no. 1888.1.13.807. Babahoyo, Ecuador, Aug. 1859. Collected by L. Fraser. Sclater Collection.

 The female syntype is also in the collection.
- occidentalis—Oryzoborus occidentalis P. L. Sclater, 1860 . . . 59
 Proc. zool. Soc. Lond., 1860 : 276. HOLOTYPE, Adult male. Reg.
 no. 1885.2.10.86. Babahoyo, Ecuador, Aug. 1859. Collected by
 L. Fraser. Sclater Collection.
- occidentalis—Passer occidentalis Shelley, 1883 61

 Ibis, (5) 1:548, New name for Pyrgita simplex Swainson, 1837.

 which is preoccupied. The specimen referred to by Sharpe, Cat.

 Bds. Brit. Mus., 12 (1888):337 as the type is that examined by Shelley when proposing the new name.

occidentalis—Procnias occidentalis P. L. Sclater, 1855 Proc. zool. Soc. Lond., 1854 (publ. 1855): 249. HOLOTYPE, Adult male. Reg. no. 1885.6.12.5. Bogota, Colombia, 1853. Collected for Argent. Sclater Collection.

Since the description refers only to the male and makes no mention of the very different female plumage it would appear that the female specimen listed by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 52, is not

a syntype.

- occidentalis—Pterythius xanthochloris occidentalis Harington, 1913, 32B Bull. Br. Orn. Cl., 33:82 HOLOTYPE, Adult male. Reg. no. 1888. 5.30.237. Dehra, north-west Himalayas. Collected by G. F. L. Marshall. Tweeddale Collection.
- occidentalis—Salpornis salvadorii occidentalis Clarke, 1914 Bull, Brit. Orn. Cl., 33:94. HOLOTYPE, Adult. Reg. no. 1914.1.9. 1. Ankpa, northern Nigeria. Collected and presented by S. R. Clarke.
- occidentalis—Schetba rufa occidentalis Delacour, 1931. Oiseau, (n.s.) 1:485. HOLYTYPE, Adult male. Reg. no. 1931.8.18. 3249. Tabiky, west of Ankosoabo, south-west Madagascar, 11 Nov. 1929. Collected by J. Delacour on the Archbold-Vernay Expedition.
- occidentis—Zosterops palpebrosa occidentis Ticehurst, 1927 . Bull. Brit. Orn. Cl., 47:88. HOLOTYPE, Adult. Reg. no. 1886. 12.1.1613. Simla, north-west Himalayas, 19 Oct. 1880. Collected by W. Davison. Hume Collection.
- occipitale—Pyrgisoma occipitale Salvin, 1878 . Ibis, (4) 2:446. SYNTYPE, Adult (= immature, vide Sclater and Salvin, Biol. Centr.-Amer., 1 (1886): 404). Reg. no. 1885.12.14. 1364. Savana Grande, Guatemala, 1862. Salvin Godman Collection.

This specimen was figured in Sclater and Salvin, Exotic Ornithology (1868): 64, and was referred to as the type by the author. There are other syntypes in the collection.

- occipitales—Yuhina occipitales Hodgson, 1836 . 32B Asiat. Res., 19:166. SYNTYPE, Adult (relaxed mount). Reg. no. 1843.1.13.371. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
- occipitalis—Chlamydodera occipitalis Gould, 1875 69 Ann. Mag. nat. Hist., (4) 16: 429. HOLOTYPE, Adult male. Reg. no. 1881.5.1.1779. New Guinea (Corrected to Port Albany, northern Australia vide Gould, Birds of New Guinea, 1 (1879): text to pl. 45. Jardine given as collector). Purchased of E. Jansen for Gould Collection.

- ocellata—Spermophila ocellata P. L. Sclater and Salvin, 1866. 59

 Proc. zool. Soc. Lond., 1866: 181. SYNTYPE, Adult male. Reg.
 no. 1885.2.10.164. Nauta, Ucayuli River, East Peru, (17 April 1865, on label). Collected by E. Bartlett. Sclater Collection.

 Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 131.

 This is the author's selected male syntype, his selected female syntype and other male syntypes are also in the collection.
- ocellatus—Harporhynchus ocellatus P. L. Sclater, 1862. . . . 29

 Proc. zool. Soc. Lond., 1862: 18, pl. 3. HOLOTYPE, Adult male.

 Reg. no. 1886.8.2.209. Oaxaca, southern Mexico, Nov. 1860.

 Collected by A. Boucard for A. Sallé. Sclater Collection.

A female syntype is also in the collection. No holotype was designated and the specimen listed as the "type" by Hartert, *Novit.* Zool., 26 (1919): 170, was one of a series of syntypes which included the above specimens.

- ochracea—Prionops concinnata ochracea Lynes, 1920 23
 Bull. Br. Orn. Cl., 41:18. HOLOTYPE, Adult male. Reg. no. 1920.
 12.22.692. 10 miles east of Nahud, western Kordofan, Sudan, 22
 May 1920. Collected and presented by Admiral H. Lynes.
- ochraceiceps—Hylophilus ochraceiceps P. L. Sclater, 1859 . . . 56

 Proc. zool. Soc. Lond., 1859: 375. HOLOTYPE, Adult female. Reg.
 no. 1886.9.15.313. Playa Vicente, Oaxaca, southern Mexico,
 April 1859. Collected by A. Boucard. Sclater Collection.
- ochraceiceps—Pomatorhinus ochraceiceps Walden, 1873. . . . 32B

 Ann. Mag. nat. Hist., (4) 12: 487. SYNTYPE, Adult. Reg. no. 1888.
 4.20.717. Karen Hills, Burma, (Jan. 1872 on label). Collected by R. G. W. Ramsay. Tweeddale Collection.

 There are two other syntypes in the collection.
- ochraceus—Contopus ochraceus P. L. Sclater and Salvin, 1869. . 9
 Proc. zool. Soc. Lond., 1869: 419. HOLOTYPE, Adult. Reg. no. 1899.4.20.2626. Costa Rica, 1869. Collected by J. Carmiol. Salvin-Godman Collection.
- ochraceus—Drymocataphus tickelli ochraceus Kinnear, 1934. 32B Bull. Br. Orn. Cl., 55: 53. New name for D. t. olivaceus Kinnear,

- 1924 (q.v.) on the grounds of its preoccupation by *Mixornis olivaceus* Tickell, 1859.
- ochrocephalus—Muscicapa ochrocephalus Gmelin, 1789. . . . 36

 Systema Naturae, 1, pt. 2., : 944. Holotype, unsigned unfinished painting. Dusky Sound? Queen Charlotte Sound, South Island, New Zealand, 4 April 1774. No. 157 in Forster, Unpublished water-colour drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1, (1959): 308.
- ochrofulvescens—Planesticus fumigatus ochrofulvescens Chubb, 1918 31 Bull. Br. Orn. Cl., 38: 54. HOLOTYPE, Adult. Reg. no. 1905.6.28. 124. Trinidad, Jan. 1863. Cavendish Taylor Bequest.
- ochrogularis—Seicercus ruficapilla ochrogularis Moreau, 1941. . 39

 Bull. Brit. Orn. Cl., 61:24. Holotype, Adult male. Reg. no. 1940.11.1.1. Forest of Kungwe Mountain, 6,500′, West Tanganyika, 22 August 1940. Collected by C. Abdullah for R. E. Moreau. Presented by R. E. Moreau.

Listed as type by P. L. Sclater, Cat. Bds. Brit. Mus., 14 (1888): 466.

This is the described male. The described female syntype is also in the collection.

ocularis—Euscarthmus occularis Salvin, 1876)
ocularis—Motacilla ocularis Swinhoe, 1860)
ocularis—Phrygilus ocularis P. L. Sclater, 1859)
ocularis—Ploceus ocularis Smith, 1828 61 South African Commercial Advertiser no. 144, (vide Macdonald and Grant, Ann. Transvaal Mus., 23 (1953): 98, 202). SYNTYPE Adult male (relaxed mount). Reg. no. 1845.7.6.26. Northward, of the tropic of Cancer. (South-east Africa, vide Smith, Illustration of the zoology of South Africa, Aves, (1839) pl. 30). Collected by and purchased of A. Smith. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 428.	
ocularius—Drymoica ocularius Smith, 1843	;
oenops—Garrulus oenops Whitaker, 1898	
ogilvie-granti—Anthreptes ogilvie-granti Bannerman, 1921	
oglei—Actinura oglei Godwin-Austen, 1877	
oinops—Ocyris oinops Hodgson, 1845	

- Reg. no. 1880.1.1.4680. Nepal. Collected by B. H. Hodgson. Presented by the Honourable East India Company Museum. The bill is incomplete. There are other syntypes in the collection. This was selected by N. B. Kinnear and is not one of the specimens listed as types by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 489. okuensis—Alseonax minimus okuensis Bates, 1926. Bull. Brit. Orn. Cl., 46:90. HOLOTYPE. Adult male. Reg. no. 1926.8.8.5. Oku, west of Kumbo, Cameroons, 7 Feb. 1925. Collected by and purchased of G. L. Bates. okuensis—Coracina caesia okuensis Bannerman, 1938 . 20 Bull. Br. Orn. Cl., 58:76. HOLOTYPE, Adult male. Reg. no. 1926.8.8.611. Oku, 7,000', west of Kumbo, Cameroons, 14 Feb. 1925. Collected by and purchased of G. L. Bates. okuensis—Phormophlectes insignis okuensis Bannerman, 1943. Bull. Br. Orn. Cl., 63:64. HOLOTYPE, Adult female. Reg. no. 1926.8.8.561. Oku, 7,000', west of Kumbo, Cameroons, 11 Feb. 1925. Collected by and purchased of G. L. Bates. oldeani—Turdus olvaceus oldeani W. L. Sclater & Moreau, 1935 Bull. Br. Orn. Cl., 56:13. HOLOTYPE, Adult male. Reg. no. 1935.10.11.1. Oldeani Forest, 6,500', Mbulu District, Tanganyika, 6 Sept. 1934. Collected for and purchased of R. E. Moreau. oleaginea—Eucometes oleaginea Salvin, 1886 51 Ibis, (5) 4:500. HOLOTYPE, Adult female. Reg. no. 1885.6.8.444. Twek-quay Mountain, Carimang River, British Guiana (17 Nov. 1885, on label). Collected by H. Whitely. Salvin-Godman Collection. oleagineus—Anabazenops oleagineus P. L. Sclater, 1883. . Proc. zool. Soc. Lond., 1883:654. SYNTYPE, Adult male. Reg.

no. 1889.5.20.379. Sierra de Totoral, Catamarca, Argentina, 29

July 1880. Collected by E. W. White. Sclater Collection.

- oleagineus—Ostinops oleagineus P. L. Sclater, 1883 57

 Ibis, (5) 1:154. HOLOTYPE, Adult male. Reg. no. 1885.7.12.33.

 Venezuela ? 1857. Collected for Verreaux. Sclater Collection.
- olimotiensis—Pinarochroa sordida olimotiensis Elliott, 1945 . . . 31 Bull. Br. Orn. Cl., 66:19. HOLOTYPE, Adult male. Reg. no. 1945.19.1. Olimoti Mountain, 9,000', north of Ngorongoro, north-east Tanganyika, 15 July 1944. Collected and presented by H. F. I. Elliott.

Proc. zool. Soc. Lond., 1837: 7. SYNTYPE, Adult (relaxed mount). Reg. no. 1855.12.19.127. Galapagos Archipelago. Collected by C. Darwin. Purchased from the Zoological Society of London. There are several other syntypes in the collection.
olivacea—Cyornis olivacea Hume, 1877
olivacea—Hypsepetes olivacea Jardine & Selby, 1837 21 Illustrations of Ornithology, (n.s.): pl. 2. HOLOTYPE, Adult. Reg. no. 1886.6.24.254. ?Mauritius. Collected by Dyce. Jardine Collection; purchased of E. Gerrard. See also Jardine & Selby. Illustrations of Ornithology, 1st ser., 3 (1835): 148.
olivacea—Pachycephala olivacea Vigors and Horsfield, 1827 39 Trans. Linn. Soc. Lond., 15: 241. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1863.7.7.41. No locality (= Tasmania, vide Mathews, Birds of Australia, 8 (1920): 261). Collected by G. Caley. Presented by the Linnean Society.
olivacea—Phragamaticola olivacea Jerdon, 1844
olivacea—Prinia olivacea Strickland, 1844
olivacea—Sericornis olivacea Salvadori, 1896
olivaceiceps—Chlorospingus olivaceiceps Underwood, 1898 51 Bull. Brit. Orn. Cl., 7:59. Holotype, Adult female. Reg. no. 1898.12.14.2417. Carrillo, Costa Rica, 24 Nov. 1897. Collected by C. F. Underwood. Salvin-Godman Collection.
olivaceiceps—Criniger olivaceiceps Shelley, 1896 21 Ibis, (7) 2: 179. HOLOTYPE, Adult female. Reg. no. 1896.3.14.35. Mount Chiradzulu, Shiré Highlands, Nyasaland, July 1895. Collected by A. Whyte and presented by Sir H. H. Johnston.

olivaceum—Dicaeum olivaceum Walden, 1875 Ann. Mag. nat. Hist., (4) 15: 401. Syntype, Adult. Reg. no. 1888. 7.12.115. Karen Hills, Burma, (Nov. 1874, on label). Collected by R. G. Wardlaw Ramsay. Tweeddale Collection. Four specimens were referred to but this is labelled as type by the author. There is another syntype in the collection.
olivaceus—Agrilorhinus olivaceus Fraser, 1840
olivaceus—Cephalopyrus flammiceps olivaceus Rothschild, 1923 . 41 Novit. Zool., 30: 263. Holotype, Adult male. Reg. no. 1923.11. 11.424. Tengyueh, 8,000', north-west Yunnan, March 1922. Collected for G. Forrest. Presented by Stephenson Clarke.
olivaceus—Coccothraustes olivaceus Fraser, 1842 59 Proc. zool. Soc. Lond., 1842 : 144. Holotype, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 12 (1888) : 435). Reg. no. 1847.3.12.2. Clarence, Fernando Po. Collected by and purchased of L. Fraser.
olivaceus—Cyrtostomus frenatus olivaceus Mathews, 1926
olivaceus—Drymocataphus tickelli olivaceus Kinnear, 1924 32B Bull. Br. Orn. Cl., 45:11. HOLOTYPE, Adult male. Reg. no. 1924. 12.21.173. Bao Ha, Tonkin, 31 Oct. 1923. Collected by H. Stevens and presented by the Percy Sladen Trust and Godman Exploration Fund. Renamed Drymocataphus tickelli ochraceus Kinnear, 1934 on grounds of preoccupation by Mixornis olivaceus Tickell, 1859.
olivaceus—Microtarsus olivaceus Moore, 1854
olivaceus—Prionochilus olivaceus Tweeddale, 1877
olivaceus—Synallaxis olivaceus Eyton, 1851

oliviae—Sylviella oliviae Boyd Alexander, 1908 35 Bull. Brit. Orn. Cl., 23:16. HOLOTYPE, Adult male. Reg. no. 1911.12.23.2573. Bamingui River, French Sudan, 22 August 1905. Collected by W. Boyd Alexander. Boyd Alexander Bequest.
olivascens—Cinnicerthia olivascens Sharpe, 1881
oliviae—Sylvia oliviae Boyd Alexander, 1908 31 Bull. Br. Orn. Cl., 23:15. Syntype, Adult male. Reg. no. 1911. 12.23.2249. Near Yo, Lake Chad, French Equatorial Africa, 21 Nov. 1904. Collected by Boyd Alexander on the Alexander- Gosling Expedition. Boyd Alexander Bequest. This is the author's selected syntype. The female syntype is also in the collection.
olivina—Elainea olivina Salvin and Godman, 1884 9 Ibis, (5) 2:446. SYNTYPE, Adult male. Reg. no. 1888.1.1.787. Roraima, 3,500', British Guiana, 19 Nov. 1883. Collected by H. Whitely. Salvin-Godman Collection. There are four other syntypes in the collection.
olivina—Tanagra olivina P. L. Sclater, 1865
ombriosus—Parus ombriosus Meade-Waldo, 1890 42 Ann. Mag. nat. Hist., (6) 5: 103. Syntype, Adult male. Reg. no. 1890.7.13.4. El Pinal, El Heirro Island, Canary Islands, 22 Nov. 1899. Collected and presented by E. G. Meade-Waldo. There are several other syntypes in the collection, and one in the University Museum of Zoology, Cambridge.
omdurmanensis—Alaemon alaudipes omdurmanensis Hachisuka, 1924 17 Bull. Br. Orn. Cl., 44:86. HOLOTYPE, Adult male. Reg. no. 1916.9.20.1297. Omdurman, Sudan, 28 Nov. 1902. Collected and presented by A. L. Butler.
omoensis—Cossypha omoensis Sharpe, 1900
ongumaensis—Calendrella cinerea ongumaensis C. H. B. Grant & Mackworth-Praed, 1955

South-West Africa, 7 Oct. 1936. Collected by and purchased W. Hoesch.	
ophthalmica—Spermophila ophthalmica P. L. Sclater, 1860. Proc. zool. Soc. Lond., 1860: 276. SYNTYPE, Adult male. R no. 1885.2.10.148. Babahoyo, Ecuador, July or Aug. 1859. Clected by L. Fraser. Sclater Collection; registered as the type The female syntype is also in the collection.	teg. Col-
opistherythra—Rhipidura opistherythra P. L. Sclater, 1883. Proc. zool. Soc. Lond., 1883: 197. Syntype, Adult female. R no. 1883.5.30.84. Larat Island, Tenimber Group, 3 August 18 Collected by H. O. Forbes. Presented by the Timor Laut Comittee of the British Association. This specimen is labelled as the type. A second syntype is also the collection.	Reg. 882. om-
oreas—Centrites oreas P. L. Sclater and Salvin, 1869 Proc. zool. Soc. Lond., 1869: 154. Syntype, Adult male. Reg. 1888.1.13.213. Tinta, 11,000', Peru, 6 May 1868. Collected H. Whitely. Sclater Collection. There are several other syntypes in the collection.	no.
oreas—Pitta oreas Swinhoe, 1864	
oreobates—Cinclodes oreobates Scott, 1900	bia,
oreocincloides—Turdus oreocincloides Hodgson, 1844. in J. E. Gray, Zoological Miscellany, 1844: 83 (nomen nudu. A specimen in the collection was listed as type by Seebohm, 6 Bds. Brit. Mus., 5 (1881): 160.	. 31 un). Cat.
oreophila—Cettia oreophila Sharpe, 1888	cted
oreophila—Saxicola oreophila. See appendix, Corrections Additions, vol. 2	and . 31
orientalis—Accentor orientalis Sharpe, 1883	. 30 881. lon.

- orientalis 407 orientalis—Alauda leiopus vel orientalis Hodgson, 1843. in J. E. Grav. Zoological Miscellany, 1843: 84 (nomen nudum). Two specimens in the collection were listed as types by Sharpe. Cat. Bds. Brit. Mus., 13 (1890): 575. orientalis—Arses orientalis Salvadori, 1890. Ann. Mus. civ. Stor. nat. Giacomo Doria, (2) 9:566. SYNTYPE, Adult male. Reg. no. 1897.8.29.1. Rigo District, South-east New Guinea, 24 July 1889. Collected by the Marquis L. Loria. Purchased of Genoa Museum. This is specimen b. of the five syntypes listed. orientalis—Chlamydodera orientalis Gould, 1879. 69 Ann. Mag. nat. Hist., (5) 4:74. SYNTYPE, Adult male. Reg. no. 1881.5.1.1774. Port Denison, Queensland, (20 July, 1863 on label). Gould Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 392. The three specimens in the Australian Museum, listed as paratypes by Hindwood, Records of the Australian Museum, 21 (1946): 393. were probably not seen by Gould. orientalis—Cisticola (mentalis) orientalis Sharpe, 1883.
- orientalis—Cisticola (mentalis) orientalis Sharpe, 1883 35
 Cat. Bds. Brit. Mus., 7: 245. Syntype, Adult. Reg. no. 1895.5.1.
 1161. (Ushanbala Mountain, on label), Pangani River, East Tanganyika, 1879. Collected by J. Kirk. Shelley Collection.
 There is a second syntype in the collection.
- orientalis—Culicicapa ceylonensis orientalis E. C. S. Baker, 1923 . 37A Bull. Brit. Orn. Cl., 44:11. HOLOTYPE, Adult female. Reg. no. 1914.5.6.261. Szechuan, West China. Collected by F. W. Styan. Styan Collection.

- orientalis—Merula torquata orientalis Seebohm, 1888 31

 Ibis, (5) 6:311. Syntype, Adult male. Reg. no. 1887.11.2.2.

 Kislovodsk, Caucasus, 12 March 1886. Probably collected by G. Radde (vide Seebohm, Monograph of the Turdidae, 2 (1902):33), and received in exchange from the Petersburg Museum.

 There are two other syntypes in the collection. (See also appen-

There are two other syntypes in the collection. (See also appendix, Corrections and Additions, vol. 2.)

orientalis—Munia malacca orientalis E. C. S. Baker, 1925 . . . 60 Bull. Brit. Orn. Cl., 45:58. HOLOTYPE, Adult male. Reg. no.

- 1887.7.1.1937. Madras, South India, March 1877. Hume collection.
- orientalis—Pomatorhinus ferruginosus orientalis Delacour, 1927 . 32B Bull. Br. Orn. Cl., 47:159. HOLOTYPE, Adult male. Reg. no. 1927.6.5.23. Tam Dao, central Tonkin, 2 Dec. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China.
- orientalis—Pratincola torquata orientalis W. L. Sclater, 1911 31

 Ibis, (9) 5: 409. Syntype, Adult male. Reg. no. 1905.12.29.766.

 Umfolosi Station, Zululand. Collected by C. H. B. Grant and presented by C. D. Rudd.
 - There are many other syntypes in the collection.
- orientalis—Siva sordida orientalis Robinson & Kloss, 1919 . . . 32B Ibis, (11) 1:587. Syntype, Adult male. Reg. no. 1919.12.20.322. Langbian Peaks, 6–7,500′, south Annam, 18 April 1918. Collected and presented by H. C. Robinson and C. B. Kloss.

This is the authors' selected male. Their selected female syntype is also in the collection.

Two specimens were originally received but only one was recorded subsequently, *vide* Sharpe, *Cat. Bds. Brit. Mus.*, 3 (1877): 100.

orostruthus—Phyllastrephus orostruthus Vincent, 1933. Bull. Br. Orn. Cl., 53:133. HOLOTYPE, Adult male. Reg. no. 1933.3.1.5. Namuli Mount, 4.800', 15°21'S., 37°04'E., Portuguese East Africa, 8 Aug. 1932. Collected and presented by J. Vincent. osburni—Laletes osburni P. L. Sclater, 1861. Proc. zool. Soc. Lond., 1861:72, pl. 14. SYNTYPE, Adult male. Reg. no. 1861.9.16.2. Freeman's Hall, Trelawny, Jamaica, 28 Apr. 1859. Collected by W. Osburn. Presented by P. L. Sclater. There are two other syntypes in the collection. osiris—Nectarinia osiris Finsch, 1872. Trans. zool. Soc. Lond., 7:230. SYNTYPE, Adult male. Reg. no. 1888.7.1.199. Senafé, North-east Abyssinia, 22 May 1868. Colby W. Jesse. Tweeddale Collection. The second syntype is also in the collection. ottolanderi-Pomatorhinus montanus ottolanderi Robinson, 1918 . 32B J. F. M. S. Mus., 7:235. HOLOTYPE, Adult male. Reg. no. 1927. 4.18.990. Sodong Gerok, 3,900', Idjen Massif, near Banjoewangi, eastern Java, 1 April 1916. Collected and presented by H. C. Robinson. oustaleti—Cinclodes oustaleti Scott, 1900 Bull. Br. Orn. Cl., 10:62. SYNTYPE, Adult. Reg. no. 1892.2.10. 774. Central Chile (Valparaiso, vide Hellmayr, Catalogue of Birds of the Americas, 4 (1925): 32). Collected and presented by H. Berkeley James. There are several other syntypes in the collection. ovamboensis—Erythropygia munda ovamboensis Neumann, 1920 . 31 J. Orn., 68:83. SYNTYPE, Adult male. Reg. no. 1876.5.23.460. Ombongo, Ovampoland, South-West Africa, 21 June 1859. Collected by C. J. Andersson. Sharpe Collection. The second syntype is also in the collection. ovampensis—Prinia superciliosa ovampensis Macdonald, 1941. Bull. Brit. Orn. Cl., 62:28. HOLOTYPE, Adult male. Reg. no. 1876.5.23.231. Ovaquenyama, Damaraland, 23 May 1867. Collected by C. J. Andersson. R. B. Sharpe Collection. owstoni—Nucifraga owstoni Ingram, 1910 71 Bull. Brit. Orn. Cl., 25:86. SYNTYPE, Adult female. Reg. no. 1910.6.4.2. Mount Arizan, Central Formosa, 18 Nov. 1909. Collected and presented by A. Owston. There are two other syntypes in the collection. oxyrhyncha—Poospiza oxyrhyncha P. L. Sclater, 1865 . Proc. zool. Soc. Lond., 1864 (publ. 1865): 608. HOLOTYPE, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 766). Reg. no. 1885.2.10.314. Curytiba, Paraná, Brazil, 20 Nov. 1820. Collected by J. Natterer. Sclater Collection; obtained by Sclater in exchange

with the Vienna Museum.

Sclater used Natterer's MS. name.

pacifica — Coereba pacifica P. R. Lowe, 1912 Bull. Brit. Orn. Cl., 29:85. HOLOTYPE, Adult male. Reg. no. 1884.7.31.198. Pacasmayo, Peru, 4 June 1877. Collected by J. Stolzmann. Sclater Collection. paena — Erythropygia paena Smith, 1836 Report of the Expedition for exploring central Africa, 1834: Appendix: 46. SYNTYPE, Adult male (relaxed mount). Reg. no. 1838.6.9.95. North of Kuruman, Latakoo. Collected by A. Smith and purchased of the South African Museum. There is a second syntype in the Edinburgh Museum. palawanensis — Broderipus palawanensis Tweeddale, 1878 Proc. zool. Soc. Lond., 1878: 616. HOLOTYPE, Adult female. Reg. no. 1887.11.20.514. Puerto Princesa, Palawan, Philippines, Jan. 1878. Collected by A. H. Everett. Tweeddale Collection. palawanensis — Criniger palawanensis Tweeddale, 1978	
Report of the Expedition for exploring central Africa, 1834: Appendix: 46. Syntype, Adult male (relaxed mount). Reg. no. 1838.6.9.95. North of Kuruman, Latakoo. Collected by A. Smith and purchased of the South African Museum. There is a second syntype in the Edinburgh Museum. Palawanensis — Broderipus palawanensis Tweeddale, 1878. 66. Proc. zool. Soc. Lond., 1878: 616. Holotype, Adult female. Reg. no. 1887.11.20.514. Puerto Princesa, Palawan, Philippines, Jan. 1878. Collected by A. H. Everett. Tweeddale Collection. palawanensis — Criniger palawanensis Tweeddale, 1978	1884.7.31.198. Pacasmayo, Peru, 4 June 1877. Collected by J.
Proc. zool. Soc. Lond., 1878: 616. HOLOTYPE, Adult female. Reg. no. 1887.11.20.514. Puerto Princesa, Palawan, Philippines, Jan. 1878. Collected by A. H. Everett. Tweeddale Collection. palawanensis — Criniger palawanensis Tweeddale, 1978 21 Proc. zool. Soc. Lond., 1878: 618. SYNTYPE, Adult male. Reg. no. 1888.4.1.440. Puerta Princesa, Palawan, Philippines, Jan. 1878. Collected by A. Everett. Tweeddale Collection. The female syntype is also in the collection. palawanensis — Dicrurus palawanensis Tweeddale, 1878 64 Proc. zool. Soc. Lond., 1878: 614. SYNTYPE, Adult female. Reg. no. 1887.11.20.739. Puerto Princesa, Palawan, Philippines, Dec. 1877. Collected by A. H. Everett. Tweeddale Collection. There were several syntypes, a male being sent to Dresden Museum by the author. palawanensis — Mainatus palawaneensis Sharp, 1890 62 Cat. Bds. Brit. Mus., 13: 104. SYNTYPE, Adult female. Reg. no. 1888.9.20.320. Puerto Princesa, Palawan, Philippines, 3 Dec. 1877. Collected by A. H. Everett. Tweeddale Collection. The immature syntype is also in the collection. The immature syntype is also in the collection. palawanensis — Phyllornis palawanensis Sharp, 1876	Report of the Expedition for exploring central Africa, 1834: Appendix: 46. Syntype, Adult male (relaxed mount). Reg. no. 1838.6.9.95. North of Kuruman, Latakoo. Collected by A. Smith and purchased of the South African Museum.
Proc. zool. Soc. Lond., 1878: 618. SYNTYPE, Adult male. Reg. no. 1888.4.1.440. Puerta Princesa, Palawan, Philippines, Jan. 1878. Collected by A. Everett. Tweeddale Collection. The female syntype is also in the collection. Palawanensis — Dicrurus palawanensis Tweeddale, 1878	Proc. zool. Soc. Lond., 1878:616. HOLOTYPE, Adult female. Reg. no. 1887.11.20.514. Puerto Princesa, Palawan, Philippines,
Proc. zool. Soc. Lond., 1878: 614. SYNTYPE, Adult female. Reg. no. 1887.11.20.739. Puerto Princesa, Palawan, Philippines, Dec. 1877. Collected by A. H. Everett. Tweeddale Collection. There were several syntypes, a male being sent to Dresden Museum by the author. palawanensis — Mainatus palawaneensis Sharp, 1890 62 Cat. Bds. Brit. Mus., 13: 104. SYNTYPE, Adult female. Reg. no. 1888.9.20.320. Puerto Princesa, Palawan, Philippines, 3 Dec. 1877. Collected by A. H. Everett. Tweeddale Collection. The immature syntype is also in the collection. palawanensis — Phyllornis palawanensis Sharpe, 1876	Proc. zool. Soc. Lond., 1878: 618. SYNTYPE, Adult male. Reg. no. 1888.4.1.440. Puerta Princesa, Palawan, Philippines, Jan. 1878. Collected by A. Everett. Tweeddale Collection.
Cat. Bds. Brit. Mus., 13: 104. SYNTYPE, Adult female. Reg. no. 1888.9.20.320. Puerto Princesa, Palawan, Philippines, 3 Dec. 1877. Collected by A. H. Everett. Tweeddale Collection. The immature syntype is also in the collection. The immature syntype is also in the collection. palawanensis — Phyllornis palawanensis Sharpe, 1876	Proc. zool. Soc. Lond., 1878: 614. SYNTYPE, Adult female. Reg. no. 1887.11.20.739. Puerto Princesa, Palawan, Philippines, Dec. 1877. Collected by A. H. Everett. Tweeddale Collection. There were several syntypes, a male being sent to Dresden
Nature, 14: 297. SYNTYPE, (Adult female, vide Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 33). Reg. no. 1876.8.28.7. Palawan, Philippines, 1874. Collected by and purchased of J. B. Steere. See Sharpe, Trans. Linn. Soc. Lond. (Zool.) 2nd ser., 1 (1877): 333 for further notes on the type material. palawanorum — Dicaeum pygmaeum palawanorum Hachisuka, 1926. 45 Bull. Brit. Orn. Cl., 47: 55. Holotype, Adult male. Reg. no. 1911.11.16.305. Iwahig, Palawan, Philippines, 25 June 1907. Collected and presented by W. P. Lowe. pallatangae — Elainea pallatangae P. L. Sclater, 1862 Proc. zool. Soc. Lond., 1861 (publ. 1862): 407, pl. 41. Syntype, Adult female. Reg. no. 1888.1.13.601. Pallatanga, Ecuador, (Nov. 1858, on label). Collected by L. Fraser. Sclater Collection.	Cat. Bds. Brit. Mus., 13: 104. SYNTYPE, Adult female. Reg. no. 1888.9.20.320. Puerto Princesa, Palawan, Philippines, 3 Dec. 1877. Collected by A. H. Everett. Tweeddale Collection.
Bull. Brit. Orn. Cl., 47:55. HOLOTYPE, Adult male. Reg. no. 1911.11.16.305. Iwahig, Palawan, Philippines, 25 June 1907. Collected and presented by W. P. Lowe. pallatangae — Elainea pallatangae P. L. Sclater, 1862	Nature, 14: 297. SYNTYPE, (Adult female, vide Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 33). Reg. no. 1876.8.28.7. Palawan, Philippines, 1874. Collected by and purchased of J. B. Steere. See Sharpe, Trans. Linn. Soc. Lond. (Zool.) 2nd ser., 1 (1877): 333
Proc. zool. Soc. Lond., 1861 (publ. 1862): 407, pl. 41. SYNTYPE, Adult female. Reg. no. 1888.1.13.601. Pallatanga, Ecuador, (Nov. 1858, on label). Collected by L. Fraser. Sclater Collection.	Bull. Brit. Orn. Cl., 47:55. HOLOTYPE, Adult male. Reg. no. 1911.11.16.305. Iwahig, Palawan, Philippines, 25 June 1907.
	Proc. zool. Soc. Lond., 1861 (publ. 1862): 407, pl. 41. SYNTYPE, Adult female. Reg. no. 1888.1.13.601. Pallatanga, Ecuador, (Nov. 1858, on label). Collected by L. Fraser. Sclater Collection.

- pallescens Anthus pallescens Vigors & Horsfield, 1827 . . . 19
 Trans. Linn. Soc. Lond., 15:229. HOLOTYPE, (Immature, vide Sharpe,
 Cat. Bds. Brit. Mus., 10(1885):616) (mounted). Reg. no. 1863.7.7.
 31. Australia. Presented by the Linnean Society.
- pallescens Cercomacra tyrannina pallescens Chubb, 1918. . . . 4
 Bull. Br. Orn. Cl., 38:85. SYNTYPE, Adult male. Reg. no. 1889.
 9.20.412. Esmeraldas, Ecuador, Nov. 1859. Collected by L. Fraser. P. L. Sclater Collection.
 The female syntype is in the collection.
- pallescens Stachyridopsis rufifrons pallescens Ticehurst, 1932 . 32B Bull. Br. Orn. Cl., 53:18. HOLOTYPE, Adult female. Reg. no. 1932.10.20.1. Gamon Chaung, 500 ft., Sandoway District, Arakan, Burma, 9 Feb. 1930. Collected by E. Henrichs and presented by J. K. Stanford.
- palliata—Plagiospiza superciliosa palliata van Rossem, 1938 . . . 49
 Bull. Brit. Orn. Cl., 58: 127. Holotype, Adult male. Reg. no. 1899.2.1.3591. Tutuaca, 9,500', Chihuahua, Mexico, 30 March 1888. Collected by W. Lloyd. Salvin-Godman Collection.
- palliata Vermivora superciliosa palliata van Rossem, 1939 . . . 54
 Proc. Biol. Soc. Wash., 52: 12. HOLOTYPE, Adult male. Reg. no. 1892.4.1.338. Sierra Nevada de Colima Jalisco, Mexico, 7 April 1889. Collected by W. B. Richardson (W. Lloyd, on label). Salvin-Godman Collection.

There is also a female syntype in the collection. Renamed Galerida cristata superflua by Hartert (q.v.) on the grounds of preoccupation.

pallida — Cactornis pallida P. L. Sclater and Salvin, 1870 53

Proc. zool. Soc. Lond., 1870 : 327. SYNTYPE, Adult (male, vide Sharpe, Cat. Bds. Brit. Mus., 12 (1888) : 21). Reg. no. 1885.12.

nudum.

pallida — Otocorys pallida Sharpe, 1890.

Collected by A Habel. Salvin-Godman Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 21. There is another syntype in the collection.
pallida — Chalcoparia singalensis pallida Chasen, 1934
pallida — Drymoica pallida Smith, 1842
pallida — Mirafra pallida Sharpe, 1902
pallida — Molpastes haemorrhous pallida E. C. S. Baker, 1917 . 21 Bull. Br. Orn. Cl., 38: 15. SYNTYPE, Adult male. Reg. no. 1886.9.1. 1722. Deesa, Bombay, 16 March 1876. Collected by E. A. Butler. Hume Collection. This is the author's selected male. His selected female syntype is also in the collection.
pallida — Munia pallida Wallace, 1864
pallida — Myiagra pallida Ramsay, 1879
of the types. Gould, Birds of New Guinea, 2 (1884): pl. 36, implies that the type was at Sydney, but no holotype was designated in the original

description. The reference in Nature, 20 (1879): 125 is a nomen

Cat. Bds. Brit. Mus., 13:533. SYNTYPE, Adult male. Reg. no. 1890.1.29.54. Kashgar, east Turkestan, Dec. 1873. Collected by J. Biddulph. Seebohm Collection. (cont.)

17

14.562. Indefatigable Island, Galapagos Archipelago, 9 Sept. 1868.

- There are several other syntypes in the collection. Renamed O. diluta Sharpe, 1890 on the grounds of preoccupation by O. alpestris pallida Dwight, 1890.

- pallidiceps Ploceus aureoflavus pallidicpes Vincent, 1933 . . . 61

 Bull. Br. Orn. Cl., 53: 147. Holotype, Adult male. Reg. no. 1933.3.1.17. Mocuba, 700', ("near Lugela River ferry" on label) Quelimane Province, Portuguese East Africa, 1 Feb.1932. Collected and presented by J. Vincent.
- pallidigula Xenocichla pallidigula Sharpe, 1897 . 2

 Bull. Br. Orn. Cl., 7:7. HOLOTYPE, Adult male. Reg. no. 1897.10.31.4. Ntebi, Uganda, 3 Aug. 1895. Collected and presented by F. J. Jackson.
- pallidior Culicicapa ceylonensis pallidior Ticehurst, 1927 . . . 37A Bull. Brit. Orn. Cl., 47: 108. HOLOTYPE, Adult male. Reg. no.

1886.4.1.3089. Simla, India, 1 Oct. 1880. Collected by W. Davison. Hume Collection.
pallidior — Estrilda pallidior Jackson, 1910 60 Bull. Brit. Orn. Cl., 27:6. HOLOTYPE, Adult male. Reg. no. 1910.10.8.3. Northern Uaso, (3,000', on label), Nyero, Kenya, 10 Aug. 1909. Collected and presented by F. J. Jackson.
pallidior — Neomixis striatigula pallidior Salomonsen, 1934 321 Ann. Mag. nat. Hist., (10) 14:65. HOLOTYPE, Adult male. Reg. no. 1931.8.18.2928. Ampotaka, South Madagascar, 20 March 1930. Collected on the Archbold Vernay Expedition.
pallidior — Philydor atricapillus pallidior Chubb, 1919
pallidior — Pnoepyga albiventer pallidior Kinnear, 1924 321 Bull. Brit. Orn. Cl., 45: 10. HOLOTYPE, Adult male. Reg. no. 1924.10.9.1. Dharmsala, 4,000', Punjab, 1 Dec. 1921. Collected and presented by H. Whistler.
pallidior — Poliospiza pallidior Lort Phillips, 1898 59 Ibis, (7) 4:398. SYNTYPE, Adult male. Reg. no. 1898.7.27.21. Wagga, Somaliland, 27 Feb. 1897. Collected by and purchased of E. Lort Phillips. This specimen was labelled "Type" and is the only definite male of the type series. There are four other syntypes in the collection.
pallidiora — Ianthia cyanura pallidiora E. C. S. Baker, 1924 3 The Fauna of British India; Birds (2nd ed.), 2:101. HOLOTYPE, Adult female. Reg. no. 1886.4.1.2144. Simla, (29 Nov. 1880 on label). Collected by A. O. Hume. Hume Collection.
pallidipes — Phylloscopus pallidipes Blanford, 1872 3. J. Asiat. Soc. Bengal, 41, no. 2:162, pl. 7. HOLOTYPE, Adult. Reg. no. 1876.10.17.3. Sikkim. Collected and presented by L. Mandelli.
pallidirostris — Collyriocincla pallidirostris Sharpe, 1877 39 Cat. Bds. Brit. Mus., 3:293. SYNTYPE, Adult (relaxed mount). Reg. no. 1842.1.13.35. Port Essington, Northern Australia, 13 May 1840. Collected and presented by W. Chambers. The second syntype is also in the collection.
pallidiventris — Anthus pallidiventris Sharpe, 1885

Sharpe.

There are two other syntypes in the collection.

pallidus — Criniger pallidus Swinhoe, 1870
Ibis, (2) 6: 252. SYNTYPE, Adult. Reg. no. 1898.10.2.90. Hainan, Feb. 1868. Collected by R. Swinhoe. Seebohm Collection.
pallidus — Drymaoedus pallidus Sharpe, 1883
pallidus — Eurylaimus javanicus pallidus Chasen, 1935
pallidus — Horeites pallidus Brooks, 1871
pallidus — Myiobius pallidus Gosse, 1847
pallidus — Pericrocotus peregrinus pallidus E. C. S. Baker, 1920 . 20 Bull. Br. Orn. Cl., 40:115. Holotype, Adult male. Reg. no. 1886.4.1.483. Near Larkhana, Sind, India, 4 Feb. 1871. Collected by A. O. Hume. Hume Collection.
pallidus — Tephrodornis pondicerianus pallidus Ticehurst, 1920 . 20 Bull. Br. Orn. Cl., 41:56. HOLOTYPE, Adult male. Reg. no. 1886.3.1.2423. Larkhana, Sind, 4 Feb. 1871. Hume Collection.
pallidus — Troglodytes pallidus Hume, 1875
palliseri — Brachypteryx palliseri Blyth and Kelaart, 1851 35 J. Asiat. Soc. Bengal., 20: 178. SYNTYPE, Adult female. Reg. no. 1852.11.26.17. Ceylon. (Nuwera Elia and Dimboola, vide Layard, Ann Mag. nat. Hist., (2) 12 (1853): 269). Collected and presented by Dr Kelaart.

palmae — Fringilla palmae Tristram	, 1889 .			. 58
Ann. Mag. nat. Hist., (6) 3: 489.	SYNTYPE,	Immature	male.	Reg.
no. 1905.12.22.158. Palma, (San	ta Cruz, 3,0	000', on lal	bel), Ca	anary
Islands, 14 April 1889. Collecte	ed by H. B.	Tristram a	and E.	G. B.
Meade-Waldo. Presented by E.	G. B. Mea	de-Waldo.		

There were many syntypes. Tristram, Catalogue of a collection of birds, (1889): 226, stated that "types" were in his collection. This is the "juvenile" described, and there are also three adult male syntypes in the collection, and a male and a female in the University Museum

of Zoology, Cambridge.

University Museum of Zoology, Cambridge.

Gould used Jerdon's MS. name. It seems almost certain that this specimen is the one given by Jerdon to Hume, who described it in Stray Feathers, 1 (1872): 4, but, in a footnote, accepted Gould's description as the earlier. As there is no specimen in the collection from Gould, the latter presumably returned to Jerdon the specimen he used in the original description.

palustris — Megalurus palustris Horsfield, 1821 35

Trans. Linn. Soc. Lond., 13: 159. SYNTYPE, Adule male (relaxed mount). Reg. no. Old Vellum Cat. 14, 412 a. Java. Collected by T. Horsfield. Presented by the Honourable East India Company Museum.

This specimen was regarded as the type when relaxed as a mounted specimen in 1883, but other specimens were also available to the author at the time of the original description.

Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 270, refers to types in Seebohm's possession. This specimen and a female are both marked

^ *
as "type" on the label but are part of an original large syntypical series.
pammelaena — Myzomela pammelaena P. L. Sclater, 1877
panayensis — Artamides panayensis Steere, 1890
panayensis — Edolisoma (Graucalus) panayensis Steere, 1890 20 List of Birds and Mammals collected by the Steere Expedition to the Philippines: 14 (reprinted in Ibis, (6) 3 (1891): 310). SYNTYPE, Adult male. Reg. no. 1896.6.6.64. Conception, Panay, Philippines, 17 Jan. 1888. Collected by and purchased of J. B. Steere. This is the author's selected male. His selected female and two other syntypes are also in the collection.
panayensis — Orthotomus panayensis Steere, 1890 39 List of Birds and Mammals collected by the Steere Expedition to the Philippines: 20. HOLOTYPE, Adult (female, on label). Reg. no. 1896.6.6.240. Panay, Philippines, (Conception, 12 Jan., on label). Collected by J. B. Steere. Steere Collection.
pandoo — Petrocincla pandoo Sykes, 1832
panerythrus — Philydor panerythrus P. L. Sclater, 1862
pannosa — Sylvicola pannosa Gosse, 1847
papua — Hylochelidon nigricans papua Mathews, 1928 18

- papuenis Chaetorhynchus papuensis A. B. Meyer, 1874

 Sitzungsb. K. Akad. Wien, math.-naturwiss., 69 (1): 493. Syntype,
 Adult female. Reg. no. 1878.10.23.1. Arfak Mountains, c. 3,550',
 New Guinea, July 1873. Collected by A. B. Meyer. Purchased of
 E. Gerrard as duplicates of the Dresden Museum.

 Listed as "one of the Typical specimens" by Sharpe, Cat. Bds.
 Brit. Mus., 3 (1877): 243.

- paraguayensis Dacnis cayana paraguayensis Chubb, 1910 51 Ibis, (9) 4: 619. SYNTYPE, Adult. Reg. no. 1905.10.12.803. Sapucay, Paraguay, 6 June 1904. Collected by and purchased of W. Foster.

Listed as the type by Hellmayr and Conover, Catalogue of birds of the Americas, 8 (1935): 270.

paraguayensis — Embernagra platensis paraguayensis Chubb, 1918
 Ibis, (10) 6:5. SYNTYPE. Adult male. Reg. no. 1903.12.22.90.
 Sapucay, Paraguay, 19 April 1903. Collected by W. Foster, and purchased of Miss A. Foster.

This is the syntype measured for the description, and listed as the type by Hellmayr and Conover, *Catalogue of birds of the Americas*, 11(1938): 635. There are several other syntypes in the collection.

paraguayensis — Merula albicollis paraguayensis Chubb, 1910. . . 31 Ibis, (9) 4:608. Syntype, Adult male. Reg. no. 1905.10.12.276. Sapucay, Paraguay, 7 June 1904. Collected by and purchased of W. Foster.

There are several other syntypes in the collection.

paraguayensis — Nemosia pileata paraguayensis Chubb, 1910 . . . 51
 Ibis, (9) 4:629. Syntype, Adult male. Reg. no. 1905.10.12.850.
 Sapucay, Paraguay, 2 May 1904. Collected by and purchased of W. Foster.

This specimen was labelled as the type by the author, and listed as such by Hellmayr and Conover, *Catalogue of birds of the Americas*, 9 (1936): 370.

Sapucay, Paraguay, 6 July 1904. Collected and presented by W. Foster.

There are eight other syntypes in the collection.

paraguayensis—Sporophila pileata paraguayensis Chubb, 1910 . 59 Ibis, (9) 4:634. Syntype, Adult male. Reg. no. 1905.10.12.878. Sapucay, Paraguay, 24 March 1903. Collected and purchased of W. Foster.

The other two syntypes are also in the collection.

- paranensis Lepthasthenura paranensis P. L. Sclater, 1861 3

 Proc. zool. Soc. Lond., 1861 : 377. HOLOTYPE, Adult. Reg. no. 1889.5.20.104. Parana, Argentina (error; Mendoza, vide Hellmayr, Catalogue of Birds of the Americas, 4 (1925) : 69). Collected by H. Burmeister. Sclater Collection.
- pardus Campylorhynchus pardus P. L. Sclater, 1858 28

 Proc. zool. Soc. Lond., 1857 (publ. 1858): 271, HOLOTYPE, Adult.

 Reg. no. 1886.9.15.41. Santa Martha, Colombia. Collected for G. N. Lawrence. Sclater Collection.
- parkini Passer domesticus parkini Ticehurst, 1920 6

 Bull. Br. Orn. Cl., 41:13. HOLOTYPE, Adult male. Reg. no. 1949.1.9830. Srinagar, 5,000', Cashmere, 5 April 1920. Collected and presented by H. Whistler.

The type was stated to be in the British Museum in 1920 but was actually received, with the Whistler collection, in 1949.

- parsoni—Hirundo tahitica parsoni C. M. N. White, 1936 . . . 18

 Bull. Br. Orn. Cl., 57:27. HOLOTYPE, Adult male. Reg. no.
 1901.3.20.85. Bartle Frere, Queensland, 30 May 1900. Collected and presented by H. C. Robinson.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 605. There are two other syntypes in the collection.

- parvirostris Jabouilleia danjoui parvirostris Delacour, 1927 . . . 32B Bull. Br. Orn. Cl., 47:160. Holotype, Adult male. Reg. no. 1927.6.5.15. Bana, central Annam, 14 Aug. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China.

parvirostris — Myiobius parvirostris Gould, 1839
Zoology of the voyage of H.M.S. Beagle, 3, Birds: 48, SYNTYPE, Adult (relaxed mount). Reg. no. 1856.3.15.16. Tierra Del Fuego,
Chile, and La Plata (Santa Cruz, Patagonia, on label: queried by
Hellmayr, Catalogue of birds of the Americas, 5(1927): 401, footnote).
(The date 4 Jan. 1837 on the label appears to refer to the date when
Gould received it). Collected and presented by C. Darwin.
Gould prepared the description, but the generic name, originally <i>Tyrannula</i> , was changed by G. R. Gray when the text was edited
by Darwin.
parvirostris — Parus afer parvirostris Shelley, 1900
Birds of Africa, 2:223, 241. HOLOTYPE, Adult male. Reg. no.
1896.3.15.31. Salisbury, Mashonaland, Southern Rhodesia, 12
April 1895. Collected and presented by G. A. K. Marshall. parvirostris — <i>Picolaptes parvirostris</i> P. L. Sclater, 1889
Proc. zool. Soc. Lond., 1889: 33 HOLOTYPE, (Immature, vide
Proc. zool. Soc. Lond., 1889: 33 HOLOTYPE, (Immature, vide Hellmayr, Catalogue of Birds of the Americas, 4 (1925): 325).
South-east Brazil (locality no doubt erroneous, vide Hellmayr, loc.
cit). Salvin-Godman Collection.
parvirostris — Polioptila parvirostris Sharpe, 1885
Cat. Bds. Brit. Mus., 10: 448. HOLOTYPE, Adult male. Reg. no. 1881.5.1.2204. Chamicuros, Amazon River, Eastern Peru. Col-
lected by J. Hauxwell. Gould Collection.
parvirostris — Pyrocephalus parvirostris Gould, 1839
Zoology of the voyage of H.M.S. Beagle, 3, Birds: 46, pl. 6. Syn-
TYPE, Adult male (relaxed mount). Reg. no. 1856.3.15.17. Buenos
Aires. Collected and presented by C. Darwin. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 213.
parvissima — Colluricincla parvissima Gould, 1872
Ann. Mag. nat. Hist., (4) 10:114. HOLOTYPE, Adult. Reg. no.
1881.5.1.451. Rockingham Bay, Queensland. Gould Collection.
parvula—Colluricincla parvula. See appendix, Corrections and
Additions, vol. 2
parvula — Geospiza parvula Gould, 1837
Proc. zool. Soc. Lond., 1837: 6. SYNTYPE, Adult male, (relaxed mount). Reg. no. 1855.12.19.194. Galapagos Archipelago, (James
Island, vide Zoology of the voyage of H.M.S. Beagle, 3, Birds,
Island, vide Zoology of the voyage of H.M.S. Beagle, 3, Birds, (1841): 102; Chatham Island, on label but presumably not
written until 1855). Collected by C. Darwin. Purchased of the
Zoological Society of London. There are two other syntypes in the collection. Swarth, Occ. Pap.
California Acad. Sci., 18 (1931): 229 confirms that the locality given
on the label – Chatham Island – is an error and attributes this to
Sharpe.
parvulus — Orpheus parvulus Gould, 1837
Proc. zool. Soc. Lond., 1837: 27. HOLOTYPE, Adult (relaxed mount).
Reg. no. 1855.12.19.92. (not 93 as labelled for many years; 93 is a

- specimen of *Mimus orpheus*). Galapagos Islands (Albemarle Island, on label). Collected by C. Darwin on the voyage of *H.M.S. Beagle* and purchased of the Zoological Society of London.
- parvus—Zosterops aureiventer parvus Hachisuka, 1926 47

 Bull. Brit. Orn. Cl., 47:56. HOLOTYPE, Adult male. Reg. no. 1898.9.30.223. Kina Balu, North Borneo, (1,000 ft,. on label), 30 March 1887. Collected by J. Whitehead. Seebohm Collection.
- passerina Mirafra passerina Gyldenstolpe, 1926 17

 Ark. Zool., 19: 24. Lectotype, Adult. Reg. no. 1876.5.23.704.

 Damaraland, Collected by C. J. Andersson. Sharpe Collection.

 Designated lectotype by C. H. B. Grant & Praed, Bull. Br. Orn.

 Cl., 69 (1949): 30.
- passerinus Ploceus passerinus Reichenow, 1886 . . . 61
 Zoologisches Jahrbuch, 1:156. New name proposed for Ploceus
 atrigula (q.v.) on the grounds of alleged preoccupation.
- patae Certhilauda albescens patae Macdonald, 1953. 17
 Bull. Brit. Mus. nat. Hist. (Zool.), 1:344. HOLOTYPE, Adult male.
 Reg. no. 1950.50.936. Grootderm, 400', Little Namaqualand,
 South-West Africa, 17 Dec. 1949. Presented by the South-West
 Africa Expedition.
- patagonicus—Orpheus patagonicus Lafresnaye and d'Orbigny, 1837 29 Mag. Zool., 7, cl. 2: 19. A specimen in the collection, B.M. Reg. no. 1855.12.19.311, collected by C. Darwin and purchased of the Zoological Society of London, was listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 352. G. R. Gray, in Darwin, Zoology of the Voyage of H.M.S. Beagle, 3, Birds (1839): 60, referred to d'Orbigny's specimens. According to Hellmayr, Catalogue of Birds of the Americas, 7 (1934): 322, the type is in the Paris Museum.
- paucimaculatus Thryothorus paucimaculatus Sharpe, 1881 . . . 28
 Cat. Bds. Brit. Mus., 6:238. HOLOTYPE, Adult. Rcg. no.
 1886.3.6.355. Balzar Mountains, Ecuador. Collected by
 Illingworth. Salvin-Godman Collection.

pearsoni — Dryodromas pearsoni Neave, 1909 3. Ann. Mag. nat. Hist., (8) 4:130. HOLOTYPE, Adult (male, vide, Ibis, (9) 4 (1910):151). Reg. no. 1909.12.31.206. Katanga, southeast Congo, (Lufupa River, 30 Oct. 1907, vide, Ibis, (9) 4 (1910):151). Collected and presented by S. A. Neave.
peasei — Estrilda peasei Shelley, 1903
pectoralis — Corvus pectoralis Gould, 1836
pectoralis — Dicrurus pectoralis Wallace, 1862 6 Proc. zool. Soc. Lond., 1862 : 342. SYNTYPE, Adult male. Reg. no. 1873.5.12.1971. Sula Islands, 1861. Collected by A. R. Wallace. Wallace Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 3 (1877) : 240. There is another syntype in the collection.
pectoralis — Drymoica pectoralis Smith, 1843 3 Illustrations of the Zoology of South Africa, Aves, : pl. 75, fig. 1. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1843.2.28.93. Latakoo. Collected by A. Smith. Purchased of the South African Society. The name is preoccupied by Malcorus pectoralis Smith, 1829 (q.v.), which appears on the same plate under the name of D. ocularius. The specimen does not appear to have been given a new name but was sunk into the synonymy of Prinia flavicans by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 186.
pectoralis — Erythropygia pectoralis Smith, 1836 3 Report of the Expedition for exploring central Africa, 1834; Appendix : 46. SYNTYPE, Adult male (relaxed mount). Reg. no. 1845.7.6.45. Between Orange River and Kurrichane. Collected by and purchased of A. Smith. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 75. There is a second syntype in the Edinburgh Museum, vide Scottish Naturalist, 1930: 150.
pectoralis — Gerygone pectoralis Davison, 1892
pectoralis — Glyciphila pectoralis G. R. Gray, 1869 4 Handlist of Birds, 1:155. This is a nomen nudem and the specimen listed as the type by Gadow, Cat. Bds. Brit. Mus., 9 (1884):212 has no type status.

- pectoralis Glyphorhynchus pectoralis P. L. Sclater and Salvin, 1860
 Proc. zool. Soc. Lond., 1860: 299. SYNTYPE, Adult. Reg. no. 1889.5.14.549. Choctum, Vera Paz, Guatemala, Jan. 1860. Collected by O. Salvin. Salvin-Godman Collection. There is another syntype in the collection.
- pectoralis Graucalus pectoralis Jardine and Selby, 1828 . . . 20 Illustrations of Ornithology, 1: pl. 57. HOLOTYPE, Adult male. Reg. no. 1886.6.24.115. Sierra Leone. Jardine Collection; purchased of E. Gerrard.
- pectoralis Herpsilochmus pectoralis P. L. Sclater, 1857.

 Proc. zool. Soc. Lond., 1857: 132. Sclater mentioned two specimens, one in the Academy of Natural Sciences, Philadelphia, from the Massena Collection, and another unlocalised specimen in the British Museum, thought also to be from the Massena Collection. The latter syntype cannot now be traced in the collection. The Wucherer specimen from Bahia, listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 247, is not accepted as syntypical material.

See also Sclater, Ibis, (4) 5 (1881): 298.

Smith redescribed this in *Illustrations of the Zoology of South Africa*, Aves, (1843): pl. 75., under the name of Drymoica ocularius possibly confusing it with the other specimen illustrated which he described as D. pectoralis (q.v.). The original description was reprinted in Ann. Transvaal Mus., 22 (1953): 201.

drawings Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 134.
Vol. 1, no. 50; vol. 2, no. 53 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 8(1920): 212.
pectoralis — Nectarinia pectoralis Horsfield, 1821 4 Trans. Linn. Soc. Lond., 13:167. SYNTYPE, Adult male. Reg. no. 1880.1.1.1998. Java. Collected by T. Horsfield. Presented by the Honourable East India Company Museum. There are other syntypes in the collection.
pectoralis — Oreocincla pectoralis Legge, 1876
pectoralis — Pellorneum pectoralis Godwin-Austen, 1877 321 J. Asiat. Soc. Bengal, 46, pt. 2:41. Syntype, Adult male. Reg. no. 1895.7.14.1723. (Manbum, on label), Saddya, Assam, 25 Jan. 1877. Collected by M. J. Ogle. Godwin-Austen Collection. This is the author's selected male. His selected female and two other syntypes are also in the collection.
peguensis — Alauda peguensis Oates, 1875
peguensis — Ploceus manyar peguensis E. C. S. Baker, 1925 6 Bull. Br. Orn. Cl., 45:58 HOLOTYPE, Adult male. Reg. no. 1897.12.10.717. Pegu, Berma, ('19 Aug. 1880' on label). Collected by E. W. Oates. Presented by J. Biddulph.
peguensis — Zosterops palpebrosa peguensis E. C. S. Baker, 1922 . 4 Ibis, (11) 4: 144. HOLOTYPE, Adult male. Reg. no. 1886.12.1.1700. Moulmein, Southern Burma, 4 Dec. 1878. Collected by W. Davison. Hume Collection.
pekinensis — Alauda pekinensis Swinhoe, 1863
pekinensis — Drymoeca (?) pekinensis Swinhoe, 1868 3 Ibis, (2) 4:62. HOLOTYPE, Adult female. Reg. no. 1898.10.3.126. Pekin, China, (7 April, 1867, on label). Collected by Pére David. Seebohm Collection.

pekinensis — Parus pekinensis David, 1870	2
pellotis — Timalia pellotis Hodgson, 1836	В
pelopus — Anthus pelopus J. E. Gray, 1846	9
pelzelni — Euphonia nigricollis pelzelni P. L. Sclater 1886 5 Cat. Bds. Brit. Mus., 11:61. SYNTYPE, Adult male. Reg. no. 1885.6.12.36. Govinda, West Ecuador. Collected by C. Buckley. Sclater Collection, having apparently been purchased through Gerrard by Berlepsch and given to Sclater (vide specimen label). Sclater used a Berlepsch MS. name. There are two other syntypes in the collection.	1
pelzelni — Euscarthmus pelzelni P. L. Sclater and Salvin, 1881. Ibis, (4) 5: 268. HOLOTYPE, Immature male (vide Hellmayr, Catalogue of birds of the Americas, 5 (1927): 319). Reg. no. 1888.1. 13.273. Cuyaba, Matto Grosso, Brazil, 16 June. Collected by Natterer. Sclater Collection.	9
pelzelni—Granatellus pelzelni P. L. Sclater, 1865 5. Proc. zool. Soc. Lond., 1864 (publ. 1865): 600, pl. 37. HOLOTYPE, Adult male. Reg. no. 1884.5.15.516. Madiera River, Brazil, 19 Sept. 1829. Collected by J. Natterer. Sclater Collection, received by him in exchange with the Vienna Museum.	4
pelzelni — Myiarchus pelzelni Berlepsch, 1883	9
pelzelni — Myrmeciza pelzelni P. L. Sclater, 1890	4

1889.9.20.497. Marabitanas, Rio Negro, Brazil, March 1831. Collected by J. Natterer. Sclater Collection. There are several other syntypes in the collection.
pelzelni — Sycalis pelzelni P. L. Sclater, 1872
pelzelni — Tityra pelzelni Salvin and Godman, 1890
pembae — Cyanomitra olivacea pembae Granvik, 1934 40 Revue Zool. Bot. Afr., 25: 142. HOLOTYPE, Adult male. Reg. no. 1926.5.2.19. Pemba Island, Tanganyika, 27 May 1925. Collected and presented by J. H. Vaughan.
penduliger — Cephalopterus penduliger P. L. Sclater, 1859 Ibis, (1) 1:114, pl. 3. Syntype, Adult male. Reg. no. 1888.1.13. 1686. Pallatanga, Ecuador. Collected by L. Fraser. Sclater Collection. There is another syntype in the collection.
penicillata — Alauda penicillata Gould, 1837
penicillatus — Pycnonotus penicillatus Kelaart, 1851
peninsulae — Trochalopterum peninsulae Sharpe, 1887 32E Proc. zool. Soc. Lond., 1887 : 436, pl. 37. Syntype, Adult male. Reg. no. 1886.12.22.12. Larut Range, c, 4.400', Perak, Malaya, Sept.—Oct. 1886. Collected and presented by L. Wray. The female syntype is also in the collection.
peninsularis — Cyornis peninsularis Robinson, 1909 3' J. Fed. Malay Sts. Mus., 2:163. SYNTYPE, Adult male. Reg. no. 1936.4.12.2951. Telôm, 3,800', Perak-Pahang Boundary, South Perak, Malaya, 28 Nov. 1908. Collected by H. C. Robinson and C. B. Kloss. Presented by the Raffles Museum, Singapore. The female syntype is also in the collection

- peninsularis Dicrurus macrocercus peninsularis Ticehurst, 1932 . 64

 Bull. Brit. Orn. Cl., 53: 20. HOLOTYPE, Adult male. Reg. no.
 1932.10.19.1. Madras City, India, 7 May 1931. Collected for the
 Madras Museum. Presented by H. Whistler.
- peninsularis Melanocichla peninsularis Sharpe, 1888 32B Proc. zool. Soc. Lond., 1888 : 274. HOLOTYPE, Adult male. Reg. no. 1888.3.18.42. Gunong Batu Putch, 3,400', Perak, Malaya. Collected and presented by L. Wray.
- peracensis Bhringa remifer peracensis E. C. S. Baker, 1918 . . . 64
 Bull. Brit. Orn. Cl., 39: 18. HOLOTYPE, Adult male. Reg. no.
 1905.2.1.248. Telôm, Perak-Pehang Border, Malaya. 20 Jan. 1902.
 Collected by H. C. Robinson. Presented by the Royal Society.

- percivali Rhynchostruthus percivali Ogilvie-Grant, 1900 . . . 59
 Bull. Brit. Orn. Cl., 11:30. Holotype, Adult male. Reg. no.
 1900.12.16.1. Yeshbun, Hadramut, 150 miles north-west of
 Aden, South Arabia, 17 Jan. 1900. Collected by G. W. Bury.
 Presented by A. Blayney Percival.
- percivali Telephonus percivali Ogilvie-Grant, 1900 23

 Bull. Br. Orn. Cl., 10:50. HOLOTYPE, Adult male. Reg. no. 1900.8.5.231. Arabia, (Al Milah, north of Lahej, 16 Aug. 1899 on label). Collected by W. Dodson on the Percival-Dodson Expedition.

- periophthalmicus Monarcha periophthalmicus Sharpe, 1882 . . . 38 J. Linn. Soc. Lond., 16:318, 430. HOLOTYPE, Adult. Reg. no. 1882.3.8.59. Morocco District, Astrolabe Mountains, south-east New Guinea. Collected by A. Goldie and purchased of E. Gerrard.

- perplexa—Cisticola aridula perplexa C. M. N. White, 1947 . . . 35 Ostrich, 18:174. HOLOTYPE, Adult male. Reg. no. 1909.11.21. 170. Chambezi Valley, (3,900', on label), Northern Province, Northern Rhodesia, 25 Oct. 1908. Collected by and purchased of S. A. Neave.
- perplexus Phylloscopus armandii perplexus Ticehurst, 1934 . . . 35

 Bull. Brit. Orn. Cl., 54:96. Holotype, Adult male. Reg. no. 1921.7.5.399. Chien-Chuan Valley, north-west Yunnan, 26°40'N., May 1918. Collect by G. Forrest. Presented by Col. Stephenson Clarke.
- perplexus Pycnonotus simplex perplexus Chasen and Kloss, 1929. 21 J. Orn., 1929 (2): 116. HOLOTYPE, Adult male. Reg. no. 1947.60. 47. Balambangan Island, British North Borneo, 12 Sept. 1927. Collected by F. N. Chasen and C. B. Kloss and presented by the Raffles Museum, Singapore.

persieus 427
perplexus — Spinus pinus perplexus van Rossem, 1938 59 Bull. Brit. Orn. Cl., 58: 134. HOLOTYPE, Adult female. Reg. no. 1899.2.1.2116. San Andres, Chiapas, Mexico, 11 May 1877. Collected by W. B. Richardson. Salvin-Godman Collection.
perretti — Aethopyga mystacalis perretti Harrisson and Hartley, 1934
persepolis — Sturnus vulgaris persepolis Ticehurst, 1928 62 Bull. Br. Orn. Cl., 48:117. HOLOTYPE, Adult male. Reg. no. 1923.12.23.187. Baba Hazi, 19 miles south-east of Shiraz, Persia, 7 Oct. 1918. Collected by J. E. B. Hotson. Presented by the Bombay Natural History Society.
persica — Alaudula persica Sharpe, 1890
persica — Motacilla personata var. persica Blanford, 1876 19 The Zoology and Geology of Eastern Persia, 2:232 SYNTYPE, Adult male. Reg. no. 1874.11.23.60. Near Niriz, 5,000', east of Shiriz, Persia, 4 June 1872. Collected by W. T. Blanford of the Persian Boundary Commission and received in exchange with the Indian Museum, Calcutta. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 479. There are two other syntypes in the collection.
persica — Saxicola persica Seebohm, 1881
persica — Sitta earopaea persica Witherby, 1903
persicus — Parus (Cyanistes) persicus Blanford, 1873

W. L. Sclater. <i>Ibis</i> , (6)4(1892): 74, states that the type is at Calcutta, but four specimens were referred to and this is one of the syntypes.
personata — Apalis personata Sharpe, 1902
personata — Gerygone personata Gould, 1866 36 Proc. zool. Soc. Lond., 1866: 217. Holotype, Adult (male, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 230). Reg. no. 1868.1.27.19. Cape York, Queensland. Collected by J. Jardine for C. Coxan. Purchased of J. Gould.
personata — Haemophila personata Salvin, 1895
personata — Motacilla personata Gould, 1861
personata — Otocompsa personata Hume, 1873
personata — Spizocorys personata Sharpe, 1895
personata — Tityra personata Jardine and P. J. Selby, 1827 7 Illustrations of Ornithology I, pt. 2: pl. 24. HOLOTYPE, Adult male. Reg. no. 1886.6.24.472. Real del Monte, Mexico. (Guatemala, in B. M. register). Collected by Jenkins for W. Swainson who presented it to Jardine. Labelled "type of figure" in Jardine's hand.
perspicillata — Myiagra perspicillata G. R. Gray, 1859 38 Proc. zool. Soc. Lond., 1859: 161. HOLOTYPE, Immature male. Reg. no. 1859.4.19.12. Island of Nu, Port de France, New Calidonia. Collected by J. Macgillivray. Purchased of H. Cuming.
perspicillata — Zosterops perspicillata Shelley, 1889

- Reg. no. 1889.3.21.55. Kilimanjaro, 5,000-6,000, East Africa, 11 Aug. 1888. Collected and presented by M. C. V. Hunter. The male syntype is also in the collection.
- peruvianus Leptopogon peruvianus P. L. Sclater and Salvin 1868 . 9
 Proc. zool. Soc. Lond., 1867 (publ. 1868): 757. HOLOTYPE, Adult
 male. Reg. no. 1888.1.12.411. Chyavetas, East Peru. Collected
 by E. Bartlett. Sclater Collection.
- peruvianus Sclerurus mexicanus peruvianus Chubb, 1919 . . . 3

 Bull. Br. Orn. Cl., 39:41. HOLOTYPE, Adult female. Reg. no. 1889.5.14.485. Yurimaguas, east Peru, 19 March 1866. Collected by E. Bartlett. Salvin-Godman Collection.

- petrophila Cisticola petrophila Boyd Alexander, 1907. . . . 35

 Bull. Brit. Orn. Cl., 17: 104. Syntype, Adult male. Reg. no.
 1911.12.23.2465. Northern Nigeria, (Pettu, 27 August 1904, on label). Collected and bequeathed by W. Boyd Alexander.

 Labelled as the type; but there are several other syntypes in the collection.

- phaeocephalus Cyphorhinus phaeocephalus P. L. Sclater, 1860 . 28

 Proc. zool. Soc. Lond., 1860: 291. SYNTYPE, Adult (male, vide Sclater, Catalogue of American Birds, (1862): 19). Reg. no. 1886.9. 15.70. Ecuador (Esmeraldas, vide Sclater, 1862, loc. cit.). Collected by L. Fraser. Sclater Collection.

 The other, female, syntype is also in the collection.
- phaeocephalus Rhectes phaeocephalus von Reichenow, 1900 . 39

 Orn. Mber., 8:187. Holotype, Adult male. Reg. no. 1901.4.
 28.37. Builalla River, south-west New Guinea, Aug. 1898. Collected by and purchased of E. Weiske.
- phaeocercus Mitrephorus phaeocercus P. L. Sclater, 1859.

 Proc. zool. Soc. Lond., 1859: 44. SYNTYPE, Adult. Reg. no. 1888.1.13.917. Cordova, Vera Cruz, Mexico. Collected by A. Sallé. Sclater Collection.

 Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 219. There is a second syntype in the collection.
- phaeochromus Oriolus phaeochromus G. R. Gray, 1860 63

 Proc. zool. Soc. Lond., 1860 : 351. SYNTYPE, Adult. Reg. no. 1862.9.11.17. East Gililo (=Halmahera), Moluccas, 1859. Collected by A. R. Wallace. Wallace Collection.

 Listed as type by Sharpe, Cat. Bds. Brit. Mus., 3 (1877) : 212.
- phaeonotus Myiarchus phaeonotus Salvin and Godman, 1883. Ibis, (5) 1:207. Syntype, Adult male. Reg. no. 1888.1.1.1587. Merume Mountains, British Guiana, 8 June 1881. Collected by H. Whitely. Salvin-Godman Collection.

 There are several other syntypes in the collection.
- - W. L. Sclater, *Ibis*, (6) 4 (1892): 74, states that the type is in Calcutta. The above specimen was listed as the type by Gadow, *Cat. Bds. Brit. Mus.*, 8 (1883): 44. Three specimens were referred to in the original description.
- phaeopleurus Buarremon phaeopleurus P. L. Sclater, 1856. . . . 49
 Proc. zool Soc. Lond., 1856: 85. HOLOTYPE, Adult. Reg. No. 1885.6.12.890. Venezuela, 1848. Sclater Collection.

- phaeopleurus Catharus phaeopleurus P. L. Sclater and Salvin, 1875. 31 Proc. zool. Soc. Lond., 1875: 541. HOLOTYPE, Adult. Reg. no. 1886.8.2.18. Medellin, Antioquia, 1875. Collected by T. K. Salmon. Sclater Collection.
- phaeoptila—Phonipara phaeoptila Salvin and Godman, 1884. . . 50 Ibis, (5) 2:445. HOLOTYPE, Adult male. Reg. no. 1885.12.14.632. Roraima Mountains, 3,500', British Guiana (29 Nov. 1883, on label). Collected by H. Whitely. Salvin-Godman Collection.
- philbyi Calandrella blanfordi philbyi Bates, 1936.
 Bull. Brit. Orn. Cl., 56: 130. HOLOTYPE, Adult male. Reg. no. 1935.1.5.4. Ashaira (desert), 4,000', Arabia. 13 Jan. 1935. Collected and presented by H. Philby.

- philippensis Hyloterpe philippensis Walden, 1872 39

 Ann. Mag. nat. Hist., (4) 10: 252. HOLOTYPE, Adult male. Reg.
 no. 1888.5.30,337. Luzon, Philippines (Jan. 1872, on label).
 Collected by A. B. Meyer. Tweeddale Bequest.
- philippensis Poliolophus urostictus philippensis Hachisuka, 1934 21 Tori, 8:220. HOLOTYPE, Adult male. Reg. no. 1888.4.1.364. Dinagat, Mindanao, Philippines, June 1877. Collected by A. H. Everett. Tweeddale Collection.

Reg. no. 1876.8.28.8. Panay, Philippines. Collected by J. B. Steere and purchased of him as one of the types.

Listed as the type by Hachisuka, Birds of the Philippine Islands,

2 (1935): 292.

philippinensis—Mirafra philippinensis Wardlaw Ramsay, 1886 . 17
 Ibis, (5) 4:160. Syntype, Adult. Reg. no. 1888.9.20.1080.
 Manilla, Philippines. Collected by F. Maitland-Heriot. Tweed-dale Collection.

There are several other syntypes in the collection.

- phillipsi Dumetia hyperythra phillipsi Whistler, 1941 32B Ibis, (14) 5:319. Holotype, Adult male. Reg. no. 1940.12.3.10. Kumbalgamna, 3,000', Central Province, Ceylon, 8 Oct. 1937. Collected by E. C. Fernando of the British Museum.—Ceylon Ornithological Survey.

There are three other syntypes in the collection.

- phoeniceus Cardinalis phoeniceus Bonaparte, 1838 59

 Proc. zool. Soc. Lond., 1837 (publ. 1838) : 111. Bonaparte used a
 Gould MS, name and his type specimen is referred to by Hartert,
 Novit. Zool., 26 (1919) : 155 and is now presumably in the American
 Museum of Natural History. The Gould specimen listed as the
 type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888) : 167, appears to
 have no type status.
- phoenicomerus Euplectus phoenicomerus G. R. Gray, 1862 . . . 61

 Ann. Mag. nat. Hist., (3) 10:444. HOLOTYPE, Adult male. Reg. no. 1862.12.1.13. Cameroons, 7,000′, Collected by R. Burton and presented by Mrs Burton.

phoenicomitra — Myiobius phoenicomitra Taczanowski and Berlepsch, 1885
Proc. zool. Soc. Lond., 1885: 91. SYNTYPE, Adult male. Reg. no. 1888.1.13.857. Mapoto, Ecuador, 4 Feb. 1884. Collected by Stolzmann. Sclater Collection.
Taczanowski and Berlepsch used a Stolzmann MS. name. A female syntype is also in the collection.
phoenicura — Tyrannula phoenicura P. L. Sclater, 1855 9 Proc. zool. Soc. Lond., 1854 (publ. 1855): 113, pl. 66. ?HOLOTYPE, Adult. Reg. no. 1854.5.25.8. Rio Napo, eastern Ecuador. Collected for J. Gould. Gould Collection. Sclater described the species from a collection received by Gould. This specimen appears to have been part of that collection, was registered in the year that the description was made, and would seem to have been the only one available.
phoenicuroides — Bradypterus phoenicuroides J. E. Gray, 1846 . 31 Catalogue of Mammalia and Birds of Nepal and Thibet presented by B. H. Hodgson to the British Museum, Appendix: 70, 163. Syn- TYPE, Adult male (relaxed mount). Reg. no. 1845.1.9.433. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
phoenicuroides — Ruticilla phoenicuroides Moore, 1854 31 in Horsfield & Moore, Catalogue of Birds in the Museum of the Honourable East India Company, 1:301. SYNTYPE, Adult male (relaxed mount). Reg. no. 1879.12.28.701. Shikarpore, Afghanistan. Collected by W. Griffiths and presented by the Honourable East India Company Museum. Listed as type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881):343.
phoenicurus — Eremobius phoenicurus Gould, 1839 3 Zoology of the Voyage of H.M.S. Beagle, 3, Birds: 69, pl. 21. SYNTYPE, Adult (relaxed mount). Reg. no. 1855.12.19.73. Port Desire, Patagonia. Collected by Charles Darwin and purchased of the Zoological Society of London. Listed as type by P. L. Sclater, Cat. Bds. Brit. Mus., 15(1890): 27. There is at least one other syntype in the collection.
phrygius — Merops phrygius Latham, 1801
piaggiae — Turdus piaggiae Bouvier, 1877

Tana, Abyssinia, vide Jackson, Birds of Kenya and Uganda, 2 (1938)
: 954). Collected by Piaggia. Purchased of A. Bouvier. Listed as the type by Seebohm, <i>Cat. Bds. Brit. Mus.</i> , 5 (1881): 171.
picaecolor — Hemipus picaecolor Hodgson, 1845
picata — Muscicapa picata Sykes, 1832
picata — Melanodryas picata Gould, 1865
picata — Rhipidura picata Gould, 1848
picaum — Todirostrum picaum P. L. Sclater, 1858 9 Proc. zool. Soc. Lond., 1858 : 70. HOLOTYPE, Adult male. Reg. no. 1888.1.13.257. Upper Rio Napo, Ecuador. Collected by Don Villavicencio for Verreaux. Sclater Collection.
picta — Setophaga picta Swainson, 1829
picticollis — Garrulax picticollis Swinhoe, 1872
pictum — Todirostrum pictum Salvin, 1897
pieaoides — Sibia pieaoides Hodgson, 1839

pileata — Atticora pileata Gould, 1858
There is another syntype in the collection.
pileata — Leptasthenura pileata P. L. Sclater, 1881
pileata — Spermophila pileata P. L. Sclater, 1865
pileata — Timalia pileata Horsfield, 1821
India Company Museum. A second syntype was formerly in the collection, vide Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 509.
pileatus — Chlorospingus pileatus Salvin, 1865
pinetorum — Spizella pinetorum Salvin, 1863
pinicola — Turdus pinicola P. L. Sclater, 1859
pinwilli — Pomatorhinus pinwilli Sharpe, 1883 32B Cat. Bds. Brit. Mus., 7:413. SYNTYPE, Adult. Reg. no. 1876.2. 12.842. North-west Himalayas. Collected and presented by Capt. Stackhouse Pinwill. The immature syntype is also in the collection.
pipiens — Cisticola pipiens Lyncs, 1930

23 Feb. 1927. Collected by Admiral H. Lynes and B. B. Osmaston. Presented by Admiral H. Lynes.
placens — Elainea placens P. L. Sclater, 1859
placida — Xenocichla placida Shelley, 1889
placidus — Bradornis infuscatus placidus Clancey, 1958 3' Durban Mus. Novit., 5:130, 135. HOLOTYPE, Adult male. Reg. no. 1957.36.1. Kakia, 3,000', Southern Bechuanaland Protectorate, 24°55'S., 23°25'E., 12 June 1957. Collected by B. P. Hall on the Barlow 1957 Kalahari Expedition.
placidus — Calornis placidus Wallace, 1862 62 Proc. zool. Soc. Lond., 1862 : 343. HOLOTYPE, Adult male, Reg. no. 1873.5.12.1976. (Mesmon Island, Lat. 0°60,S., Long. 129°50'E., on label), Moluccas, 1860. Collected by A. R. Wallace. Wallace Collection. Wallace used a G. R. Gray MS. name.
plagosus — Dendrocolaptes plagosus Salvin & Godman, 1883 Ibis, (5) 1:210. SYNTYPE, Adult male. Reg. no. 1889.5.14.824. Camacusa, British Guiana, 8 April 1882. Collected by H. Whitely. Salvin-Godman Collection. The female syntype is in the collection.
platyrhyncha — Phyllomyias platyrhyncha P. L. Sclater and Salvin, 1873
Nomencl. Av. Neotrop., 1873: 158. Holotype, Adult (female, on label). Reg. no. 1888.1.13.656. Goiaz, Brazil (Aug. 1823, on label). Collected by Natterer. Sclater Collection.
pleurostictus — Thryothorus pleurostictus P. L. Sclater, 1860 28 Ibis, (1) 2: 30 HOLOTYPE, Adult. Reg. no. 1886.9.15.110A. Vera Paz, Guatemala. Collected by Skinner. Sclater Collection; obtained by Sclater from J. Gould.
plowesi — Fringillaria capensis plowesi Vincent, 1950

plumbea — Myiagra plumbea Vigors and Horsfield, 1827 Trans. Linn. Soc. Lond., 15:254. HOLOTYPE, Adult (male, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 374), (relaxed mount). Reg. no. 1863.7.7.47. Australia. Probably collected by Caley. Presented by the Linnean Society as the type. plumbea — Stenostira plumbea Hartlaub, 1858 Casamanca River, Senegambia. Purchased of Verreaux. Hartlaub used a Verreaux MS. name. Gray marked as type this specimen, which is probably the type of Verreaux's MS, name. Hartlaub refers to a specimen in Bremen but subsequently (vide J. Orn., 1861: 169) states that the British Museum specimen was omitted and should have been included. plumbeiceps — Staphida plumbeiceps Godwin-Austen, 1877 Ann. Mag. nat. Hist., (4) 20: 519. HOLOTYPE, Adult male. Reg. no. 1895.7.14.1862. Near Sadya, eastern Assam, 26 Jan. 1877. Collected by M. J. Ogle. Godwin-Austen Collection. plumbeitarsus — Phylloscopus plumbeitarsus Swinhoe, 1861 . Ibis, (1) 2:330. HOLOTYPE, Adult male. Reg. no. 1898.9.1.372. (near Pekin, vide Swinhoe, Proc. zool. Soc. Lond., 1863: 297) China. Collected by R. Swinhoe. Seebohm Bequest. plumbeus — Myrmelastes plumbeus P. L. Sclater, 1858 . Proc. zool. Soc. Lond., 1858: 274, pl. 143. SYNTYPE, Adult male. Reg. no. 1858.1.11.13. Rio Javarri, Upper Amazon, Brazil. Collected by H. W. Bates and purchased of S. Stevens. The female syntype is in the collection. plumigenis—Tropidorhynchus plumigenis G. R. Gray, 1858 Proc. zool. Soc. Lond., 1858: 174. SYNTYPE, Adult female. no. 1873.5.12.1015. Ké Islands, New Guinea, 1857. Collected by A. R. Wallace. Wallace Collection. The immature male syntype is also in the collection. plumosus — Pinarornis plumosus Sharpe, 1876 in Layard, Birds of South Africa, (2nd ed.): 230. HOLOTYPE, Adult. Reg. no. 1876.7.5.3. Victoria Falls, Zambezi (error, Matopos Hills, Matabeleland, vide Irwin, Bull. Br. Orn. Cl., 77 (1957): 9). Collected by and purchased of Cutter. poecilocerca — Serpophaga poecilocerca P. L. Sclater and Salvin, 1873 Nomencl. Av. Neotrop., 1873: 158. HOLOTYPE, Adult (female, on label). Reg. no. 1888.1.13.100. Puellaro, 8,000', Ecuador, (April 1859, on label). Collected by L. Fraser. Sclater Collection. poeciloma — Hirundo poeciloma Gosse, 1847 18

The Birds of Jamaica: 64. SYNTYPE, Adult. Reg. no. 1846.10.26.8.

Jamaica. Collected by P. H. Gosse. Gosse Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 196.
poecilopsis — Lophophanes poecilopsis Sharpe, 1902
poecilorhyncha — Garrulax poecilorhyncha Gould, 1862 32E Proc. zool. Soc. Lond., 1862: 281. SYNTYPE, Adult. Reg. no.
1898.10.3.635. (Forest ranges near Tamsuy vide Swinhoe, Ibis, (1) 5 (1863): 283), (April 1862, on label). Collected by R. Swinhoe. Seebohm Bequest. There are several other syntypes in the collection.
poecilotis — Leptopogon poecilotis P. L. Sclater, 1862
1.13.407. Bogota, Colombia. Sclater Collection. The second syntype is in the collection.
poecilurus — Empidochanes poecilurus P. L. Sclater, 1862
poensis — Alseonax poensis Boyd Alexander, 1903 3' Bull. Brit. Orn. Cl., 14 (Oct. 1903): 17. SYNTYPE, Adult male. Reg. no. 1911.12.23.1385. Near Bakaki, Fernando Po, 17 Nov. 1902. Collected and bequeathed by Boyd Alexander. This specimen is labelled as the "type male" and is also the specimens referred to as A. obscura Sjost in Ibis, (8) 3 (July 1903): 383. There are several other syntypes in the collection.
poensis — Amadina poensis Fraser, 1843
Reg. no. 1881.2.18.31. Clarence, Fernando Po. Collected by L. Fraser. Eyton Collection. The other syntypes are also in the collection.
poensis — Batis poensis Boyd Alexander, 1903
Bull. Brit. Orn. Cl., 13:34. SYNTYPE, Adult male. Reg. no. 1911.12.23.1460. (Bakaki, Fernando Po, West Africa, 14 Nov. 1902). Collected and bequeathed by Boyd Alexander. The paragraph giving locality and date were evidently omitted from the description through a printer's error, but are quoted in Ibis, (8) 3 (1903):386. The female syntype is also in the collection.
poensis — Calamocichla poensis Boyd Alexander, 1903 3: Bull. Brit. Orn. Cl., 13:37. HOLOTYPE, Adult (male, vide, Ibis,
(8) 3 (1903): 376). Reg. no. 1923.12.23.2844. Bilelipi, Fernando Po, 24 Nov. 1902. Collected and bequeathed by Boyd Alexander.

poensis — Callene poensis Boyd Alexander 1903
poensis — Cossypha poensis Strickland, 1844
poensis — Cyanomitra poensis Boyd Alexander, 1903
poensis — Dryoscopus poensis Boyd Alexander, 1903 23 Bull. Brit. Orn. Cl., 13:37. HOLOTYPE, Adult male. Reg. no. 1911.12.23.1998. Mount St. Isabel, Fernando Po, 25 Nov. 1902. Collected and bequeathed by Boyd Alexander.
poensis — Hylia poensis Boyd Alexander, 1903
poensis — Illadopsis cleaveri poensis Bannerman, 1934 32B Bull. Brit. Orn. Cl., 54:107 HOLOTYPE, Adult male. Reg. no. 1911.12.23.2371. Near Bakaki, Fernando Po, 19 Nov. 1902. Collected and bequeathed by Boyd Alexander.
poensis — Macrosphenus poensis Boyd Alexander, 1903 35 Bull. Brit. Orn. Cl., 13: 36. HOLOTYPE, Adult male. Reg. no. 1911.12.23.2123. Mount St. Ysabel, Fernando Po, 25 Nov. 1902. Collected and bequeathed by Boyd Alexander.
poensis — Phyllostrophus poensis Boyd Alexander, 1903 21 Bull. Brit. Orn. Cl., 13:35. HOLOTYPE, Adult male. Reg. no. 1911.12.23.1736. Bakaki, Fernando Po, 18 Nov. 1902. Collected and bequeathed by Boyd Alexander.
poensis — Psalidoprocne poensis Boyd Alexander, 1903 18 Bull. Brit. Orn. Cl., 13:34. SYNTYPE, Adult male. Reg. no. 1911.12.23.1313. Bakaki, Fernando Po, 12 Nov. 1902. Collected and bequeathed by Boyd Alexander. This is the author's selected male. There is another male syntype in the collection.
poensis — Stelgidillas poensis Boyd Alexander, 1903 21 Bull. Brit. Orn. Cl., 13:35. SYNTYPE, Adult male. Reg. no. 1911.12.23.1760. Mount Sipopo, Fernando Po, 8 Nov. 1902. Collected and bequeathed by Boyd Alexander. The female syntype is also in the collection.

- poensis Sycobrotus poensis Boyd Alexander, 1903 Bull. Brit. Orn. Cl., 13:38 SYNTYPE, Adult male. Reg. no. 1911.12.23.3642. Bakaki, 4.000', Fernando Po, 15 Nov. 1902. Collected and bequeathed by Boyd Alexander. There are two other syntypes in the collection. poensis — Turdus poensis Boyd Alexander, 1903 Bull. Brit. Orn. Cl., 13:37. SYNTYPE, Adult male. Reg. no. 1911.12.23.1784. Bakaki, Fernando Po. 19 Nov. 1902. and bequeathed by Boyd Alexander. There is a second male syntype in the collection. poensis — Zosterops stenocricota poensis Bannerman, 1915 Bull. Brit. Orn. Cl., 35:54. HOLOTYPE, Adult male. Reg. no. 1904.6.22.228. Banterbari, Fernando Po, 7 March 1904. Collected by E. Seimund. Presented by Mrs Sladen. poliocephalus — Emberizoides poliocephalus G. R. Gray, 1841 . . . 49 Zoology of the voyage of H.M.S. Beagle 3. Birds: 98. SYNTYPE. Adult. Reg. no. 1885.12.14.1325. Monte Video, Uraguay. Collected by C. Darwin. Salvin-Godman Collection, received from J. Gould in 1869. Listed as the type by Sharpe, Cat. Bds. Brit Mus., 12 (1888):759. There is also a syntype collected by Fitzrov in the collection. poliochlamys — Laniarius poliochlamys Gadow, 1883 Cat. Bds. Brit. Mus., 8:155, pl. 3. SYNTYPE, Adult. Reg. no. 1875.4.9.66. Fantee, Gold Coast. Collected by S. T. Aubinn. Sharpe Collection. The other two syntypes are also in the collection. . 32B poliogaster — Stachyris poilogaster Hume, 1880 Stray Feathers, 9:116. HOLOTYPE, Adult male. Reg. no. 1886.10.1.5959. At the foot of Gunong Pulai, Johore, Malaya, 14 March 1880. Collected by W. Davison. Hume Collection. poliogastra — Ochthoeca poliogastra Salvin and Godman, 1880 Ibis. (4) 4:123. SYNTYPE, Adult male. Reg. no. 1888.1.1.83. Sierra Nevada de Santa Maria, Colombia, 19 July 1879. Collected by F. A. A. Simons. Salvin-Godman Collection. The three other syntypes are also in the collection. poliogenys — Spermospiza poliogenys Ogilvie-Grant, 1906 Bull. Brit. Orn. Cl., 19:32. HOLOTYPE, Adult female. Reg. no. 1906.12.23.214. 20 miles north of Fort Beni, 3,000', Semliki Valley, Western Ruwenzori, 11 Aug. 1960. Collected by D. Carruthers. Presented by the subscribers to the Ruwenzori Expedition.
- poliogyna Brachypteryx poliogyna Ogilvie-Grant, 1895 . . . 31 Bull. Brit. Orn. Cl., 4:40. Syntype, Adult female. Reg. no. 1897.5.13.262. Lepanto District, north-west Luzon, Philippines

(Mount Data on label), 25 Jan. 1895. Collected by J. Whitehead and presented by the subscribers to his expedition. The male syntype is also in the collection. poliogyna — Loxia curvirostra poliogyna Whitaker, 1898 Ibis, (7) 4:625. SYNTYPE, Adult male. Reg. no. 1939.12.14.33. Bou-Chebka, 3,000', El-Oubira, West-central Tunisia, May 1897. Collected and presented by J. I. S. Whitaker. polionota — Lagonosticta polionota Shelley, 1873 . Ibis., (3)3:141. SYNTYPE, Adult male. Reg. no. 1889.7.20.338. Cape Coast Castle, Fantee, West Africa, 6 March 1872. Collected by G. E. Shelley. Shelley Collection. The other two syntypes are also in the collection. polionota — Merula polionota Sharpe, 1900 . in Seebohm, Monograph of the Turdidae, 2, pt.9:85, pl. 103. Syntype, Adult female. Reg. no. 1885.3.2.332. Roriama, 3,500', British Guiana, 7 Jan. 1884. Collected by H. Whitely. Salvin-Godman Collection. The male syntype is also in the collection. polionota — Ochthoeca polionota P. L. Sclater and Salvin, 1870 Proc. zool. Soc. Lond., 1869 (publ. 1870): 599. SYNTYPE, Adult. Reg. no. 1888.1.13.58. Pitumarca, South Peru, 8 April 1869. Collected by H. Whitely. Sclater Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888):21. There are two other syntypes in the collection. polionota — Spermophila polionota Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:118. SYNTYPE, Adult male. Reg. no. 1853.3.7.9. Brazil. Purchased of Verreaux. This is the described specimen (vide specimen label). There are two other syntypes in the collection. poliophrys — Alethe poliophrys Sharpe, 1902 Bull. Br. Orn. Cl., 13:10. HOLOTYPE, Adult female. Reg. no. 1902.12.8.14. Ruwenzori, Uganda, 12 Feb. 1902. Collected and presented by F. J. Jackson. poliopsa — Edoliisoma poliopsa Sharpe, 1882 J. Linn. Soc. Lond., 16:318, 434. SYNTYPE, Adult female. Reg. no. 1882.3.8.102. Astrolabe Mountains, south-east New Guinea. Collected by A. Goldie and purchased of E. Gerrard. polioptera — Campophaga polioptera Sharpe, 1878 Mitt. K. zool. Mus., Dresden, 1:370. SYNTYPE, Adult male. Reg. no. 1878.6.19.30. Saigon, Cochinchina, May 1867. Collected and presented by M. E. Pierre.

The two other syntypes are also in the collection. In Peters, Checklist of Birds of the World, 9 (1960): 193, the reference is given

as Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 59, 69, pl. 2.

Proc. zool. Soc. Lond., 1858: 179. HOLOTYPE, Adult female. Reg. no. 1858.3.10.49. Aru Islands. Collected by A. R. Wallace. Wallace Collection.
polysticta — Dendrornis polysticta Salvin and Godman, 1883 2 Ibis, (5) 1:210. Syntype, Adult female. Reg. no. 1889.5.14.644. Bartica Grove, British Guiana, 15 Oct. 1879. Collected by H. Whitely. Salvin-Godman Collection. There are several other syntypes in the collection.
pontifex—Dicoeum hypoleucum pontifex. See appendix, Corrections and Additions, vol. 2
 poonensis — Muscicapa poonensis Sykes, 1832
popayanus — Cassicus (Ocyalus) popayanus Waterhouse, 1841 . 57 Proc. zool. Soc. Lond., 1840 (publ. 1841): 183. Holotype, Adult male (relaxed mount). Reg. no. 1855.12.19.316. Popayan, Colombia (considered erroneous by Hellmayr and Conover, Catalogue of birds of the Americas, 10 (1937): 1). Collected for J. Gould. Purchased of the Zoological Society of London.
porphyrocephala — Iridornis porphyrocephala P. L. Sclater, 1857 . 51 Proc. zool. Soc. Lond., 1855 (publ. 1857): 227, pl. 110. SYNTYPE, Adult, Reg. no. 1857.10.16.1. Near Quito, Ecuador. Purchased of J. Gould. There is another syntype in the collection.
porphyrolaema — Nectarinia porphyrolaema Wallace, 1865
Proc. zool. Soc. Lond., 1865: 479. SYNTYPE, Adult male. Reg. no. 1873.5.12.864. Makassar, South Celebes. Collected by A. R. Wallace. Wallace Collection.
This specimen was listed as the type by Gadow, Cat. Bds. Brit. Mus., 9 (1884): 73. There are two other syntypes in the collection.
porphyronotus — Sturmus porphyronotus Sharpe, 1888 62 Ibis, (5) 6: 438. Syntype, Adult. Reg. no. 1875.9.2.42. Afghanistan (Yarkand, Central Asia, vide Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 38). Collected by E. Chapman. Sharpe Collection. There is one other syntype in the collection (vide Sharpe, Cat, Bds. Brit. Mus., 13 (1890): 38).
porphyrurus — Lamprocolius porphyrurus Hartlaub, 1874 62 Abhandl. nat. Ver. Bremen, 4 (1874): 60. SYNTYPE, Adult. Reg. no.
1874.5.1.130. Accra, West Africa. Collected by Haynes. Sharpe Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 180.

THO POWEIII
powelli — Pinarolestes powelli Salvin, 1879
praecognitus — Stachyris praecognitus Swinhoe, 1866 32H Ibis. (2) 2: 310. HOLOTYPE, Adult. Reg. no. 1898.10.3.766. Formosa, Dec. 1865. Collected for R. Swinhoe. Seebohm Bequest.
praedi — Malimbus rubricollis praedi Bannerman, 1921 61 Bull. Brit. Orn. Cl., 41:78. HOLOTYPE, Adult male. Reg. no. 1909.8.5.234. N'dalla Tando, northern Angola, 29 Oct. 1908. Collected by and purchased of W. J. Ansorge.
praepectoralis — Neocossyphus praepectoralis Jackson, 1906 31 Bull. Br. Orn. Cl., 16:90. HOLOTYPE, Adult male. Reg. no. 1906.6.3.5. Kibera, Toro, west Uganda, 14 Sept. 1905. Collected and presented by F. J. Jackson.
praetermissa — Alauda praetermissa Blanford, 1869
prateri — Sitta castanea prateri Whistler and Kinnear, 1932 43 J. Bombay nat. Hist. Soc., 35:522. SYNTYPE, Adult male. Reg. no. 1937.12.21.414. Anantagiri, 3,000', Vizagapatam, Madras, 5 Feb. 1930. Collected by V. S. La Personne. Presented by the Bombay Natural History Society. There are several other syntypes in the collection.
pratincola — Sturnella magna pratincola Chubb, 1921
princei — Chamaetylas princei Sharpe, 1873
princeps — Grallaria princeps P. L. Sclater and Salvin, 1869 4 Proc. zool. Soc. Lond., 1869: 418. SYNTYPE, Adult female. Reg. no. 1889.7.10.856. Calovevora, Veragua, 1869. Collected by E. Arcé. Salvin-Godman Collection. This is the 'second and more adult specimen' (loc. cit.) The other syntype is also the type of G. guatemalensis Sclater and Salvin, (q.v.).

pringlii — Dryoscopus pringlii Jackson, 1893
pritzbueri — Turdus pritzbueri Layard, 1878
problematicus — Copsychus problematicus Sharpe, 1876 31 Ibis, (3) 6:36. HOLOTYPE, Adult male. Reg. no. 1875.8.16.34. Sibu Island, Bruit River, Sarawak, Borneo. Collected by A. Everett and purchased of T. Higgins.
procas—Nesocichla eremita procax H. F. I. Elliott, 1954 31 Bull. Brit. Orn. Cl., 74:22. HOLOTYPE, Adult male. Reg. no. 1953.55.129. Nightingale Island, South Atlantic, April 11 1950. Collected and presented by H.F.I. Elliott.
propinqua — Chibia propinqua Tristram, 1889
propinqua — Uroloncha propinqua Sharpe, 1890
propinquus — Brachyurus propinquus Sharpe, 1877 12 Trans. Linn. Soc. Lond., (2) 1:330. SYNTYPE, Immature male. Reg. no 1876.8.28.38. Dumalon, Mindanao, Philippines. Collected by and purchased of J. B. Steere. The location of the figured adult specimen is not known but this immature specimen was referred to in the description.
proserpina — Nectarinia proserpina Wallace, 1863

prostheleucus — Scytalopus prostheleucus P. L. Sclater, 1857 28 Proc. zool. Soc. Lond., 1856 (publ. 1857) : 290. SYNTYPE, Adult.

Reg. no. 1857.7.30.18. Cordova, Vera Cruz, Mexico. Collected by and purchased of A. Sallé.
prosthemelas — Camarhynchus prosthemelas P. L. Sclater and Salvin, 1870
provincialis — Otocompsa jocosa provincialis Whistler, 1931, 21 Bull. Br. Orn. Cl., 52:40. HOLOTYPE, Adult male. Reg. no. 1886.9.1.2386. Kumaon Bhabar, Nepal, 24 March 1871. Hume Collection.
provocator — Ptilotis provocator Layard, 1875
pryeri — Dicaeum pryeri Sharpe, 1881
pryeri — Megalurus pryeri Seebohm, 1884
psittacea — Fringilla psittacea Gmelin, 1789
psittacula — Camarhynchus psittacula Gould, 1837 53 Proc. zool. Soc. Lond., 1837 : 6. SYNTYPE, Adult female (relaxed mount). Reg. no. 1855.12.19.22. Galapagos Archipelago, (James Island, vide, Zoology of the voyage of H.M.S. Beagle, 3, Birds, (1841) : 103). Collected by C. Darwin. Presented by the Zoological Society of London. There is another syntype in the collection.
7 71

- pulchellum Todirostrum pulchellum P. L. Sclater, 1874 Proc. zool. Soc. Lond., 1873 (publ. 1874): 781. HOLOTYPE, Adult (male, on label). Reg. no. 1888.1.13.247. Cosnipata, Cuzco, south-east Peru, (27 May 1861, on label). Collected by H. Whitely. Sclater Collection. pulchellus — Aegithaliscus pulchellus Rippon, 1900 40 Bull. Brit. Orn. Cl., 11:11. SYNTYPE, Adult. Reg. no. 1900.12. 20.283. Nanoi, 5.300', Loi Mai, South Shan States, 27 March 1899. Collected and presented by G. Rippon. There are three other syntypes in the collection. pulchellus — Vireolanius vulchellus P. L. Sclater and Salvin, 1859 . 56 Ibis. (1) 1:12. HOLOTYPE, Adult (relaxed mount). Reg. no. 1857. 10.16.7. Guatemala. Collected by G. U. Skinner. Purchased of J. Gould. pulcher — Eupetes pulcher Sharpe, 1882 . 32A J. Linn. Soc. Lond., 16: 319, 440. SYNTYPE, Adult male. Reg. no. 1882.3.8.58. Astrolabe Mountains, south-east New Guinea. lected by A. Goldie and purchased of E. Gerrard, junior. There are five other syntypes in the collection. pulcher — Myiobius pulcher P. L. Sclater, 1861 Proc. zool. Soc. Lond., 1860 (publ. 1861): 464. SYNTYPE, Adult. Reg. no. 1888.1.1.3 856. (Vicinity of Ouito, vide Sclater, Catalogue of a collection of American birds, (1862): 227), Ecuador. Collected for J. Gould. Sclater Collection. The second syntype is in the collection. pulcherrima — Dacnis pulcherrima P. L. Sclater, 1853 51 Revue Mag. Zool. 1853: 480. HOLOTYPE, Adult. Reg. no. 1884.7.31.105. Colombia. 1853. Collected for Argent. Sclater Collection. Described from a single specimen (vide, Proc. zool. Soc. Lond., (1855): 137-8), and since this specimen is also labelled "Nemosia" torquata De Bus" it is clearly the one described and referred to on p. 138 (op. cit.) pulcherrimus — Propasser pulcherrimus Moore, 1856 59 in Horsfield and Moore, Catalogue of the birds in the Museum of the Honourable East India Company, (21st Jan. 1856): 460. (the description in Proc. zool. Soc. Lond., 1855: 216, was not published
- until Feb. 1856). SYNTYPE, Adult male. Reg. no. 1880.1.1.4581. Nepal (Collected by B. H. Hodgson, on label: Strachy, in museum register.) Presented by the Honourable East India Company Museum.

Moore used a nomen nudum of Hodgson in J. E. Gray, Zooligocal Miscellany, (1844): 85. There are several other syntypes in the collection.

pulchra — Apalis pulchra Sharpe, 1891
pulchra — Haemophila pulchra P. L. Sclater, 1886 49 Ibis, (5) 4: 259, pl. 8. HOLOTYPE, Adult. Reg. no. 1886.8.1.23. Metucana, 7,000–8,000', Valley of the Rimac River, Peru. Collected for W. Nation. Sclater Collection.
pulchra — Macgregoria pulchra De Vis, 1897
pulchra — Porphyrospiza pulchra Sharpe, 1888 50 Cat. Bds. Brit. Mus., 12:625. SYNTYPE, Adult male. Reg. no. 1885.2.10.308. Furnas, Minas Geraes, Brazil, 15 June 1823. Collected by J. Natterer. Sclater Collection. The other two syntypes are in the collection.
pulchrala — Abrornis pulchrala Hodgson, 1844
pulchrius — Dicaeum pulchrius Sharpe, 1884
pulla — Stachyris poliocephala pulla Kloss, 1931
pumilus — Bradyornis pumilus Sharpe, 1895

punctata — Cinclidia punctata Gould, 1837
mount). Reg. no, 1855.12.19.391. Himalayas (restricted to Kalka, Baghat State, Simla Hill States, by Deignan, <i>Smiths. Misc. Coll.</i> , 107 (14) (1947): 5). Collected by J. Farell. Purchased from the Zoological Society Museum on its dispersal.
Although purchased as the type it was not listed by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 520, but may have been on display as a mounted specimen at that time.
puncticeps — Dysithamnus puncticeps Salvin, 1866
puncticeps — Picolaptes puncticeps P. L. Sclater and Salvin, 1873 . 2 Nomencl. Av. Neotrop.: 160. HOLOTYPE, Adult. Reg. no. 1889. 5.20.573. Interior of Guiana (Cayenne on label). Obtained through J. Verreaux. Sclater Collection.
puncticeps — Thanmophilus puncticeps P. L. Sclater, 1890 4 Cat. Bds. Brit. Mus., 15:212. SYNTYPE, Adult male. Reg. no. 1889.9.20.167. Tilotilo, Yungas, Bolivia. Collected by C. Buckley. Sclater Collection. There are three other syntypes in the collection.
puncticollis — Dendrocolaptes puncticollis P. L. Sclater and Salvin,
1868
punctulata — Calliste punctulata P. L. Sclater and Salvin, 1876 Proc. zool. Soc. Lond., 1876: 353, HOLOTYPE, Adult male. no. 1885.6.4.384. Tilotilo, Yungas Province, Bolivia, 1876. Collected by C. Buckley. Salvin-Godman Collection.
punctulatus — Chlorospingus punctulatus P. L. Sclater and Salvin,
1869
punensis — Grallaria punensis Chubb, 1918

Hodgson's paper and used Hodgson MS. names but there is no evidence that he saw the specimens which were listed as types by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 433, and which are therefore not regarded as valid. punjaubi — Alauda gulgula punjaubi Whistler, 1936	punicea — Fringilla punicea Horsfield, 1821
J. Bombay nat. Hist. Soc., 38:767. HOLOTYPE, Adult male. Reg. no. 1938.8.21.1. River Sutlej, Ferozepore, India, 28 April 1912. Collected and presented by H. Whistler. purpurascens — Calornis purpurascens G. R. Gray, 1870	in Hodgson, J. Asiat. Soc. Bengal, 13:953. Blyth completed Hodgson's paper and used Hodgson MS. names but there is no evidence that he saw the specimens which were listed as types by Sharpe, Cat. Bds. Brit. Mus., 12 (1888):433, and which are therefore not
Handlist of the genera and species of birds in the British Museum. 2:26 Gray refers to plate 33 in Gould, Supplement to the Birds of Australia. (1851). Since no single specimen of a number from Cape York which were available can be identified with the plate the latter is regarded as the type. Sharpe, Cat. Bds. Brit. Mus., 13 (1890):140, lists two females collected by McGillivray as types but these cannot be accepted. purpurascens — Chlorophanes purpurascens P. L. Sclater and Salvin, 1873	J. Bombay nat. Hist. Soc., 38:767. HOLOTYPE, Adult male. Reg. no. 1938.8.21.1. River Sutlej, Ferozepore, India, 28 April
Nomencl. Av. Neotrop.: 16, 157. Holotype, Adult. Reg. no. 1884.7.31.137. Caraccas, Venezuela, 1872. (Collected by O. Boucard, on label.) Sclater Collection. purpurascens — Sturnus purpurascens Gould, 1868	2:26 Gray refers to plate 33 in Gould, Supplement to the Birds of Australia. (1851). Since no single specimen of a number from Cape York which were available can be identified with the plate the latter is regarded as the type. Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 140, lists two females collected by McGillivray as types but
Proc. zool. Soc. Lond., 1868: 219. SYNTYPE, Adult. Reg. no. 1881.5.1.3223. Erzeroum, Asia Minor. (Collected by Dickson and Ross, on label.) Gould Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 38. Of the two syntypes this is the only one that can be traced. purpureiceps — Calornis purpureiceps Salvadori, 1878	Nomencl. Av. Neotrop.: 16, 157. HOLOTYPE, Adult. Reg. no. 1884.7.31.137. Caraccas, Venezuela, 1872. (Collected by O.
Atti. R. Accad. Sci. Torino, 13:535. SYNTYPE, Adult male. Reg. no. 1880.11.18.108. Admiralty Islands, Collected on the voyage of H.M.S. Challenger. Presented by the Treasury. There are three other syntypes in the collection. purus — Graucalus purus Sharpe, 1891	Proc. zool. Soc. Lond., 1868: 219. SYNTYPE, Adult. Reg. no. 1881.5.1.3223. Erzeroum, Asia Minor. (Collected by Dickson and Ross, on label.) Gould Collection. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 38.
Atti. R. Accad. Sci. Torino, 13:535. SYNTYPE, Adult male. Reg. no. 1880.11.18.108. Admiralty Islands, Collected on the voyage of H.M.S. Challenger. Presented by the Treasury. There are three other syntypes in the collection. purus — Graucalus purus Sharpe, 1891	purpureiceps — Calornis purpureiceps Salvadori, 1878 62
Ibis, (6) 3:121. SYNTYPE, Adult male. Reg. no. 1893.12.1.31. Mount Elgon, East Africa, 25 Feb. 1890. Collected and presented	Atti. R. Accad. Sci. Torino, 13:535. SYNTYPE, Adult male. Reg. no. 1880.11.18.108. Admiralty Islands, Collected on the voyage of H.M.S. Challenger. Presented by the Treasury.
This is the sole male syntype, selected by the author.	Ibis, (6) 3:121. SYNTYPE, Adult male. Reg. no. 1893.12.1.31. Mount Elgon, East Africa, 25 Feb. 1890. Collected and presented by F. J. Jackson.

pusilla — Pnoepyga pusilla Hodgson, 1845
pusillum — Eupsilostoma pusillum P. L. Sclater, 1860
pusillus — Cnipolegus pusillus P. L. Sclater and Salvin, 1873
pusillus — Corvus pusillus Tweeddale, 1878
pusillus — Cyphorhinus pusillus P. L. Sclater, 1859
pusillus — Drymocataphus pusillus Delacour, 1927 321 Bull. Br. Orn. Cl., 47: 161. HOLOTYPE, Adult male. Reg. no. 1927.6.5.22. Tam Dao, central Tonkin, 12 Dec. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China. See Delacour, Bull. Brit. Orn. Cl., 48 (1928): 136. In the original reference the location of the specimen was incorrectly given as the Paris Museum.
pusillus — Xiphorhynchus pusillus P. L. Sclater, 1860
pycnopygius — Sphenaeacus pycnopygius P. L. Sclater, 1852 3: in Jardine, Contributions to Ornithology: 148. HOLOTYPE, Adult (relaxed mount). Reg. no. 1852.5.1.35. Damaraland, South-west Africa. Collected by C. J. Andersson. Purchased of Bartlett.

- pygmaeus Zosterornis pygmaeus Ogilvie-Grant, 1896 . . . 32B Bull. Brit. Orn. Cl., 6: 18. Syntype, Adult male. Reg. no. 1897. 12.11.17. Barugo, north Leite, Philippines, 5 Aug, 1896. Collected by J. Whitehead and presented by the subscribers to the Whitehead Expedition.

This is the author's selected male. His selected female syntype is

also in the collection.

- pyrrhonota Hirundo pyrrhonota Vigors and Horsfield, 1827 . . . 18

 Trans. Linn. Soc. Lond., 15:190. HOLOTYPE, Adult. Reg. no. 1863.7.7.9. New South Wales (error, western Gulf of Carpentaria, 9 Dec. 1802, vide R. Brown's MS in the museum and Kinnear, Proc. Linn. Soc. Lond., 1932:36). Collected by R. Brown on the voyage of H.M.S. Investigator and presented by the Linnean Society. Vigors & Horsfield used Latham's MS. name.
- pyrrhophrys Euphonia pyrrhophrys P. L. Sclater, 1851. . . . 51 in Jardine, Contributions to Ornithology: 89, pl. 75. SYNTYPE, Adult (female, vide Sclater, Catalogue of a collection of American birds, (1862): 55). Reg. no. 1885.6.12.25. Bogota, Colombia, 1847. (Collected by Williams, on label.) Sclater Collection. Listed as the type by Sclater (op. cit. 1862).
- pyrrhops Euscarthmus pyrrhops Cabanis, 1874.

 J. Orn. 1874: 98. The original description was based on a female in the Berlin Museum, vide Taczanowski, Ornithologie du Perou, 2 (1874): 232, and the male specimen listed as "authentic" by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 83, has no type status.

** *
pyrrhops — Stachyris pyrrhops Blyth, 1844
pyrrhoptera — Certhia pyrrhoptera Latham, 1801
pyrrhopygia — Acanthiza pyrrhopygia Vigors and Horsfield, 1827 . 36 Trans. Linn. Soc. Lond., 15: 227. HOLOTYPE, Adult female (relaxed mount). Reg. no. 1863.7.7.28. Australia. Presented as the type by the Linnean Society of London.
pyrrhoura — Myzornis pyrrhoura Blyth, 1843
pyrrhulinus — Schoenichus pyrrhulinus Swinhoe, 1876
quadricolor—Prionochilus quadricolor Tweeddale, 1878 45 Proc. zool. Soc. Lond., 1877 (publ. 1878): 762, pl. 77. HOLOTYPE, Adult. Reg, no. 1888.7.12.133. Cebu, Philippines, April 1877. Collected by A. H. Everett. Tweeddale Collection.
quelimanensis—Seicercus ruficapilla quelimanensis Vincent, 1933 . 35 Bull. Brit. Orn. Cl., 53:136. Holotype, Adult male. Reg. no. 1933.3.1.14. Namuli Mountains, 5,600', Quelimane Province, Portuguese East Africa, 25 July 1932. Collected by and purchased of J. Vincent.
quercinus—Vireolanus melitophrys quercinus Griscom, 1935 56

Ibis, (13) 5:551. HOLOTYPE, Adult. Reg. no. 1859.3.10.200. (mistakenly numbered 290). Oak Forest, Volcan de Fuego, Guate-

mala, 18 Sept. 1859. Collected by O. Salvin. Salvin-Godman Collection.
questi—Nesospiza acunhae questi P. R. Lowe, 1923 4 Ibis, (11) 4: 520. HOLOTYPE, Adult male. Reg. no. 1922.12.6.125. Nightingale Island, South Atlantic, 21 May 1922. Collected by G. H. Wilkins on the Shackleton-Rowett Expedition of the Quest. Presented by J. Q. Rowett.
quindiuna—Cyanocitta armillata var. quindiuna P. L. Sclater and and Salvin, 1876
quinquestriata—Zonotrichia quinquestriata P. L. Sclater and Salvin,
Proc. zool. Soc. Lond., 1868: 323. HOLOTYPE, Adult. Reg. no. 1885.2.10.440. Mexico. Collected by Floresi (vide Salvin and Godman, Biol. CentrAmer., Aves, 1 (1886): 368. Sclater Collection, received from Gould.
quiscalina—Campephaga quiscalina Finsch, 1869 2
Ibis, (2) 5: 189. HOLOTYPE, Adult (in bad condition). Reg. no. 1875.4.9.78. Fantee, Gold Coast. Collected for E. T. Higgins. Sharpe Collection.
radcliffei—Actinodura radcliffei Harington, 1910
radcliffei—Leioptila melanoleuca radcliffei E. C. S. Baker, 1922 . 321 The Fauna of British India (2nd ed.); Birds, 1:300. SYNTYPE, Reg. no. 1910.10.7.3. (Kyatpyin, Mogôk Road, Ruby Mines, 24 March 1910, on label), north-east central Burma. Collected by H. D. Radcliffe and presented by H. H. Harington. There is a second syntype in the collection.
radiolatus—Dendrocolaptes radiolatus P. L. Sclater, 1867
Proc. zool. Soc. Lond., 1867: 755. HOLOTYPE, Adult. Reg. no. 1889.5.20.653. Yurimaguas, Huallaga River, 60 miles from Amazon River. Collected by E. Bartlett and purchased with the Sclater Collection.
rafflesi—Prinia rafflesi Tweeddale, 1877
Tamesi—Prima rafflesi Tweeddale, 1877

raisulii—Certhia brachydactyla raisulii Bannerman, 1926
rajputanae—Salpornis spilonotus rajputanae R. Meinertzhagen, 1926 44A Bull. Brit. Orn. Cl., 46:83. HOLOTYPE, Adult male. Reg. no. 1886.11.1.1742. Sambhur, Rajputana, western India, 2 June 1873. Collected by R. M. Adam. Hume Collection.
rama—Sylvia rama Sykes, 1832
ramsayi—Actinura ramsayi Walden, 1875
ramsayi—Myzomela ramsayi Finsch, 1886
ramsayi—Petroica ramsayi Sharpe, 1879
ramsayi—Trochalopterum ramsayi Ogilvie-Grant, 1904 32E Bull. Brit. Orn. Cl., 14:92. HOLOTYPE, Adult male. Reg. no. 1888.4.20.581. Karen-nee, Salween District, Burma, 1 April 1874. Collected by R. G. Wardlaw Ramsay. Tweeddale Collection.
raymondi—Onychognathus tenuirostris raymondi R. Meinertzhagen, 1936
razae—Spizocorys razae Boyd Alexander, 1898

2932. Raza Island, Cape Verde Islands, May 1897. Collected and bequeathed by Boyd Alexander. This is the author's selected male. His selected female and fifteen other syntypes are also in the collection. rectirostris—Limnornis rectirostris Gould, 1839 . Zoology of the Voyage of H.M.S. Beagle, 3, Birds: 80, pl. 26. Syntype, Adult (relaxed mount). Reg. no. 1855.12.19.77. Maldonado, Uruguay. Collected by Charles Darwin and purchased of the Zoological Society of London. Listed as type by P. L. Sclater, Cat. Bds. Brit. Mus., 15 (1890): 77. There is another syntype in the collection. regia—Erythrospiza regia P. L. Sclater, 1881 60 *Ibis*, (5) 4: 544. HOLOTYPE, Adult female. Reg. no. 1891.1.28.2. Api, New Hebrides, August 1880. Collected by K. Howard. Presented by P. L. Sclater. reguloides—Acanthiza reguloides Vigors and Horsfield, 1827. . . . 36 Trans. Linn. Soc. Lond., 15:226. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.26. Australia. (Paramatta, New South Wales, vide Gould, Birds of Australia, 3 (1848): pl. 62). Collected by G. Caley. Presented as the type by the Linnean Society of London. reguloides—Phyllopneuste reguloides Blyth, 1842. J. Asiat. Soc. Bengal, 11:191. SYNTYPE, Adult (relaxed mount). Reg. no. 1844.3.4.4. (Calcutta, on label.) Presented by E. Blyth. regulus—Grallaria regulus P. L. Sclater, 1860 Proc. zool. Soc. Lond., 1860: 66. HOLOTYPE, Adult. Reg. no. 1889.9.20.609. Pallatanga, west Ecuador, 1858. Collected by L. Fraser. Sclater Collection. reichenowi—Cinnyris reichenowi Sharpe, 1891 Ibis, (6) 3:444, HOLOTYPE, Adult male. Reg. no. 1893.12.1.13. Sotik, Kenya, 3 Oct. 1889. Collected and presented by F. J. Jackson. reichenowi—Trochocercus reichenowi Sharpe, 1904 38 Ibis, (8) 4:630. SYNTYPE, Adult. Reg. no. 1875.4.9.333. Fantee, West Africa. Collected by H. T. Ussher. Sharpe Collection. There are at least two other syntypes in the collection. reidi-Fringillaria reidi Shelley, 1902 . Birds of Africa, 3:158. SYNTYPE, Adult male. Reg. no. 1892.5.6. 178. Ingagane River, Natal, 14 July 1881. Collected by and purchased of S. G. Reid. The female syntype is also in the collection. remota—Dendroica graciae remota Griscom, 1935 *Ibis*, (13) 5:548. HOLOTYPE, Adult male. Reg. no. 1892.4.20.900.

Volcan Viego, over 4,000', Chinandega, Nicaragua, 16 April 1891.

Collected by W. B. Richardson, Salvin-Godman Collection.

- remotum—Piprisoma modestum remotum Robinson and Kloss, 1915 45 J.F.M.S. Mus., 6: 29. Holotype, Adult male. Reg. no. 1927.4. 18.1011. Bukit Tanngga, 1,500', Negri Sembilan, Federated Malay States, 27 Jan. 1913. Collected by a native collector for the Federated Malay States Museum. Presented by the Raffles Museum, Singapore.
- repetens—Vireo solitarius repetens van Rossem, 1939 50
 Proc. Biol. Soc. Wash., 52 (1939):11. Holotype, Adult male.
 Reg. no. 1891.11.3.357. Tixtla, 6,000', (Amula) Guerrero, Mexico,
 Aug. 1888. Collected by D. W. Smith. Salvin-Godman Collection.
- restricta—Geothlypis restricta Maynard, 1887 5.

 American Exchange and Mart. Jan. 15, 1887 : 33. HOLOTYPE, Adult male. Reg. no. 1906.12.7.3409. Nassau, Bahamas, 2 Feb. 1884. Collected by C. J. Maynard. Purchased of G. S. Miller. Description repeated by Palmer, Auk, 17 (1900) : 225.
- restricta—Tchitrea viridis restricta Salomonsen, 1933 38

 Bull. Brit. Orn. Cl., 54:48. HOLOTYPE, Adult male. Reg. no.
 1929.1.7.22. Nkose Island, 3,700', Victoria Nyanza, 26 May 1928.
 Collected and presented by C. R. S. Pitman.

- reynoldsi—Notiospar curaeus reynoldsi W. L. Sclater, 1939 . . . 57 Ibis, (14) 3:143. HOLOTYPE, Adult male. Reg. no. 1928.7.21.12. Viamonte, Tierra del Fuego, 10 Aug. 1927. Collected and presented by P. W. Reynolds.

•
rhizophorae—Arachnechthra rhizophorae Swinhoe, 1869 46 Ann. Mag. nat. Hist., (4) 4: 436. SYNTYPE, Adult male. Reg. no. 1895.5.1.2381. Hainan, Southern China, (Feb. 1868, on label). Collected by R. Swinhoe. Shelley Collection. This specimen was labelled as the type by the author. There are other syntypes in the collection.
rhodesiae—Anthoscopus ansorgei rhodesiae W. L. Sclater, 1932 . 41 Bull. Brit. Orn. Cl., 52:143. HOLOTYPE, Adult male. Reg. no. 1932.6.5.3. Mt. Sunzu, 6,300', near Abercorn, Northern Rhodesia. 18 Oct. 1931. Collected and presented by Admiral H. Lynes.
rhodesiae—Cinnyris oustaleti rhodesiae Benson, 1955
rhodesiae—Pinaornis rhodesiae Chubb, 1908 31 Bull. Br. Orn. Cl., 21:110. HOLOTYPE, Adult. Reg. no. 1908.11. 6.2. Manzinyama, Gambo Kraal, Rhodesia, 17 March 1908. Collected by F. P. Mennel and E. C. Chubb and presented by the Bulawayo Museum.
rhodesiae—Quelea cardinalis rhodesiae C. H. B. Grant and Mackworth-Praed, 1944
rhodinolaema—Phoenicothraupis rhodinolaema Salvin and Godman, 1883
rhodolaema—Anthreptes rhodolaema Shelley, 1877
rhodometopus—Propasser rhodometopus Biddulph, 1881 59 Ibis, (9) 5: 156, pl. 6. Syntype, Adult male. Reg. no. 1897.12.10. 79. Kashgar, Yarkand, Jan. 1874. Collected and presented by J. Biddulph. This specimen is labelled "Type and figured" but the date and locality given in the description are from another specimen, reg. no. 1897.12.10.78., now missing from the collection. A female syntype is also in the collection.

- richardi—Cercomela familiaris richardi Macdonald, 1953 . . . 31

 Ibis, 95:73. HOLOTYPE, Adult male. Reg. no. 1950.50.373.

 Springbok, 2,600′, Little Namaqualand, 8 Dec. 1949. Collected by J. D. Macdonald on the British Museum South-West Africa Expedition.
- richardsoni—Chlorospingus opthalmicus richardsoni Griscom, 1935. 51

 Ibis, (13) 5:816. HOLOTYPE, Adult male. Reg. no. 1898.12.14.

 568. Sierra de las Minas, Guatemala, 11 July 1897. Collected by W. B. Richardson. Salvin-Godman Collection.
- richardsoni—Icterus graduacauda richardsoni W. L. Sclater, 1939 . 57 Ibis, (14) 3:141. HOLOTYPE, Adult male. Reg. no. 1890.12.14. 2085. Chimalapa, 9,000′, 41 miles north-east of Tehuantepec, Mexico, 28 March 1890. Collected by W. B. Richardson. Salvin-Godman Collection.
- richardsoni—Saltator grandis richardsoni van Rossem, 1938 . . . 50

 Bull. Brit. Orn. Cl., 58: 124. HOLOTYPE, Adult female. Reg. no.
 1894.7.1.1180. Plains of Colima, Mexico, 19 Jan. 1890. Collected
 by W. B. Richardson. Salvin-Godman Collection.
- richardsoni—Spermophila richardsoni Salvin and Godman, 1891 . 59

 Ibis, (6) 3:611. Syntype, Adult male. Reg. no. 1899.2.1.425.

 San Benito, Chiapas State, Mexico, 1 June 1897. Collected by W. B. Richardson. Salvin-Godman Collection.

 Listed as the type by Hellmayr, Catalogue of birds of the Americas, 11 (1938): 214. There are several other syntypes in the collection.
- ricketti—Paradoxornis webbiana ricketti Rothschild, 1922 . . . 33 Bull. Br. Orn. Cl., 43:11. HOLOTYPE, Adult female. Reg. no. 1922.12.7.359. Yangtze Valley, 6,000', north-west Yunnan, 23 Oct. 1921. Collected by G. Forrest and presented by Stephenson Clarke.

Proc. zool. Soc. Lond., 1887: 50. HOLOTYPE, Adult. Reg. no. 1888.1.13.944. Bogota, Colombia, Sclater Collection. Purchased by Sclater from Stevens in 1855 (on label). See also Van Rossem, Trans. San Diego Soc. Nat. Hist., 7 (1934): 350. ridleyana—Elainea ridleyana Sharpe, 1888	Bull. Brit. Orn. Cl., 14:92. SYNTYPE, Adult male. Reg. no. 1903.7.3.641. (Ah Ch'ung, on label), Fohkien Province, south China, 8 Dec. 1899. Collected and presented by C. B. Rickett.
no. 1888.6.1.1a. Fernando Norohna (= Noronha) Island, off Brazilian Coast, (30 Aug. 1887, on label). Collected by H. N. Ridley. Presented by the Royal Society. This specimen was labelled "type". There are two other syntypes in the collection. riisii—Elainea riisii P. L. Sclater, 1860	Proc. zool. Soc. Lond., 1887: 50. HOLOTYPE, Adult. Reg. no. 1888.1.13.944. Bogota, Colombia, Sclater Collection. Purchased by Sclater from Stevens in 1855 (on label). See also Van Rossem, Trans. San Diego Soc. Nat. Hist., 7 (1934):
Riise. Sclater Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 141. There are two other syntypes in the collection. rileyi—Stachyris nigriceps rileyi Chasen, 1936	no. 1888.6.1.1a. Fernando Norohna (= Noronha) Island, off Brazilian Coast, (30 Aug. 1887, on label). Collected by H. N. Ridley. Presented by the Royal Society. This specimen was labelled "type". There are two other syntypes in the collection.
Bull. Br. Orn. Cl., 56: 115. New name for S. nigriceps dilutus Robinson and Kloss, 1919 (q.v.), on the grounds of its preoccupation by S. poliocephala diluta Robinson and Kloss, 1918. ripponi—Abrornis ripponi Sharpe, 1902	Riise. Sclater Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 141.
Bull. Brit. Orn. Cl., 13:11. SYNTYPE, Adult. Reg. no. 1903.8.8. 233. Gyi-dzin-shan, 8,000', east of Tali-fu, West Yunnan, 3 April 1902. Collected and presented by G. Rippon. There are two other syntypes in the collection. ripponi—Actinodura ripponi Ogilvie-Grant and La Touche, 1907. 32B Ibis, (9) 1:186. SYNTYPE, Adult. Reg. no. 1905.9.10.286. Mount Victoria, (8,000', on label), South Chin Hills, 25 March 1904. Collected and presented by Col. G. Rippon. There are several other syntypes in the collection. ripponi—Certhia familiaris ripponi Kinnear, 1929	Robinson and Kloss, 1919 $(q.v.)$, on the grounds of its preoccupation
Ibis, (9) 1:186. SYNTYPE, Adult. Reg. no. 1905.9.10.286. Mount Victoria, (8,000', on label), South Chin Hills, 25 March 1904. Collected and presented by Col. G. Rippon. There are several other syntypes in the collection. ripponi—Certhia familiaris ripponi Kinnear, 1929	Bull. Brit. Orn. Cl., 13:11. SYNTYPE, Adult. Reg. no. 1903.8.8. 233. Gyi-dzin-shan, 8,000', east of Tali-fu, West Yunnan, 3 April 1902. Collected and presented by G. Rippon.
Bull. Brit. Orn. Cl., 49: 109. New name for C. himalayana intermedia Kinnear, 1921 (q.v.), on the grounds of preoccupation by C. intermedia Müller, 1776. ripponi—Pericrocotus solaris ripponi E. C. S. Baker, 1924 20 The Fauna of British India, (2nd ed.) 2: 327. HOLOTYPE, Adult male.	Ibis, (9) 1:186. SYNTYPE, Adult. Reg. no. 1905.9.10.286. Mount Victoria, (8,000', on label), South Chin Hills, 25 March 1904. Collected and presented by Col. G. Rippon.
The Fauna of British India, (2nd ed.) 2:327. HOLOTYPE, Adult male.	Bull. Brit. Orn. Cl., 49: 109. New name for C. himalayana intermedia Kinnear, 1921 (q.v.), on the grounds of preoccupation by C.
	The Fauna of British India, (2nd ed.) 2:327. HOLOTYPE, Adult male.

See B. P. Hall, Bull. Br. Orn. Cl., 73 (1953): 63–4 for a note on the holotype.
ripponi—Pomatorhinus ripponi Harington, 1910
ripponi—Proparus ripponi Harington, 1913
ripponi—Propasser ripponi Sharpe, 1902
ripponi—Suthora ripponi Sharpe, 1905
ripponi—Trochalopterum ripponi Oates, 1900
roberti—Callene roberti Boyd Alexander, 1903 3: Bull. Br. Orn. Cl., 13: 37. Holotype, Adult female. Reg. no. 1911.12.23.1874. Near Bakaki, 4,000′, Fernando Po, 17 Nov. 1902. Collected and bequeathed by Boyd Alexander.
roberti—Pnoepyga roberti Godwin-Austen and Walden, 1875 321 Ibis, (3) 5:252. SYNTYPE, Adult. Reg. no. 1895.7.14.1248. Chakha, Munipur Hills, Collected by W. Robert. Godwin-Austen Collection.

States, 15 April 1898. Collected and presented by Col. G. Rippon.

Godwin-Austen Collection.

The other syntype, from Konchungbum, is also in the collection, and was listed as type by Sharpe, *Cat. Bds. Brit. Mus.*, 6 (1881): 284. Only Hemes Peak was specified in the original description.

roberti—Sphenocichla roberti Godwin-Austen and Walden, 1875 . 32B Ibis, (3) 5:251. Syntype, Adult. Reg. no. 1895.7.14.3400. Hemes Peak, North Cachar Hills. Collected by H. Robert.

- robertsi—Certhilauda albofasciata robertsi Macdonald, 1953. . 17

 Proc. zool. Soc. Lond., 1953: 1003, pl. 2. HOLOTYPE, Adult male.

 Reg. no. 1928.10.25.65. Fountains, near Pretoria, Transvaal, 18

 Dec. 1926. Collected and presented by H. Lynes.
- robini—Garrulax pectoralis robini Delacour, 1927. 32B Bull. Br. Orn. Cl., 47:157. HOLOTYPE, Adult male. Reg. no. 1927.6.5.9. Tam Dao, central Tonkin, 2 Dec. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China.

- robinsoni—Enicurus maculatus robinsoni E. C. S. Baker, 1922 . . . 31 Bull. Br. Orn. Cl., 43: 19, 84. HOLOTYPE, Adult male. Reg. no. 1919.12.20.356. Langbian Peaks, 5,500-5,600', south Annam, 23 April 1918 (on label; May in text). Collected and presented by H. C. Robinson and C. B. Kloss.
- robinsoni—Myiophoneus robinsoni Ogilvie-Grant, 1905. . . . 31 Bull. Br. Orn. Cl., 15:69. HOLOTYPE, Adult male. Reg. no. 1905.4.4.2. Gunung Menkuanghebah, 4,700', Selangore, Malaya, 1 Feb. 1905. Collected and presented by H. C. Robinson.
- robinsoni—Oriolus traillii robinsoni Delacour, 1927 63

 Bull. Brit. Orn. Cl., 47: 168. Syntype, Adult female. Reg. no. 1927.6.5.21. Djiring, South Annam, 1 March 1927. Collected by J. Delacour, P. Jabouille, and W. P. Lowe, on their 3rd Expedition to French Indo-China.

There were thirty-one syntypes. This specimen was the author's selected female and his selected male syntype is also in the collection.

robinsoni—Pteruthius flaviscapis robinsoni Chasen and Kloss, 1931. 32B Bull. Raffles Mus., 5:86. HOLOTYPE, Adult male. Reg. no. 1947.60.70. (Kamborangah, on label), 7,200', Mount Kinabalu, North Borneo, 29 March 1929. Collected by F. N. Chasen and presented by the Raffles Museum, Singapore.

robinsoni—Pycnonotus robinsoni Ogilvie-Grant, 1905 21 Fasciculi Malayenses; Results of Expedition to Perak and Siamese Malay States: 85. Syntype, Adult male. Reg. no. 1905.2.1.378. Tanjong, Patani, Malay Peninsula, 1 Oct. 1901. Collected by H. C. Robinson and presented by the Royal Society and the Universities of Edinburgh and Liverpool. The female syntype is also in the collection.
robustipes—Horeites robustipes Swinhoe, 1866
roraimae—Atticora fucata roraimae Chubb, 1920
roraimae—Basileuterus roraimae Sharpe, 1885
roraimae—Chlorophonia roraimae Salvin and Godman, 1884 . 51 Ibis, (5) 2:444. SYNTYPE, Adult male. Reg. no. 1885.6.4.32. Roraima, 6.000', British Guiana, 6 Sept. 1883. Collected by H. Whitely. Salvin-Godman Collection. There are six other syntypes in the collection.
roraimae—Grallaria regulus roraimae Chubb, 1921
roraimae—Myiobius roraimae Salvin and Godman, 1883 9 Ibis, (5) 1:207. SYNTYPE, Adult male. Reg. no. 1888.1.1.1239. Roraima, British Guiana, 21 Nov. 1881. Collected by H. Whitely. Salvin-Godman Collection. There is another syntype in the collection.
roraimae—Pipromorpha oleaginea roraimae Chubb, 1919 9 Ann. Mag. nat. Hist. (9) 4:303. HOLOTYPE, Adult male. Reg. no. 1888.1.1.531. Mount Roraima, 3,500–5,000', British Guiana, 20 Dec. 1881. Collected by H. Whitely Jnr. Salvin-Godman Collection.
roraimae—Pseudochloris roraimae Chubb, 1921

1885.12.14.1262. Mount Roraima, 3,500', British Guiana, 20 Nov. 1883. Collected by H. Whitely. Salvin-Godman Collection. The described female syntype is also in the collection.
roraimae—Sporophila gutteralis roraimae Chubb, 1921 59 Ann. Mag. Nat. Hist., (9) 7:193. HOLOTYPE, Adult male. Reg. no. 1885.12.14.494. Mount Roraima, 3,500', British Guiana, 22 June 1883. Collected by H. Whitely. Salvin-Godman Collection.
roraimae—Turdus roraimae Salvin and Godman, 1884 31 Ibis, (5) 2:443. Syntype, Adult male. Reg. no. 1885.3.2.339. Roraima, 6,000', British Guiana, 22 Aug. 1883. Collected by H. Whitely. Salvin-Godman Collection. There are four other syntypes in the collection.
roraimae—Vireo roraimae Chubb, 1921
rosacea—Pyrrhula rosacea Seebohm, 1882
rosaecolor—Pyrrhulinota roseacolor Hodgson, 1844
roseaceus—Anthus roseaceus Horsfield and Moore, 1854 19 Catalogue of Birds in the Museum of the Honourable East India Company, 1:357. SYNTYPE, Adult. Reg. no. 1880.1.1.1176. Nepal. Collected by B. H. Hodgson and presented by the Honourable East India Company Museum. The authors used Hodgson's MS name. The five specimens listed as types by Sharpe, Cat. Bds. Brit. Mus., 10 (1885):591, were presented by Hodgson and are therefore not accepted.
roseipectus—Carpodacus roseipectus Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:424. Syntype, Adult male. Reg. no.
1885.12.14.1146. Oaxaca, Southern Mexico. Collected by A. Fenochio. Salvin-Godman Collection. This is the described specimen (vide specimen label), and was listed as the type by Hellmayr, Catalogue of birds of the Americas, 11 (1938): 158. There are two other syntypes in the collection.
roseires—Cisticola erythrops roseires W. L. Sclater and Mackworth-Praed, 1918

rostrata—Mellopitta lugubris rostrata Ogilvie-Grant, 1913 32A Bull. Br. Orn. Cl., 31:104. Holotype, Adult male. Reg. no. 1911.12.20.1068. Utakwa River, Camp 6b, 4,200′, Dutch New Guinea, 26 Jan. 1913. Collected by C. B. Kloss and presented by the Committee of the British Ornithologists' Union New Guinea Expedition.	
rostrata—Oreocincla rostrata Hodgson, 1845	1
rostrata—Ptilotis rostrata Wallace, 1865	8
rostratus—Megalophonus rostratus Hartlaub, 1863 17 in Gurney, Ibis, (1) 5:327, pl. 9. Syntype, Adult male. Reg. no. 1864.10.18.1. Natal. Collected by T. Ayres and presented by J. H. Gurney. Listed as "typical example" by Hartlaub in Decken's Reisen in Ost Afrika, 4 (1870):464. There is a female syntype in the collection.	7
	9
rothschildi—Cochoa rothschildi E. C. S. Baker, 1924	
rothschildi—Diphyllodes rothschildi Ogilvie-Grant, 1915 70 Ibis, (10) 2, Jubilee Supplement: 24. HOLOTYPE, Adult male. Reg. no. 1881.5.1.1737. Salawatti Island, New Guinea, Feb. 1870. Gould Collection.	
rothschildi—Hemichelidon sibirica rothschildi E. C. S. Baker, 1923 . 3' Bull. Brit. Orn. Cl., 43:156. Holotype, Adult male. Reg. no. 1921.7.15.463. Lichiang Range, North-west Yunnan, (10,000–12,000', Lat. 27°20' N., July 1918, on label). Collected by G. Forrest. Presented by Col. Stephenson Clarke.	7

rothschildi—Serinus rothschildi Ogilvie-Grant, 1902 59 Bull. Brit. Orn. Cl., 13:21. SYNTYPE, Adult male. Reg. no. 1902.12.13.42. Ichaf Ravine, Hameirah, Yemen Frontier of Southern Arabia, 8 August. Collected by and purchased of G. W. Bury. The female syntype is also in the collection.
rotumae—Lalage nigra rotumae Neumann, 1927
rovumae—Parus rovumae Shelley, 1892
rowei—Geokichla piaggiae rowei C. H. B. Grant and Mackworth-Praed, 1937
rubecula—Geocichla rubecula Gould, 1836
rubecula—Poospiza rubecula Salvin, 1895
rubecula—Todus rubecula Latham, 1801
rubeculoides—Accentor rubeculoides Moore, 1854

rubeculoides—Myiagra rubeculoides Vigors and Horsfield, 1827 . 3 Trans. Linn. Soc. Lond., 15: 253. HOLOTYPE, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 374), (relaxed mount). Reg. no. 1863.7.7.46. Australia. Presented by the Linnean Society as the type.	8
Proc. zool. Soc. Lond., 1845: 36. SYNTYPE, Adult male. Reg. no. 1845.1.9.304. (Icon. ined. no. 831.) Nepal. Collected and presented by B. H. Hodgson. There are other syntypes in the collection.	59
rubeculoides—Saxicola rubeculoides Sykes, 1832	37
rubescens—Locustella rubescens Blyth 1845	35
rubescens—Sylvia rubescens Blanford, 1874	35
rubeus—Otocorys alpestris rubeus Henshaw, 1884	17
rubicunda—Cassinia rubicunda Hartlaub, 1860	31
rubicunda—Neopipo rubicunda P. L. Sclater and Salvin, 1869. Proc. zool. Soc. Lond., 1869: 438, pl. 30. SYNTYPE, Adult male. Reg. no. 1870.6.14.17. Chamicuros, eastern Peru, 18 Aug. 1867. Collected by and purchased of E. Bartlett. The two other syntypes are also in the collection.	8

- rubiginosa—Ortygocichla rubiginosa P. L. Sclater, 1881 . . . 35 Proc. zool. Soc. Lond., 1881: 452, pl. 39. HOLOTYPE, Adult female. Reg. no. 1882.7.5.14. Blanche Bay District, New Britain, 26 Nov. 1880. Collected by T. Kleinschmidt. Purchased of the Museum Godeffroy, Hamburg.
- rubiginosus—Anabates rubiginosus P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1856 (publ. 1857): 288. HOLOTYPE, Adult male. Reg. no. 1889.5.20.328. Cordova, Vera Cruz, southern Mexico. Collected by A. Sallé. Sclater Collection.
- rubriceps—Pyranga rubriceps G. R. Gray and Mitchell, 1844. . . 51
 The Genera of birds, 2:364, pl. 89. HOLOTYPE, Adult (later identified as female) (relaxed mount). Reg. no. 1843.2.9.12. (Bogota, vide British Museum register.) Purchased of Mr Turner.

 This specimen is known, from its date of accession, to have been available to the author, and resembles the plate exactly.
- rubrifrons—Hylophilus rubrifrons P. L. Sclater and Salvin, 1867. 56

 Proc. zool. Soc. Lond., 1867: 569, pl. 30. HOLOTYPE, Adult.

 Reg. no. 1886.9.15.316. River Amazon (= Para, vide Hellmayr,

 Catalogue of Birds of the Americas, 8 (1935): 182). Collected by

 A. R. Wallace. Sclater Collection.
- rubrifrons—Pholidornis rubrifrons Sharpe and Ussher, 1872 60
 Ibis, (3) 2:182. Syntype, Adult male. Reg. no. 1874.5.1.54.
 Denkara, Gold Coast (Kaizaraku, 8 April 1871, on label). Collected by S. T. D. Aubinn. Sharpe Collection.
 The other two syntypes are also in the collection.

rubrocanus—Turdus rubrocanus J. E. Gray, 1846 Catalogue of Mammalia and Birds of Nepal and Thibet	. 31 pre-
sented by B. H. Hodgson to the British Museum: 81. The type specimens, presented by B. H. Hodgson, canno traced in the collection. The specimen listed as type by Seebe Cat. Bds. Brit. Mus., 5 (1881): 260 has no type status. Turubrocanus Hodgson, in J. E. Gray, Zoological Miscellany, (1844) is a nomen nudum. (See also appendix, Corrections and Additivol. 2.)	ohm, <i>irdus</i> : 83.
rubrocoronatum—Dicaeum rubrocoronatum Sharpe, 1876. Nature, 14:339. HOLOTYPE, Adult male. Reg. no. 1877.2 Port Moresby, South-East New Guinea. Collected by and chased of O. C. Stone.	.9.5.
rubrofrontata—Rhipidura rubrofrontata Ramsay, 1879. Proc. Linn. Soc. N.S.W., 4:82. Holotype, (Not fully adult, Ramsay, Proc. Linn. Soc. N.S.W., 6 (1881):178). Reg. no. 1 12.24.1. Lango, Guadalcanar, Solomon Islands. Collected J. Cockerell. Received in exchange from E. P. Ramsay of Sydney Museum.	895. l by
rubrogularis—Mesia argentauris rubrogularis Kinnear, 1925. Bull. Br. Orn. Cl., 45:75. HOLOTYPE, Adult male. Reg. 1924.12.21.136. Ngoi-Tio, Tonkin, 12 June 1924. Collected H. Stevens and presented by the Percy Sladen Trust and Sal Godman Exploration Fund.	no. d by
rubroniger—Munia rubroniger Hodgson, 1836 Asiat. Res., 19:153. SYNTYPE, Adult. Reg. no. 1843.1.13. Nepal. Collected and presented by B. H. Hodgson.	. 60 819.
rubropersonatus—Malimbus rubropersonatus Shelley, 1887 Ibis, (5) 5: 41, pl. 2. HOLOTYPE, Immature. Reg. no. 1889.7 240. (Denkera, inland forest of Fantee, vide Shelley, The Bird Africa, 4 (1905): 353), Gold Coast, 1870. Collected by Government. Shelley Collection.	7.25. ds of
rubropygia—Raya rubropygia Hodgson, 1839 J. Asiat. Soc. Beng., 8:36. SYNTYPE, Adult male (relaxed mou Reg. no. 1843.1.13.217. Nepal. Collected and presented by B Hodgson.	
rubropygium—Dicaeum trigonostigma rubropygium E. C. S. Ba 1921	. 45
ruddi—Apalis ruddi C. H. B. Grant, 1908	uese

rt	nddi—Heteronyx ruddi C. H. B. Grant, 1908	
ru	Ifa —Cyanoderma erythroptera rufa Chasen and Kloss, 1927 32 Bull. Br. Orn. Cl., 48:47. HOLOTYPE, Adult female. Reg. no. 1947.60.39. Sampit, south coast of Borneo, 113° E., 24 June 1914. Collected by C. Lumholz and presented by the Raffles Museum, Singapore.	
ru	Proc. zool. Soc. Lond., 1843	•
ru	in Byron, Voyage of H.M.S. Blonde, Appendix: 250. SYNTYPE, Adult. Reg. no. Old Vellum Cat. 20, no. 187a. Sandwich Islands. Collected by A. Bloxham. Presented by Lord Byron, R.N. A second syntype is also in the collection.	55
ru	Mirafra rufa Lynes, 1920	
ru	Ann. Mag. nat. Hist., (1) 11: 371. SYNTYPE, Adult (relaxed mount). Reg. no. 1842.2.15.35. (Cataguan, on label), Luzon, Philippines. Collected by and purchased of H. Cuming. Listed as the type by Sharpe, Cat. Bds. Brit Mus., 4 (1879): 343. A second, cleaner, syntype is also in the collection.	
ru	fescens—Anthus rufenscens Vigors and Horsfield, 1827 Trans. Linn. Soc. Lond., 15:230. HOLOTYPE, Adult. Reg. no. 1863.7.7.34. Australia. Collected by G. Caley. Presented as the type by the Linnean Society of London.	
ru	Ifescens—Atrichia rufescens Ramsay 1866	

The specimen was sent by Ramsay to Sclater and lent by him to Gould for the illustration in the Supplement to the Birds of Australia, (Dec. 1867): pl. 26, (vide Chisholm, Emu, 51 (1951): 103).

J. Gould.

rufescens—Bradypterus rufencens Sharpe and Bouvier, 1876 35 Bull. Soc. zool. France, 1:307. HOLOTYPE, Adult female. Reg. no. 1878.2.23.38. Landana, Congo, 1 June 1876. Collected by L. Petit. Purchased of A. Bouvier as the type.
rufescens—Drymoepus rufescens Hume, 1872
rufescens—Pachycephala rufescens Wallace, 1862 39 Proc. zool. Soc. Lond., 1862: 341. SYNTYPE, Adult male. Reg. no. 1873.5.12.510. Bouru Island, Moluccas, 1861. Collected by A. R. Wallace. Wallace Collection. See also Hartert, Novit. Zool., 5 (1898): 132. A syntype from Sula is also in the collection.
rufescens—Phlexis rufescens Sharpe, 1902
ruficapilla—Calliste ruficapilla P. L. Sclater, 1851
ruficapilla—Drymoica ruficapilla Smith, 1842
Illustrations of the Zoology of South Africa, Aves: pl. 73. SYNTYPE, Adult male. Reg. no. 1843.2.28.103. Interior of Cape Colony, (South Africa, on label). Collected by A. Smith. Purchased of the South African Society. See also Lynes, Ibis, (12) 6, Supplement, (1930): 658. There is another syntype in the collection.
ruficapilla—Drymoica ruficapilla Fraser, 1843
Proc. zool Soc. Lond., 1843:16. HOLOTYPE, Adult male. Reg. no. 1847.1.18.57. Nun River, mouth of Niger River, August 1841. Collected and presented by L. Fraser. Renamed D. anonyma Von Muller, 1855 on the grounds of preoccupation by D. ruficapilla Smith, 1842.
ruficapilla—Spermospiza ruficapilla Shelley, 1888 60
Proc. zool Soc. Lond., 1888: 30. HOLOTYPE, Adult female. Reg. no. 1887.9.28.83. Bellima, (Bongo country of the Upper Gazelle River, Congo, vide Shelley, Birds of Africa, 4 (1905): 296), 15 July, 1883. Collected and presented by Emin Pasha.

ruticeps 4/5
ruficauda—Erythropygia ruficauda Sharpe, 1882
Trans. Linn. Soc., (2), Zool. 1: 327. ?SYNTYPE, Adult male. Reg. no. 1876.8.28.26. Isabella de Basilan, Philippines, Oct. 1878. Collected by and purchased of J. B. Steere. Two specimens were collected. This appears to be one and the other is in the American Museum of Natural History. Hachisuka, Birds of the Philippine Islands, 2 (1935): 334, refers to the American specimen and erroneously attributes it to the Rothschild Collection.
ruficeps—Dendrocincla ruficeps P. L. Sclater and Salvin, 1868 . 2 Proc. zool. Soc. Lond., 1868: 54. HOLOTYPE, Adult. Reg. no. 1889.5.20.623. Panama, 1866. Collected by W. T. H. Chambers. Sclater Collection.
ruficeps—Dryoscopus ruficeps Shelley, 1885
ruficeps—Garrulax ruficeps Gould, 1862
ruficeps—Grallaria ruficeps P. L. Sclater, 1873 4 Proc. zool. Soc. Lond., 1873: 729. SYNTYPE, Adult. Reg. no. 1889.9.20.617. Medellin, Antioquia, Colombia, 1873. Collected by T. K. Salmon. Sclater Collection. Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 317, and so found in this (4) 1 (1877): pl 8. There are three other
and so figured in <i>Ibis</i> , (4) 1 (1877): pl.8. There are three other syntypes in the collection. ruficeps—Gymnopithys ruficeps Salvin and Godman, 1892

bia. Collected by T. K. Salmon. Salvin-Godman Collection. Listed as type by Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 302. Although the authors mentioned four specimens, now in the collection, they only specified the locality of one of them, Cauca. Of the other three one is from Remedios (Salmon) and two are from Bogota.

ruficeps—Megalurus ruficeps Tweeddale, 1877 Ann. Mag. nat. Hist., (4) 20:94. HOLOTYPE, Adult female. Reg. no. 1888.4.20.681. Monte Alban, Luzon, Philippines, Feb. 1877. Collected by A. H. Everett. Tweeddale Collection. (See also appendix, Corrections and Additions, vol. 2.)

ruficeps—Mixornis ruficeps Hodgson, 1845
ruficeps—Paradoxornis ruficeps Blyth, 1842
ruficollis—Cisticola ruficollis Walden, 1871
ruficollis—Ianthocincla ruficolis Jardine and Selby, 1838 32E Illustrations of Ornithology, 2nd ser., 1838 : pl. 21. HOLOTYPE, Adult. Reg. no. 1886.6.24.187. "Said to be Himalayas". Jardine Collection; purchased by Jardine of Carfrae, taxidermist, Edinburgh.
ruficollis—Montifringilla ruficollis Blanford, 1871
ruficollis—Pomatorhinus ruficollis Hodgson, 1836 32E Asiat. Res., 19:182. Syntype, Adult. Reg. no. 1843.1.13.621. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
ruficollis—Tanagra ruficollis Gmelin, 1789
ruficolor—Galerida theklae ruficolor Whitaker, 1898 17 Ibis, (7) 4:603. SYNTYPE, Adult male. Reg. no. 1939.12.14.22. Amamiz, Morocco, 17 May 1897. Collected by E. Dodson and presented by J. I. Whitaker. There is a female syntype in the collection.
rufidorsalis—Dryodromas rufidorsalis Sharpe, 1897 35 Bull. Brit. Orn. Cl., 6:48. HOLOTYPE, Adult. Reg. no. 1897. 10.31.8. Tsavo River, Kenya, 20 Sept. 1894. Collected and presented by F. J. Jackson

rufifrons—Cyornis rufifrons Wallace, 1865
rufifrons—Muscicapa rufifrons Latham, 1801
rufifrons—Prinia rufifrons Jerdon, 1840
with an original label can be accepted as a syntype. It was listed as the type by Sharpe, <i>Cat. Bds. Brit. Mus.</i> , 7 (1883): 247, and stated by him to have come from Madras. Renamed <i>P. buchanani</i> Blyth 1844 on the grounds of preoccupation by <i>P. rufifrons</i> Rüppell. 1835.
rufifrons—Stachyris rufifrons Hume, 1873
rufifrons—Trichostoma rufifrons Tweeddale, 1878 32B Proc. zool. Soc. Lond., 1878 : 616, pl. 38. Syntype, Adult male. Reg. no. 1888.4.20.1051. Puerto Princesa, Palawan, 10 Dec. 1877. Collected by A. Everett. Tweeddale Collection. The two other syntypes are also in the collection.
rufigaster—Dendrocitta rufigaster Gould, 1837
rufigena—Anaplectes rufigena Shelley, 1900 61

rufigene—Todirostrum rufigene P. L. Sclater and Salvin, 1877 . 9 Proc. zool. Soc. Lond., 1877: 522. HOLOTYPE, Adult. Reg. no. 1888.1.1.352. Mongi, Ecuador. Collected by C. Buckley. Salvin-Godman Collection.
rufigenis—Buarremon rufigenis Salvin, 1895
rufigenis—Calliste rufigenis P. L. Sclater, 1857
rufigula—Cyornis rufigula Wallace, 1865
rufigula—Myiagra rufigula Wallace, 1864
rufigularis—Hypsipetes rufigularis Sharpe, 1877
rufigularis—Siphia rufigularis Scully, 1879
rufilateralis—Rhipidura rufilateralis Sharpe, 1879
rufilatus—Nemura rufilatus Hodgson, 1845

rufimenta—Cinclosoma rufimenta Hodgson, 1836 Asiat. Res., 19:148. SYNTYPE, Adult. Reg. no. 1843.1.13.576. Nepal. Collected and presented by B. H. Hodgson. There are two other syntypes in the collection. rufinuchalis—Dryoscopus rufinuchalis Sharpe, 1895 Proc. zool. Soc. Lond., 1895: 479. SYNTYPE, Adult. Reg. no. 1895.7.7.11. Durro, Abyssinian Somaliland, 1 Dec. 1894. Collected and presented by Dr. A. Donaldson Smith. There is another syntype in the Academy of Natural Sciences. Philadelphia. rufipectus—Formicarius rufipectus Salvin, 1866 . Proc. zool. Soc. Lond., 1866: 73, pl. 8. SYNTYPE, Adult female. Reg. no. 1889.7.13.811. Santiago de Veragua. Collected by E. Arcé. Salvin-Godman Collection. The other syntype is in the collection. rufipennis—Cichladusa rufipennis Sharpe, 1901 Bull. Br. Orn. Cl., 12:35. SYNTYPE, Adult. Reg. no. 1895.5.1.929. Lamu, coastal East Africa. Shelley Collection. rufipennis—Pachycephala rufipennis G. R. Grav. 1858 Proc. zool. Soc. Lond., 1858: 178. HOLOTYPE, Female (immature, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883): 219). Reg. no. 1858.3. 10.71. Ké Island, south-west Pacific Ocean. Collected by A. R. Wallace Wallace Collection. rufipennis—Phacellodomus rufipennis P. L. Sclater, 1889 Proc. zool. Soc. Lond., 1889: 33. SYNTYPE, Adult. Reg. no. 1880.12.31.55. Bolivia. Collected by T. Bridges. Eyton Collection. This specimen is selected rather than the one listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 83, which appears to be wrongly labelled. rufipennis—Synallaxis rufipennis P. L. Sclater and Salvin, 1879 Proc. zool. Soc. Lond., 1879:620. SYNTYPE, Adult. Reg. no. 1889.5.14.249. Tilotilo, Bolivia, 1876. Collected by C. Buckley. Salvin-Godman Collection. There is another syntype in the collection. rufipennis—Trichastoma rufipennis Sharpe, 1872. Ann. Mag. nat. Hist., (4) 10:451. HOLOTYPE, Adult (female on label). Reg. no 1863.12.15.22. (Camma River, on label), Gaboon, West Africa. Collected by P. B. du Chaillu and purchased of S. Stevens. rufitergum—Garrulus glandarius rufitergum Hartert, 1903 Die Vögel der paläarktischen Fauna, 1:30. HOLOTYPE, Adult male. Reg. no. 1936.10.15.2. Tring, 21 Oct. 1895. Collected by W. Rothschild. Presented by the American Museum of Natural History, from the Rothschild Collection.

- rufiventer—Corythocichla brevicaudata rufiventer Delacour, 1927. 32B Bull. Br. Orn. Cl., 47:162. HOLOTYPE, Adult male. Reg. no. 1927.6.5.17. Djiring, south Annam, 12 March 1927. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China.

- rufivertix—Euphonia rufivertix Salvin, 1866 5

 Proc. zool. Soc. Lond., 1866: 71, pl. 7. Syntype, Adult male.

 Reg. no. 1885.6.4.158. (Santiago de Veragua, on label.) Collected by E. Arcé. Salvin-Godman Collection.

 There are several other syntypes in the collection.
- rufobrunneus—Ligurnus rufobrunneus G. R. Gray, 1862. . . . 59

 Ann. Mag. nat. Hist., (3) 10:444. HOLOTYPE, Adult. Reg. no. 1968.24.1. West Africa. Collected and presented by E. Sabine.
- rufocaudatus—Pycnonotus rufocaudatus Eyton, 1845 21
 Ann. Mag. nat. Hist., (1) 16:228. Syntype, Adult. Reg. no. 1846.3.4.25. Malacca. Collected and presented by Capt. A. Charlton.

There is a second syntype in the collection.

There is another syntype in the collection.

rufocinerea—Grallaria rufocinerea P. L. Sclater and Salvin, 1879 . 4

Proc. zool. Soc. Lond., 1879: 526. HOLOTYPE, Adult. Reg. no. 1889.9.20.618. Santa Elena, Antioquia. Collected by T. K. Salmon. Sclater Collection.

rufofronta—Rhissidura rufofronta?, 1879
rufofrontata—Rhipidura rufofrontata Ramsay, 1879 38 see R. rubrofrontata Ramsay, 1879. Misspelling of original name.
rufogularis—Drymoica rufogularis Fraser, 1843
rufogularis—Ianthocincla rufogularis Gould, 1835 32B Proc. zool. Soc. Lond., 1835 : 48. HOLOTYPE, Adult (relaxed mount). Reg. no. 1838.5.12.107. Himalayas. Purchased of J. Gould.
rufogularis—Minla rufogularis Mandelli, 1873 32B Stray Feathers, 1:416. Holotype, Adult. Reg. no. 1886.10.1. 6200. Bhotan Doars, north-east Bengal, Jan. (in text; Dec. 1872 in label). Collected by L. Mandelli. Hume Collection.
rufogularis—Ruticilla rufogularis Moore, 1854
rufogularis—Synallaxis rufogularis Gould, 1839
rufolateralis—Piezorhynchus rufolateralis G. R. Gray, 1858 38 Proc. zool. Soc. Lond., 1858: 176. SYNTYPE, Adult male. Reg. no. 1858.3.10.77. Aru Islands. Collected by A. R. Wallace. Wallace Collection. The female syntype is also in the collection.
rufolateralis—Smithornis rufolateralis G. R. Gray, 1864 1 Proc. zool. Soc. Lond., 1864: 143, pl. 16. HOLOTYPE, Adult. Reg. no. 1864.2.7.1. West Africa (Gold Coast, vide Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 389). Purchased of J. Gould.
rufopicta—Estrilda rufopicta Fraser, 1843

rufostrigata—Linota rufostrigata Walton, 1905 59 Bull. Brit. Orn. Cl., 15:93. HOLOTYPE, Adult male. Reg. no. 1905.12.31.281. Khamba Jong, South Tibet, 9 Oct. 1903. Collected and presented by H. J. Walton.
rufovelatus—Euplectes rufovelatus Fraser, 1842 61 Proc. zool. Soc. Lond., 1842:142. Syntype, Adult male. Reg. no. 1847.1.18.50. Clarence, Fernando Po. Early 1842. Collected by and purchased of L. Fraser. There are two other syntypes in the collection.
rufula—Prinia rufula Godwin-Austen, 1874
rufum—Heteropelma rufum Pelzeln, 1869
rugillatus—Garrulax rugillatus Swinhoe, 1860
rupicola—Hirundo rupicola Hodgson, 1836
rupicola—Sitta rupicola Blanford, 1873
russata—Rhipidura russata Tristram, 1879

russatus—Euscarthmus russatus Salvin and Godman, 1883 9 Ibis, (5) 2:445. SYNTYPE, Adult male. Reg. no. 1888.1.1.380. Roraima, 5,000', British Guiana, 7 Sept. 1883. Collected by H. Whitely. Salvin-Godman Collection. There are several other syntypes in the collection.
russula—Cettia russula Slater, 1897
rustica—Cisticola rustica Wallace, 1863
ruticilloides—Phoenicura ruticilloides Hodgson, MS 31 MS drawings of Birds of Nepal: pl. 75, nos. 403, 405. Three specimens in the collection were listed as types by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 345 but this is a nomen nudum.
ruwenzori—Cryptospiza salvadorii ruwenzori W. L. Sclater, 1925 . 30 Bull. Brit. Orn. Cl., 46:45. HOLOTYPE, Adult male. Reg. no. 1906.12.23.319. Mubuku Valley, 7,000', Eastern Ruwenzori, 12 March 1906. Collected by R. B. Woosnam and presented by the subscribers to the Ruwenzori Expedition.
ruwenzoria—Mirafra africana ruwenzoria Kinnear, 1921 17 Bull. Br. Orn. Cl., 41:139. HOLOTYPE, Adult male. Reg. no. 1906.12.23.473. Mokia, 3,400', south-east Ruwenzori, East Africa, 4 June 1906. Collected by R. E. Dent and presented by the subscribers to the Ruwenzori Expedition.
ruwenzoriae—Tchitrea perspicillata ruwenzoriae C. H. B. Grant and Mackworth-Praed, 1940
ruwenzorii—Apalis ruwenzorii Jackson, 1904
ruwenzorii—Nilaus minor ruwenzorii Bannerman, 1923 23 Ibis, (11) 5:698. HOLOTYPE, Adult male. Reg. no. 1906.12.23.

- 751. Mokia, 3,400', south-east Ruwenzori, 27 May 1906. Collected by R. E. Dent and presented by the subscribers to the Ruwenzori Expedition.
- ruwenzorii—Tarsiger ruwenzorii Ogilvie-Grant, 1906 31 Bull. Br. Orn. Cl., 19:33. SYNTYPE, Adult male. Reg. no. 1906. 12.23.1444. Mubuku Valley, 7,000', east Ruwenzori, Uganda, 14 Jan. 1906. Collected by R. B. Woosnam and presented by the subscribers to the Ruwenzori Expedition.

The female syntype is also in the collection.

- saba—Malacopteron magnum saba Chasen and Kloss, 1930. Bull. Raffles Mus., 4:75. HOLOTYPE, Adult male. Reg. no. 1947.60.53. Samawang River, near Sandakan. North Borneo, 11 July 1927. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore.
- sabini—Thamnophilus sabini J. E. Gray, 1831 23 Zoological Miscellany, 1:7. SYNTYPE, Adult male (relaxed mount). Reg. no. Old Vellum Cat. 10.28a. Sierra Leone. Collected and presented by Capt. Sabine. Listed as type by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 143.
- sabota—Mirafra sabota Smith, 1836 Report of the Expedition for exploring Central Africa . . . 1834: Appendix: 47. Syntype, Adult (relaxed mount). Reg. no. 1838. 6.9.8. Between Latakoo and the Tropic (vide Smith, Illustrations of the Zoology of South Africa, 2 (Aves): pl. 89, fig. 1; type locality restricted to Rustenberg, western Transvaal, by Macdonald, Contributions to the Ornithology of western South Africa, (1957):98). Collected by A. Smith and purchased of the South African Museum. There are two other syntypes in the collection; others are in the Royal Scottish Museum, Edinburgh, and the Academy of Natural Sciencies, Philadelphia.
- sacculatus—Criniger ochraceus sacculatus Robinson, 1915 . *Ibis*, (10) 3:746. HOLOTYPE, Adult female. Reg. no. 1936.4.12. 1883. Genting Bidai, 2,300', Selangor, 14 May 1908. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore.
- sagittata—Sylvia sagittata Latham, 1801 36 Index Ornithologicus, Suppl.: liv. Syntypes, Water-colour drawings. New South Wales. Nos. 196, 197, 198, in Watling, Original water-colour drawings. . . . Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 142.

Vol. 2, no. 9 Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2). See also Mathews,

Birds of Australia, 9 (1922): 402.

- sakalava—Ploceus sakalava Hartlaub, 1861 61
 Ornithologisches Beitrage zur Fauna Madagascar's: 54. HOLOTYPE,
 Adult male (relaxed mount). Reg. no. 1845.7.6.10. Madagascar.
 Collected by and purchased of A. Smith.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 579.

There are several other syntypes in the collection.

- salimalii—Cisticola juncidis salimalii Salim Ali and Whistler, 1936 35 J. Bombay nat. Hist. Soc., 38:487. Syntype, Adult male. Reg. no. 1936.6.18.1. Peermade 3,200', (High Range), Travancore, 25 Feb. 1933. Collected and presented by Salim Ali.

This specimen was presented as the type but a number were referred to in the description and no particular specimen was

selected.

- salimalii—Erythrina synoica salimalii R. Meinertzhagen, 1938 . 59
 Bull. Brit. Orn. Cl., 58:95. HOLOTYPE, Adult male. Reg. no.
 1965.M.39. Akrobat, 5,000', North Afghanistan, 26 April 1937.
 Collected by R. Meinertzhagen. Meinertzhagen Collection.
- sallaei—Setophaga sallaei Bonaparte and Sclater, 1856. . . . 54 in Bonaparte, C. r. Acad. Sci. Paris, 42:957. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1857.7.30.21. (Collector's no. 129.) Cordova, Vera Cruz, 8 April 1855. Collected by and purchased of A. Sallé.

Sclater, *Proc. zool. Soc. Lond.*, 1856: 292, pl. 120, redescribed as *Granatellus sallaei* Bonaparte's bird and stated that only the above single specimen was received; confirming in *Proc. Zool. Soc. Lond.*, 1858: 97, that the type was in the British Museum. His specimen listed as the type by Sharpe, *Cat. Bds. Brit. Mus.*, 10 (1885): 372, was not mentioned in Sclater's *Catalogue of a collection of American birds*, (1862) and is therefore of a later date. The Paris Museum specimen, listed by van Rossem, *Bull. Mus. Comp. Zool. Harv.*, 77 (1934): 404, cannot be a type.

See also Hellmayr, Novit. Zool., 13 (1906): 317.

salmoni—Ostinops salmoni P. L. Sclater, 1883
salmoni—Rhopias fulviventris salmoni Chubb, 1918 4 Ann. Mag. nat. Hist., (9) 2:124. Syntype, Adult male. Reg. no. 1889.7.10.349. Remedios, north Colombia. Collected by T. K. Salmon. Salvin-Godman Collection. Chubb stated the type to be a specimen collected by Salmon at Remedios. Salmon obtained two males at this locality, both of which are in the collection.
salomonis—Edoliisoma salomonis Tristram, 1879 20 Ibis, (4) 3:440. Syntype, Adult female. Reg. no. 1880.2.10.1. Makira Harbour, San Christoval, Solomon Islands, 3 Oct. 1878. Collected by Lt. G. E. Richards, R.N., and purchased of H. B. Tristram. The second adult female syntype is also in the collection; the immature female is in the Liverpool Museum.
saltator—Merula saltator Hill, 1842
salvadorii—Galeopsar salvadorii Sharpe, 1891 62 Ibis, (6) 3: 241, pl. 6. HOLOTYPE, Adult female. Reg. no. 1893. 12.1.1. Turquel, Suk, northern Kenya, 15 Jan. 1890. Collected and presented by F. J. Jackson.
salvadorii—Nectarinia salvadorii Shelley, 1903
salvadorii—Sphecotheres salvadorii Sharpe, 1877
salvadorii—Zosterops salvadorii. See appendix, Corrections and Additions, vol. 2

salvini—Cyphorhinus salvini Sharpe, 1881
Cat. Bds. Brit. Mus., 6: 292, pl. 18. HOLOTYPE, Adult. Reg. no. 1858.2.4.42. Rio Napo, Ecuador. Purchased of Verreaux.
salvini—Empidochanes salvini P. L. Sclater, 1888 9
Cat. Bds. Brit. Mus., 14:218. SYNTYPE, Adult male. Reg. no. 1888.1.1.1335. Roraima, 5,000', British Guiana, 20 Aug. 1883. Collected by H. Whitely. Salvin-Godman Collection.
There are several other syntypes in the collection.
Proc. zool. Soc. Lond., 1877: 324. SYNTYPE, Adult male. Reg. no. 1888.1.13.79. Tumbez, Peru, 17 (Caerwcal) 1876. Collected by Stolzmann. Sclater Collection. Listed by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 25, as an "authentic specimen".
salvini—Pachyrhamphus salvini Richmond, 1899
Auk, 16: 186. New name for P. similis Salvin, 1895 (q.v.) which is preoccupied by P. similis Cherrie, 1891.
salvini—Phoenicothraupis salvini Berlepsch, 1883 51
Ibis, (5) 1:487. Syntype, Adult male. Reg. no. 1885.6.7.481. Chisec, Vera Paz, Guatamala. Collected by O. Salvin. Salvin-Godman Collection. This is specimen number 3 of the nine syntypes listed.
salvini—Pseudochloris salvini Chubb, 1919
Bull. Brit. Orn. Cl., 39:70. SYNTYPE, Male (Immature, vide Hellmayr, Catalogue of birds of the Americas, 11 (1938):313, note). Reg. no. 1896.10.6.344. Vina, 5,500', Huamachuco, Northern Peru, March 1894. Collected by O. T. Baron. Salvin-Godman Collection. The female syntype is also in the collection.
salvini—Vireo pallens salvini van Rossem, 1934
Bull. Mus. Comp. Zool. Harv., 77: 418. HOLOTYPE, Adult. Reg. no. 1886.9.9.529. Holbox Island ("Jolbox", on label), Yucafan, Dec. 1885. Collected by G. F. Gaumer. Salvin-Godman Collection.
salwinensis—Abrornis superciliaris salwinensis, E. C. S. Baker, 1924 35
Bull. Brit. Orn. Cl., 44: 42. HOLOTYPE, Adult male. Reg. no. 1886.4.1.3364. Salween, Tenasserim Province, Burma, 16 Feb.
1874. Collected by W. Davison. Hume Collection.
samaliyae—Serinus mozambicus samaliyae C. M. N. White, 1947. 59
Bull Brit. Orn. Cl., 68: 11. HOLOTYPE, Adult male (female, on label). Reg. no. 1947.80.33. Pempele Pool (between Musonic and Solwezi, Southern Belgian Congo), 2 May 1939. Collected and presented by C. M. N. White. This is certainly the holotype specimen in spite of the descrepancy between description and label concerning the sex.
samarensis—Corvus samarensis Steere, 1890

- to the Philippines: 23. HOLOTYPE, Adult male. Reg. no. 1896. 6.6.18. Catbalogan, Samar, Philippines, 12 April 1890, Collected by J. B. Steere. Steere Collection samarensis—Hypothymis samarensis Steere, 1890. . 37A List of birds and mammals collected by the Steere Expedition to the Philippines: 16. SYNTYPE, Reg. no. 1896.6.6.177. Samar. Philippines, (Catbalogan, 4 April, on label). Collected by J. B. Steere on his second Expedition, and purchased of him as the type. This specimen was labelled "type" by the author, and listed as such by Hachisuka, Birds of the Philippine Islands, 2 (1935): 316. The female syntype is also in the collection. samarensis—Oriolus samarensis Steere, 1890 63 List of Birds and Mammals collected by the Steere Expedition to the Philippines: 17. HOLOTYPE, Adult male. Reg. no. 1896.6.6.41. Catbalogan, Samar, Philippines, April. Collected by D. W. Worcester. Steere Collection. samarensis—Orthotomus samarensis Steere, 1890 List of Birds and Manunals collected by the Steere Expedition to the Philinnines: 20. HOLOTYPE, Adult male. Reg. no. 1844.6.6.244. Samar, Philippines, (Catbalogan, 5 April, on label). Collected by J. B. Steere, Steere Collection. samarensis—Prionochilus samarensis Steere, 1890. 45 List of Birds and Mammals collected by the Steere Expedition to the Philippines: 22. SYNTYPE, Adult male. Reg. no. 1896.6.6.565. Catbalogan, Samar, Philippines, (31 March 1888, on label). Collected by J. B. Steere on the 2nd Expedition (vide Salomonsen, Amer. Mus. Novit., 1990 (1960): 20). The female syntype is also in the collection. samarensis—Setaria samarensis Steere, 1890 37 List of Birds and Mammals collected by the Steere Expedition to the Philippines: 16. SYNTYPE, Reg. no. 1896.6.6.191. Samar, Philippines, (Catbalogan, 2 April, on label). Collected by J. B. Steere on his second expedition, and purchased of him as the type. Listed as the type by Hachisuka, Birds of the Philippine Islands, 2 (1935): 334. The female syntype is also in the collection. samharensis—Ammomanes samharensis Shelley, 1902 17 The Birds of Africa, 3:99, pl. 21. HOLOTYPE, Adult female. no. 1869.10.16.253. Amba, Samhar desert, Abyssinia, 22 Aug. 1868. Collected by W. T. Blanford and presented by the Governor and Council of Bombay. sancta—Rhipidura sancta Sharpe, 1899 . 37A
 - Bull. Brit. Orn. Cl., 10: 29. SYNTYPE, Adult male. Reg. no. 1900. 1.10.16. Espiritu Santa Island, New Hebrides (Hoy Head, 17 Sept. 1899, vide Sharpe, Ibis, (7) 6 (1900): 340). Collected and presented by A. M. Farquhar from the voyage of H.M.S. Wallaroo.

- sanctaehelenae—Estrilda sanctaehelenae Shelley, 1903 . . . 60

 Bull. Brit. Orn. Cl., 13:74. HOLOTYPE, Adult. Reg. no. 1889.7.

 20.456. St. Helena, 4 Jan. 1874. Collected by G. E. Shelley. Shelley Collection.
- sanctaeluciae Margarops sanctaeluciae P. L. Sclater, 1880 . . . 31
 Ibis, (4) 4:73. SYNTYPE, Adult. Reg. no. 1886.8.2.175. Saint Lucia, West Indies, 1871. Collected by J. E. Semper. Sclater Collection.

The second syntype is also in the collection.

- sanctaemarthae—Rhamphocaenus sanctaemarthae P. L. Sclater, 1861 34 Proc. zool. Soc. Lond., 1861: 380. HOLOTYPE, Adult. Reg. no. 1889.9.20.391. Santa Martha, New Colombia. Collected for Verreaux. Sclater Collection.
- sanghirana—Anthreptes malacensis sanghirana Hachisuka, 1926 46 Bull. Brit. Orn. Cl., 47:68. HOLOTYPE, Adult male. Reg. no. 1895.9.9.103. Sanghir Island, North of Celebes, 8 Sept. 1876. (Collected by Laglaize, on label). Shelley Collection.
- sanguinea—Fringilla sanguinea Gould, 1838 59

 Proc. zool. Soc. Lond., 1837 (publ. 1838) : 127. HOLOTYPE, Adult male. Reg. no. 1855.12.19.10. Trebizond, Erzeroum, Persia. Collected by K. E. Abbott. Purchased of the Zoological Society of London.

The other syntypes are also in the collection.

sanguinolenta—Certhia sanguinolenta Latham, 1801 4
Index Ornithologicus, Suppl.: xxxvii. ?HOLOTYPE, Water-colour drawing. New South Wales. No. 109 in Watling, Original water-colour drawings. . . . Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 128.

Vol. 2, no. 78 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews,

Birds of Australia, 11 (1924): 305.

- sanguinolenta—Cryptospiza reichenowi sanguinolenta Vincent, 1933 60 Bull. Brit. Orn. Cl., 53:148. HOLOTYPE, Adult male. Reg. no. 1933.3.1.6. Mlanje Mountain, 6,300', Southern Nyasaland, 3 Jan. 1932. Collected and presented by J. Vincent.

The second syntype is also in the collection.

- santhome—Serinus mozambicus santhome Bannerman, 1921 . . . 59
 Bull. Brit. Orn. Cl., 41:137. HOLOTYPE, Adult male. Reg. no. 1911.12.23.3951. St. Thomas Island, Gulf of Guinea, 23 Feb. 1909. Collected by Boyd Alexander. Boyd Alexander Bequest.
- santovestris—Aplonis santovestris Harrison and Marshall, 1937. 62 Bull. Br. Orn. Cl., 57: 148. HOLOTYPE, Adult female. Reg. no. 1937.5.15.1. Mount Watiamasam, 4,000′, Espiritu Santo, New Hebrides, 25 May 1934. Collected and presented by T. H. Harrisson and A. J. Marshall.
- sarayacuensis—Corythopis sarayacuensis Chubb, 1918 5

 Bull. Br. Orn. Cl., 38:48. HOLOTYPE, Adult. Reg. no. 1889.
 7.10.961. Sarayacu, Ecuador, Feb. 1880. Collected by C. Buckley. Salvin-Godman Collection.
- sarkari—Dendrocitta formosae sarkari Kinnear and Whistler, 1930 71 Bull. Brit. Orn. Cl., 51:17. HOLOTYPE, Adult male. Reg. no. 1937.12.21.49. Anantagiri, 3,000', Vizagapatam, South India, 5 Feb. 1930. Collected by V. S. La Personne. Presented by the Bombay Natural History Society.

Tristram, Catalogue of a collection of birds, (1889): 206, claimed to have the types at that date, but this syntype came to Seebohm

later.

saturata—Amblyospiza saturata Sharpe, 1908 61 Ibis, (9) 2:353. SYNTYPE, Adult male. Reg. no. 1909.8.8.47. Ja River, Cameroons, 29 June 1906. Collected by and purchased of G. L. Bates. There are other syntypes in the collection.
saturata—Anuropsis malaccensis saturata Robinson and Kloss, 1920
saturata—Argya saturata Sharpe, 1895
saturata—Cinnyris saturata Hodgson, 1837
saturata—Erythropygia ruficauda saturata Neumann, 1920 31 J. Orn., 68:83. HOLOTYPE, Adult male. Reg. no. 1890.3.3.3. Yambuya, Aruwimi River, central Africa. Collected by J. S. Jameson and presented by Mrs Jameson.
saturata—Heterocnemis saturata Salvin, 1885
saturata—Leioptila saturata Walden, 1875
saturata—Rhipidura saturata Sharpe, 1879

saturata—Volvocivora saturata Swinhoe, 1870 20

Ibis, (2) 6:242. SYNTYPE, Adult male. Reg. no. 1897.7.2.56.

Hainan, south China, Feb. 1868. Collected by R. Swinhoe.

Seebohm Collection.

This is the author's selected male. His selected female syntype is also in the collection.

- saturatior—Cercomacra tyrannina saturatior Chubb, 1918 . . . 4

 Bull. Br. Orn. Cl., 38:85. SYNTYPE, Adult male. Reg. no. 1922.

 3.5.2142. Ituribisi R., British Guiana, Feb. 1907. Collected by F. V. McConnell. McConnell Collection.

 This is the outbor's selected male. His selected female and

This is the author's selected male. His selected female and twenty-two other syntypes are in the collection.

- saturatior—Cyornis poliogenys saturatior Robinson and Kloss, 1927 37 Bull. Brit. Orn. Cl., 48:43. HOLOTYPE, Adult female. Reg. no. 1886.4.1.4117. Near Dibrughar, Upper Assam, 2 March 1880. Collected by J. R. Cripps. Hume Collection.
- saturatior—Dendrocitta vagabunda saturatior Ticehurst, 1922. 71

 Bull. Brit. Orn. Cl., 42:56. HOLOTYPE, Adult female. Reg. no. 1886.3.1.650. Kankariyet Mountains, Amherst Distr., Lower Burma, 3 Sept. 1878. Collected by J. Darling. Hume Collection.
- saturatior—Gampsorhynchus saturatior Sharpe, 1888 32B Proc. zool. Soc. Lond., 1888 : 273. HOLOTYPE, Adult female. Reg. no. 1888.3.18.61. Gunong Batu Putch, 3,400', Perak, Malaya. Collected and presented by L. Wray.
- saturatior—Heliocorys modesta saturatior Bannerman, 1922 . . . 17
 Bull. Br. Orn. Cl., 42:141. HOLOTYPE, Adult male. Reg. no. 1922.11.25.263. Tibati, 2,800', Cameroon, 1 Aug. 1921. Collected by and purchased of G. L. Bates.
- saturatior—Hirundo senegalensis saturatior Bannerman, 1923. . 18
 Bull. Br. Orn. Cl., 43:85. HOLOTYPE, ?Subadult. Reg. no.
 1895.5.1.1983. Accra, Gold Coast, 11 Feb. 1872. Collected by
 G. E. Shelley. Shelley Collection.
- saturatior—Ixops poliotis saturatior Rothschild, 1921 32B Novit. Zool., 28:38. HOLOTYPE, Adult male (labelled as female). Reg. no. 1921.7.15.298. Shweli-Salwin Divide, 25°40′ N., 7,000′, Yunnan, May 1919. Collected by G. Forrest and presented by Stephenson Clarke.

- saturatior—Paradoxornis unicolor saturatior Rothschild, 1921 . 33
 Novit. Zool., 28: 54. SYNTYPE, Adult male. Reg. no. 1921.7.15.
 545. Shweli-Salwin Divide, 9.000, Yunnan, July 1919. Collected by G. Forrest and presented by Stephenson Clarke.
 There are two other syntypes in the collection.
- saturatior—Procarduelis rubescens saturatior Rothschild, 1923 . 59
 Bull. Brit. Orn. Cl., 43:12. HOLOTYPE, Adult male. Reg. no.
 1922.12.7.517. Shweli-Salvin Divide, 9,000', North Yunnan,
 south-west China, May 1919. Collected by G. Forrest.
 Stephenson Clarke Collection.

- saturatior—Tchitrea affinis saturatior Salomonsen, 1933 . . . 38

 Ibis, (13) 3:732. HOLOTYPE, Adult female. Reg. no. 1886.4.1.
 2907. Buxa Duars, Bhutan, Feb. 1878. Collected by L. Mandelli. Hume Collection.
- saturatus—Acanthopneuste nitidus saturatus Baker, 1924 . . . 35

 Bull. Brit. Orn. Cl., 44:62. HOLOTYPE, Adult male. Reg. no. 1919.12.20.391. Dalan, (error for Daban), South Annam, March 1918. Collected by C. B. Kloss. Presented by H. C. Robinson and C. B. Kloss.
- Saturatus—Ammomanes saturatus Ogilvie-Grant, 1900 17
 Novit. Zool., 7: 249. Holotype, Adult female. Reg. no. 1900.
 8.5.144. Jebel Manif, north of Lahej, south Arabia, 13 Aug. 1899
 (error: 14 Aug. on label). Collected by A. B. Percival on the Percival-Dodson Expedition.
- saturatus—Cephalopyrus flammiceps saturatus Whistler, 1924. 41
 Bull. Brit. Orn. Cl., 45:15. HOLOTYPE, Adult male. Reg. no.
 1886.11.1.644. Native Sikkim, March 1878. Collected for L.
 Mandelli. Hume Collection.

- saturatus—Molpastes cafer saturatus Whistler and Kinnear, 1932. 21 J. Bombay nat. Hist. Soc., 35:754. Holotype, Adult female. Reg. no. 1937.12.21.241. Anantagiri, 3,000′, Eastern Ghats, 22 Feb. 1930. Collected by V. S. La Personne and presented by the Bombay Natural History Society; collected on the Vernay Scientific Survey of the Eastern Ghats.
- saturatus—Pericrocotus peregrinus saturatus E. C. S. Baker, 1920 . 20 Bull. Br. Orn. Cl., 40:115. HOLOTYPE, Adult male. Reg. no. 1873.5.12.1122. West Java, 1863. Collected by A. R. Wallace. Wallace Collection.
- saturatus—Platyrhynchus saturatus Salvin and Godman, 1882 . 9

 Ibis, (4) 6:781 HOLOTYPE, Adult male. Reg. no. 1888.1.1.296.

 Merume Mountains, British Guiana, 28 July 1881. Collected by H. Whitely. Salvin-Godman Collection.
- saturatus—Pomatorhinus ruficollis saturatus Delacour, 1927

 Bull. Br. Orn. Cl., 47:159. HOLOTYPE, Adult male. Reg. no. 1927.6.5.12. Tam Dao, central Tonkin, 26 Nov. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indochina.

Other early specimens from Mandelli came to the collection by way of the Hume Collection. A Hume collection specimen, listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 420, cannot now be traced.

- saturatus—Tribura luteoventris saturatus Ticehurst, 1941 . . . 35

 Ibis, (14) 5:318. HOLOTYPE, female (immature, on label). Reg.
 no. 1948.80.1138. Thayetmyo-Minbu border, 5,000′, Southern
 Chin Hills, 5 Dec. 1938. Collected by B. E. Smythies. Presented
 by P. F. Garthwaite and H. C. Smith.

- scapularis—Jora scapularis Horsfield, 1821 Trans. Linn. Soc. Lond., 13:152. SYNTYPE, Adult (relaxed mount). Reg. no. 1880.1.1.4744. Java, 1811–1817. Collected by T. Horsfield and presented by the Honourable East India Company Museum. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 12. schistacea—Grandala schistacea Hodgson, 1843 J. Asiat. Soc. Bengal, 12: pl. opp. 450. HOLOTYPE, Adult female (error, Immature, vide J. E. Gray, Catalogue of Mammalia and Birds of Nepal and Thibet presented to the British Museum by B. H. Hodgson, 1846:69; relaxed mount). Reg. no. 1844.12.27.2. "North region of Cachar". Presented by B. H. Hodgson. schistacea—Hypocnemis schistacea P. L. Sclater, 1858 Proc. zool. Soc. Lond., 1858: 252. HOLOTYPE, Adult male. Reg. no. 1858.1.11.18. Rio Javarri, north-east Peru. Collected by H. W. Bates and purchased of S. Stevens. schistacea—Melaenornis schistacea Sharpe, 1895 Proc. zool. Soc. Lond., 1895: 481. HOLOTYPE, Adult. Reg. no. 1895.7.7.12. Darro Mountains, western Somaliland, 19 Nov. 1894. Collected and presented by Dr. A. Donaldson Smith. schistacea—Nigrita schistacea Sharpe, 1891 Ibis, (6) 3:118. HOLOTYPE, Adult female. Reg. no. 1893.12.1.7. Sotik, Kenya, (3 Oct. 1889, on label). Collected and presented by F. J. Jackson. schistaceiceps—Dicaeum schistaceiceps G. R. Gray, 1860 Proc. zool. Soc. Lond., 1860: 349. SYNTYPE, Adult male. Reg. no. 1860.2.4.33. Batchian, Moluccas. Collected by A. R. Wallace. Wallace Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 29. schistaceiceps—Todirostrum schistaceiceps P. L. Sclater, 1859.

reference is given.

schistaceus—Enicurus schistaceus Hodgson, 1836
schistaceus—Rhectes nigrescens schistaceus von Reichenow, 1900 39 Orn. Mber., 8: 187. HOLOTYPE, Adult male. Reg. no. 1901.4.28. 36. Aroa River, 4,000', south-east New Guinea, Dec. 1899. Collected by and purchased of E. Weiske.
schisticeps—Budytes schisticeps Hodgson, 1844
schisticeps—Culipeta schisticeps J. E. Gray, 1846 35 Catalogue of Mammalia and Birds of Nepal and Thibet presented by B. H. Hodgson to the British Museum; 67, 153. SYNTYPE, Adult. Reg. no. 1843.1.13.388. Nepal. Collected and presented by B. H. Hodgson. Gray used a Hodgson MS. name. By the register date this specimen must have been available at the time of description. It is almost tail-less. There are several
other syntypes in the collection.
schisticeps—Pomatorhinus schisticeps Hodgson, 1836 32B Asiat. Res., 19:181. SYNTYPE, Adult. Reg. no. 1843.1.13.618. Nepal. Collected and presented by B. H. Hodgson. There are several other syntypes in the collection.
schisticeps—Ruticilla schisticeps J. E. Gray, 1846
schistilatus—Orthotomus (Horeites) schistilatus Hodgson, 1845 . 35 Proc. zool. Soc. Lond., 30. Syntype, Adult. Reg. no. 1860.4.16. 359. Nepal. Presented by the Honourable East India Company Museum. Listed as the type by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 145. There are other syntypes in the collection.
scindicus—Pyctorhis altirostris scindicus Harington, 1915 32B J. Bombay nat. Hist. Soc., 23: 424. HOLOTYPE, Adult male. Reg. no. 1893.11.30.4. Mangnami (on label), near Sukkar, Sind, India, 12 Nov. 1876. Collected and presented by W. J. Blanford.
sclateri—Calandrella sclateri Shelley, 1902

sclateri—Cisticola nuchalis sclateri Bannerman, 1920 35 Bull. Brit. Orn. Cl., 41:7. HOLOTYPE, Adult male. Reg. no. 1920.12.9.9. Akonolinga, Nyong River, Cameroons, 8 Jan. 1913. Collected and presented by G. L. Bates.
sclateri—Chrysomitris sclateri Sharpe, 1888
sclateri—Cyanomitra olivacea sclateri Vincent, 1934
sclateri—Dendrocitta rufa sclateri E. C. S. Baker, 1922
Sclateri—Doliornis sclateri Taczanowski, 1874
Sclateri—Euprinodes sclateri Boyd Alexander, 1903
Sclateri—Hylophilus sclateri Salvin and Godman, 1883 56 Ibis, (5) 1:205. SYNTYPE, Adult male. Reg. no. 1885.3.10.193. Roraima, British Guiana, 3 Dec. 1881. Collected by H. Whitely. Salvin-Godman Collection. The female syntype is also in the collection.
Proc. zool. Soc. Lond., 1879: 265, pl. 25. Holotype, Adult male. Reg. no. 1881.3.29.76. New Britain, (=Palakuuru Island, off New Britain, vide Sclater, Proc. zool. Soc. Lond., 1879: 448; =Credner Island, vide Peters, Check-list of birds of the World, 12 (1967): 361), June 1878. Collected by the Rev. G. Brown. Purchased of E. Gerrard.
clateri—Onithion sclateri Berlepsch and Taczanowski, 1883 9 Proc. zool. Soc. Lond., 1883: 554. New name for Eupsilostoma pusillum P. L. Sclater, 1860, (q.v.) on the grounds of preoccupation by Myiopatis pusilla Cabanis and Heine, 1859.
clateri—Parus sclateri. See appendix, Corrections and Additions, vol. 2

sclateri—Philemon sclateri G. R. Gray, 1870
sclateri—Polioptila sclateri Sharpe, 1885
sclateri—Thripophaga sclateri Berlepsch, 1883
scotti—Zosterops scotti Neumann, 1899
scratchleyana—Munia scratchleyana Sharpe, 1898 60 Bull. Brit. Orn. Cl., 7:60. HOLOTYPE, Adult. Reg. no. 1898.5. 31.14. Mount Albert Edward, south-east New Guinea. Collected by and purchased of W. Blyth & Co. Prepared from a specimen preserved in spirit.
scullii—Hirundo scullii Seebohm, 1883
scutata—Synallaxis scutata P. L. Sclater, 1859
sechuenensis—Motacilla felix var. sechuenensis Swinhoe, 1870. 19 Proc. zool. Soc. Lond., 1870: 123, fig. 2. Syntype, Adult male. Reg. no. 1898.10.20.448. Yunyang, Szechuen, China, 23 April 1869. Collected by R. Swinhoe. Seebohm Bequest. There are several other syntypes in the collection.
secunda—Mirafra secunda Sharpe, 1890

seebohmi—Alaudula seebohmi Sharpe, 1890	
seebohmi—Cettia seebohmi Ogilvie-Grant, 1894	
Proc. zool. Soc. Lond., 1879: 177. SYNTYPE, Adult. Reg. no. 1879.5.29.26. Antananarivo, Madagascar. (Collected by Lormier, vide, Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 100). Purchased of Higgins. This specimen is labelled "Type", but there are three other syntypes in the collection.	•
Seebohmi—Lanius seebohmi Gadow, 1883	23
seebohmi—Merula seebohmi Sharpe, 1888	31
Seebohmi—Phylloscopus seebohmi Hume, 1877	35
seebohmi—Saxicola seebohmi Dixon, 1882	;
seimundi—Bradyornis infuscatus seimundi Ogilvie-Grant, 1913 Ibis, (10) 1:536. SYNTYPE, Adult male. Reg. no. 1903.3.9.298. Deelfontein, Cape Colony, (4 May 1902, on label). Collected by C. H. E. Seimund and C. H. B. Grant. Presented by Col. A. T.	,

Sloggett.

The female syntype is also in the collection.

Bull. Brit. Orn. Cl., 23: 19. Syntype, Adult male. Reg. no. 1904.6.22.302. (Banterbari, on label), Fernando Po, 31 Jan. 1904. Collected by E. Seimund. Presented by Mrs P. Sladen, the Duke of Bedford, and the Hon. W. Rothschild. This specimen was labelled as type by the author and is his selected male. His selected female syntype is also in the collection.
semibadius—Thryothorus semibadius Salvin, 1870
semicervinus—Basileuterus semicervinus P. L. Sclater, 1860 54 Proc. zool. Soc. Lond., 1860: 84. SYNTYPE, Adult female. Reg. no. 1884.5.456. Nanegal, Western Ecuador. Feb. 1859. Collected by L. Fraser. Sclater Collection. The measurements and date in the description refer to this specimen. The larger male syntype was collected in April.
semicinereus—Dysithamnus semicinereus P. L. Sclater, 1855 4 Proc. zool. Soc. Lond., 1855 : 90, pl. 97. SYNTYPE, Adult male. Reg. no. 1854.1.25.126. Bogota, Colombia. Purchased of S. Stevens. There are two other syntypes in the collection.
semicinereus—Hylophilus semicinereus P. L. Sclater and Salvin, 1867 56 Proc. zool. Soc. Lond., 1867: 570, pl. 30. HOLOTYPE, Adult. Reg. no. 1886.9.15.322. Para, lower Amazon, northern Brazil, May 1849. Collected by A. R. Wallace. Sclater Collection.
semiflava—Geothlypis semiflava Sclater, 1860
semiflavus—Tyrannulus semiflavus P. L. Sclater and Salvin, 1860 . Proc. zool. Soc. Lond., 1860 : 300. Syntype, Adult. Reg. no. 1888.1.1.607. Choctum, 1,200′, Northern Vera Paz, Guatamala. Collected by O. Salvin. Salvin-Godman Collection. There are several other syntypes in the collection.
semiflavus—Vireo semiflavus Salvin, 1863

- semifusca—Phyllomyias semifusca P. L. Sclater, 1862 9

 Proc. zool. Soc. Lond., 1861 (publ. 1862): 383, pl. 36. HOLOTYPE,

 Adult. Reg. no. 1888.1.13.436. Santa Martha, Colombia. Collected for Verreaux. Sclater Collection.
- semifuscus—Chlorospingus semifuscus P. L. Sclater and Salvin, 1873 51 Nomencl. Av. Neotrop.: 24, 157. HOLOTYPE, Adult. Reg. no. 1885.6.8.57. Near Quito, Ecuador. (Collected by Higgins, on label). Salvin-Godman Collection.
- semipagana—Elainea semipagana P. L. Sclater, 1862 9
 Proc. zool. Soc. Lond., 1861 (publ. 1862): 406. HOLOTYPE, Adult male. Reg. no. 1888.1.13.522. Babahoyo, Ecuador, Aug. 1859. Collected by L. Fraser. Sclater Collection.
- semiruber—Pericrocotus speciosus semiruber Whistler, 1933 . . . 20 J. Bombay nat. Hist. Soc., 36: 340. HOLOTYPE, Adult male. Reg. no. 1932.6.1.2. Sankarametta, 3,500', Vizagapatam, Upper East Ghats, India, 15 April 1930. Collected by V. S. La Personne and presented by the Bombay Natural History Society.
- **semirubra**—Rhipidura semirubra P. L. Sclater, 1877 37A Proc. zool. Soc. Lond., 1877 : 552. HOLOTYPE, Adult male. Reg. no. 1880.11.18.95. Admiralty Island, Pacific, March 1875. Collected by J. Murray on the voyage of H.M.S. Challenger.

There is another syntype at the Academy of Natural Sciences, Philadelphia.

- semitorquata—Garrulax semitorquata Ogilvie-Grant 1900 . . . 32B Bull. Br. Orn. Cl., 10: 49. Syntype, Adult male. Reg. no. 1900. 1.2.57. Five Fingers Mountains, Hainan, 4 April 1899. Collected by J. Whitehead. Purchased of E. Gerrard. The female syntype is also in the collection.

A second specimen in the collection may also be a syntype.

semperi—Leucopeza semperi P. L. Sclater, 1876
senegalensis—Textor senegalensis Shelley, 1896 61 The Birds of Africa, 1:34. HOLOTYPE, Adult. Reg. no. 1886.1. 30.6. Gambia River. Collected by and purchased of H. Whitely.
senex—Anthochaera senex G. R. Gray, 1860
senex—Platyrhynchus senex P. L. Sclater and Salvin, 1880 9 Proc. zool. Soc. Lond., 1880: 156. SYNTYPE, Adult. Reg. no. 1888.1.1.270. Sarayacu, Bobanaza River, East Ecuador. Collected by C. Buckley. Salvin-Godman Collection. The second syntype is also in the collection.
sepiaria—Brachypteryx sepiaria Horsfield, 1821
sepium—Orthotomus sepium Horsfield, 1821
septentrionalis—Chloropsis cyanopogon septentrionalis Robinson and Kloss, 1919
septentrionalis—Prionochilus maculatus septentrionalis Robinson and Kloss, 1921

serena—Erythrospiza serena P. L. Sclater, 1881 60 Ibis, (5) 4:544, pl. 15. SYNTYPE, Adult male. Reg. no. 1861. 6.15.2. Aneiteum, New Hebrides, 8 Feb. 1860. Collected by J. MacGillivray. Purchased of H. Cuming. The two immature syntypes are also in the collection.
serica—Pica serica Gould, 1845
sericea—Horeites sericea Walden, 1875
Sericea—Meliphaga sericea Gould, 1837
sericeogula—Raya sericeogula Hodgson, 1839
serinus—Criniger serinus J. and E. Verreaux, 1855 21 J. Orn., 1855: 105. SYNTYPE, Adult. Reg. no. 1854.5.23.24. Gaboon. Purchased of Verreaux. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 101.
serlei—Mirafra rufocinnamomea serlei White, 1960 17 Bull. Br. Orn. Cl., 80:11. HOLOTYPE, Adult male. Reg. no. 1955.59.133. Enugu, 1,000', Nigeria, 4 Oct. 1952. Collected and presented by W. Serle.
serva—Pyriglena serva P. L. Sclater, 1858
seshekeensis—Serinus atrogularis seshekeensis C. H. B. Grant and Mackworth-Praed, 1958

1955.42.65. (Collector's no. M56.) Chunga Pool, Sesheke District, south-western Northern Rhodesia, 10 May 1952. Collected by J. M. Winterbottom. Presented by the National Museums, Southern Rhodesia.
Setafer—Cinclosoma setafer Hodgson, 1836
chanensis—Certhia discolor shanensis E. C. S. Baker, 1930
hanensis—Tribura thoracica shanensis Ticehurst, 1941 35 Ibis, (14) 5: 318. HOLOTYPE, Adult male. Reg. no. 1948.80.1139. Maymyo, 3,500', Mandalay, Upper Burma, 20 Oct. 1937. Collected by P. F. Garthwaite. Presented by P. F. Garthwaite and H. C. Smith. The upper mandible is missing.
harii—Hypochera chalybeata sharii Bannerman, 1922 61 Bull. Brit. Orn. Cl., 43: 29. HOLOTYPE, Adult male. Reg. no. 1911.12.23.3308. Ratu, Gribingui River, French Equatorial Africa, 16 Sept. 1905. Collected by Boyd Alexander on the Alexander-Gosling Expedition. Boyd Alexander Bequest.

- sharpei—Cryptospiza sharpei Boyd Alexander, 1907 60
 Bull. Brit. Orn. Cl., 19:46. SYNTYPE, Adult male. Reg. no.
 1911.12.23.3297. Kaga Djirri, Kemo River, French Congo, 19
 Oct. 1905. Collected by Boyd Alexander. Boyd Alexander
 Bequest.

The bill of the specimen is imperfect. There are several other syntypes in the collection.

 sharpei—Macronyx sharpei Jackson, 1904
sharpei—Myzomela sharpei Ogilvie-Grant, 1888
sharpei—Phyllostrephus sharpei Shelley, 1880 21 Ibis, (4) 4:334. HOLOTYPE, Adult. Reg. no. 1895.5.1.744. Dares-Salaam. Collected by Dr. Kirk. Shelley Collection.
sharpei—Smithornis sharpei Boyd Alexander, 1903
sharpei—Trochalopterum sharpei Rippon, 1901
sharpii—Amadina sharpii Nicholson, 1878 60 Proc. zool. Soc. Lond., 1878: 130, pl. 10. HOLOTYPE, Adult. Reg. no. 1877.11.25.4. Abeokuta, Southern Nigeria. Collected and presented by H. Robin.
sharpii—Apalis sharpii Shelley, 1884
sharpii—Corvus sharpii Oates, 1889
Sharpii—Otyphantes sharpii Shelley, 1898
Sharpii—Pholidauges sharpii Jackson, 1898

shelleyi—Aethopyga shelleyi Sharpe, 1876
shelleyi—Amydrus morio shelleyi Hartert, 1891 62 Katalog der Vogelsammlung im Museum Senckenbergishen : 75, note. Syntype, Adult male. Reg. no. 1889.3.6.18. (Ugogo, East Africa. Collected by J. Kirk, on label.) Shelley Collection. There are several other syntypes in the collection.
shelleyi—Cinnyris shelleyi Boyd Alexander, 1899
shelleyi—Cotile shelleyi Sharpe, 1885
shelleyi—Cryptospiza shelleyi Sharpe, 1902
shelleyi—Hyphantornis shelleyi Sharpe, 1890 61 Cat. Bds. Brit. Mus., 13: 464. SYNTYPE, Adult male. Reg. no. 1860.12.31.98. Tete, Zambesi River. Collected by Sir J. Kirk. Presented by Earl Russell of the Foreign Office, from the Livingstone Expedition (vide, Sharpe in Hist. Coll. Brit. Mus., 2 (1906): 407). There are other syntypes in the collection.
shelleyi—Nesocharis shelleyi Boyd Alexander, 1903 60 Bull. Brit. Orn. Cl., 13:48. Holotype, Adult female. Reg. no. 1911.12.23.3300. Fernando Po, 12 Dec. 1902. (Pico Joaquin, on label), Moka, (10 Dec., on label). Collected by J. Lopez for Boyd Alexander. Boyd Alexander bequest.
shelleyi—Passer shelleyi Sharpe, 1891 65 Ibis, (6) 3: 256. HOLOTYPE, Adult (male, vide Shelley, The Birds of Africa, 3 (1902): 247). Reg. no. 1887.9.28.314. Lado, south Sudan (9 April 1884, vide Shelley, op. cit.). Collected and presented by Emin Pasha.
shelleyi—Saxicola shelleyi Sharpe, 1876

- shellevi—Spreo shellevi Sharpe, 1890 Cat. Bds. Brit. Mus., 13: 191. HOLOTYPE, Adult. Reg. no. 1895. 7.12.76. Somaliland. Collected by and purchased of E. Lort Phillips.
- sheppardi—Cinnyris sheppardi Jackson, 1910 Bull. Brit. Orn. Cl., 27:6. SYNTYPE, Adult male. Reg. no. 1910. 10.8.5. Kiunga, near Lamu, British East Africa, 31 July 1906. Collected for and presented by F. J. Jackson.

The described female syntype is also in the collection.

- sherriffi—Spelaeornis souliei sherriffi Kinnear, 1934 . 32B Bull. Br. Orn. Cl., 54:107. HOLOTYPE, Adult male. Reg. no. 1933.11.16.34. Dongda La, between Lingste and Trashi Yangsti. 10,500', east Bhutan, 25 July 1933. Collected by F. Ludlow and G. Sherriff and presented by the Salvin-Godman Trust Fund.
- 28.5. Aconquija, 4,000 metres, Tucuman, north-west Argentina, 7 Sept. 1918. Collected and presented by Stewart Shipton. The female syntype is also in the collection.
- shiwae—Cisticola tinniens shiwae C. M. N. White, 1947 Ostrich, 18: 174. HOLOTYPE, Adult male. Reg. no. 1931.12.21. 847. Lake Young, (5,000', on label), Shiwa Ngandu, Chinsali, Northern Rhodesia, 12 Oct. 1930. Collected and presented by Admiral H. Lynes.
- shortridgii—Zosterops shortridgii Ogilvie-Grant, 1909 . Ibis, (9) 3:663. HOLOTYPE, Adult male. Reg. no. 1905.12.26.85. Rabbit Island, Albany, south-western Australia, 17 April 1905. Collected by G. C. Shortridge. Presented by W. E. Balston.
- siamensis—Acridotheres siamensis Swinhoe, 1863. 62 Proc. zool. Soc. Lond., 1863: 303. HOLOTYPE, Adult. Reg. no. 1888.7.10.75. Siam. Collected by Sir R. Schomburgk. Seebohm Collection.
- siamensis—Graucalus macei siamensis E. C. S. Baker, 1918 20 Bull. Br. Orn. Cl., 38:69. SYNTYPE, Adult female. Reg. no. 1916. 12.27.590. Mi-Nam-Kabren, Siam, 5 Nov. 1915. Collected by C. Chunggat and presented by E. G. Herbert.

This is the author's selected female. His selected male syntype is also in the collection.

siamensis-Pomatorhinus olivaceus siamensis E. C. S. Baker, 1917 32B Bull. Br. Orn. Cl., 38:9. SYNTYPE, Adult male. Reg. no. 1916. 12.27.807. Maprit, south-west Siam, 27 Dec. 1915. Collected and presented by E. G. Herbert.

This is the author's selected male. His selected female syntype is

also in the collection.

siamensis—Zosterops siamensis Blyth, 1867
siberu—Calyptomena viridis siberu Robinson and Kloss, 1926 . 1 Ibis, (12) 2:285. HOLOTYPE, Adult male. Reg. no. 1947.60.72. Siberu Island, west Sumatra, 24 Sept. 1924. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore.
siberu—Dicrurus leucogenis siberu Chasen and Kloss, 1926 64 Ibis, (12) 2:294. HOLOTYPE, Adult male. Reg. no. 1947.60.24. Sireut Island, west Sumatra, 18 Sept. 1924. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore.
siberu—Oriolus chinensis siberu Chasen and Kloss, 1926 63 Ibis, (12) 2:294. HOLOTYPE, Adult male. Reg. no. 1947.60.29. Siberut Island, west Sumatra, 28 Sept. 1924. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore.
sibilans—Larvivora sibilans Swinhoe, 1863
sibilatrix—Phacellodomus sibilatrix P. L. Sclater, 1879
sibirica—Otocorys sibirica Swinhoe, 1871
sibutuense—Dicaeum sibutuense Sharpe, 1893
sicula—Acredula sicula Whitaker, 1901
signata—Leiothrix signata McClelland, 1840

- Birds in the Museum of the East India Company, 1 (1854): 289). Reg. no. 1880.1.1.893. Assam. Collected by J. McClelland. Presented by the Honourable East India Company Museum.
- signatum—Todirostrum signatum P. L. Sclater and Salvin, 1881 9

 Ibis, (4) 5:267. Syntype, Adult male. Reg. no. 1888.1.1.364.

 Yquitos, Upper Amazon, Peru, 26 Oct. 1878. Collected by H. Whitely. Salvin-Godman Collection.

 There is another syntype in the collection.
- signatus—Chlorospingus signatus Taczanowski and Berlepsch, 1885 51 Proc. zool. Soc. Lond., 1885: 82. SYNTYPE, Adult male. Reg. no. 1885.6.12.845. Machay, Ecuador, 3 Dec. 1883. Collected by J. Stolzmann. Sclater Collection.

 Listed by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 241, as one of the types.
- sikkimensis—Grammatoptila striata sikkimensis Ticehurst, 1924 32B Bull. Br. Orn. Cl., 44: 104. HOLOTYPE, Adult. Reg. no. 1886. 10.1.4037. Sikkim, April 1873. Collected by L. Mandelli. Hume Collection.
- sikkimensis—Irena puella sikkimensis Whistler and Kinnear, 1933. 22 J. Bombay nat. Hist. Soc., 36: 582. HOLOTYPE, Adult male. Reg. no. 1921.7.12.504. Sukna, Darjeeling, Feb. 1911. Collected and presented by H. K. Robinson.

similis—Iole maclellandi similis Rothschild, 1921
similis—Malaconotus similis Smith, 1836
similis—Pachyrhamphus similis Salvin, 1895
Renamed P . salvini Richmond, 1899 $(q.v.)$, on the grounds of preoccupation by P . similis Cherrie, 1891. There are three other syntypes in the collection.
similis—Petroica similis G. R. Gray, 1859
similis—Ptilotis similis Layard, 1876
simillima—Arachnothera simillima Hume, 1877
simillima—Pitta simillima Gould, 1868

*
simillima—Spizalauda simillima Hume, 1870
simillimus—Turdus simillimus Jerdon, 1839
simlaensis—Oreocincla mollissima simlaensis E. C. S. Baker, 1924. 31 The Fauna of British India; Birds (2nd ed.), 2:164. HOLOTYPE, Adult male. Reg. no. 1886.7.8.2317. Simla, Punjab, 5 Nov. 1878. Collected by A. O. Hume. Hume Collection.
simlaensis—Phylloscopus proregulus simlaensis Ticehurst, 1920 . 35 Bull. Brit. Orn. Cl., 41:55. HOLOTYPE, Adult. Reg. no. 1886. 7.8.1193. Simla, 2 Nov. 1880. Collected by W. Davison. Hume Collection.
simlaensis—Siva strigula simlaensis R. Meinertzhagen, 1926 32B Bull. Br. Orn. Cl., 46: 128. HOLOTYPE, Adult male. Reg. no. 1886.10.1.6739. Simla, northern India, 21 Oct. 1880. Collected by W. Davison. Hume Collection.
simlaensis—Sylviparus modestus simlaensis E. C. S. Baker, 1917 . 42 Bull. Brit. Orn. Cl., 38:8. HOLOTYPE, Adult. Reg. no. 1886. 11.1.473. Simla, north Punjab, India. 22 Oct. 1880. Collected by W. Davison (not A. L. Davidson as stated in description). Hume Collection.
simonsi—Buarremon simonsi Sharpe, 1900
simonsi—Heleodytes simonsi Chubb, 1906
simonsi—Scytalopus simonsi Chubb, 1917
simplex—Criniger simplex Wallace, 1862

- Moluccas, 1858. Collected by Allen. A. R. Wallace Collection. Listed as type by Sharpe, *loc. cit*. There are five other syntypes in the collection.
- simplex—Grallaria simplex Salvin and Godman, 1884 4

 Ibis, (5) 2:451. Syntype, Adult male. Reg. no. 1889.7.10.899.

 5,000', Roraima, British Guiana, 14 Sept. 1883. Collected by H. Whitely. Salvin-Godman Collection.

There are several other syntypes in the collection.

Proc. zool. Soc. Lond., 1874: 132. SYNTYPE, Adult male. Reg. no. 1885.2.10.212. Lima. Collected by W. Nation. Sclater Collection.

Taczanowski used a Jelski MS. name. There is another syntype in the collection. This specimen and the other British Museum syntype do not form part of the main series, which was at Warsaw, but are a part of the syntypical material referred to in the text.

There is a female syntype in the collection.

- sindianus—Crateropus terricolor sindianus Ticehurst, 1920 . . . 32B Bull. Br. Orn. Cl., 40:156. HOLOTYPE, Adult male. Reg. no. 1920.10.23.1. Karachi, India, 2 April 1919 (27 April 1919, on label). Collected and presented by C. B. Ticehurst.
- sindianus—Mirafra erythroptera sindianus Ticehurst, 1920 . . . 17

 Bull. Br. Orn. Cl., 41:56. HOLOTYPE, Adult male. Reg. no. 1926.10.20.1. Karachi, Sind, 15 April 1918. Collected and presented by C. B. Ticehurst.

sindianus—Phylloscopus sindianus Brooks, 1880
sindianus—Prinia flaviventris sindianus Ticehurst, 1920 35 Bull. Brit. Orn. Cl., 40:157. HOLOTYPE, Adult male. Reg. no. 1886.10.1.1568. Sukker, Sind, 20 Feb. 1879. Collected by E. Butler. Hume Collection.
sinensis—Alauda sinensis Waterhouse, 1839
sinensis—Brachypteryx sinensis Rickett and La Touche, 1897 31 Bull. Br. Orn. Cl., 6:50. SYNTYPE, Adult male. Reg. no. 1897. 12.8.28. Kuatun, north-west Fohkien, Oct. 1896. Collected and presented by C. B. Rickett and J. D. La Touche. The female syntype is also in the collection.
sinensis—Corvus sinensis Moore, 1856
sinensis—Cryptolopha sinensis Rickett, 1898
sinensis—Enicurus sinensis Gould, 1865
sinensis—Lusciniola pryeri sinensis Witherby, 1912 35 Bull. Brit. Orn. Cl., 31:11. Holotype, Adult male. Reg. no. 1924.12.20.7. Hankow, China, 20 March 1912. Collected by Admiral H. Lynes. Presented by H. F. Witherby.
sinensis—Monticola rufiventris sinensis A. C. Meinertzhagen, 1927 31 Bull. Br. Orn. Cl., 47: 148. HOLOTYPE, Adult female. Reg. no. 1905.12.24.349. Kuatun, north-west Fokien, China, 21 May 1897. Collected and presented by C. B. Rickett.

singapurensis—Aegithina tiphia singapurensis Chasen and Kloss,
Bull. Raffles Mus., 5:85. HOLOTYPE, Adult male. Reg. no. 1947.60.44. Pulau Ubin, Straits of Johore, near Singapore, 26 Feb. 1923. Collected by P. M. de Fontaine and presented by the Raffles Museum, Singapore.
singetra—Tchitrea mutata singetra Salomonsen, 1933 38 Bull. Brit. Orn. Cl., 53: 120,122,124. HOLOTYPE, Adult male. Reg. no. 1931.8.18.2449. Soalala, Western Madagascar, 31 May (or July, the label is indistinct) 1929. (Collected by R. Decary, on label). From the Franco-Anglo-American Archbold-Vernay Expedition.
sinlumensis—Urocichla sinlumensis Harington, 1908 321 Ann. Mag. nat. Hist., (8) 2:246. SYNTYPE, Adult male. Reg. no. 1908.8.2.23. Sinlum-kaba, Bhamo District, Upper Burma, 29 April 1908. Collected and presented by H. H. Harington. The second syntype is also in the collection.
sintaungensis—Krimnochelidon concolor sintaungensis E. C. S. Baker, 1933
sinuatus—Cardinalis sinuatus Bonaparte, 1838
sipahi—Corythus? sipahi Hodgson, 1836
sipora—Oriolus chinensis sipora Chasen and Kloss, 1926 63 Ibis, (12) 2:294. HOLOTYPE, Adult male. Reg. no. 1947.60.28. Sipora Island, West Sumatra, 5 Nov. 1924. Collected by C. B. Kloss. Presented by the Raffles Museum, Singapore.
siquijorensis—Iole siquijorensis Steere, 1890
sladeni—Turdus poliocephalus sladeni Cain and Galbraith, 1955. 31 Bull. Br. Orn. Cl., 75:92. HOLOTYPE, Adult male. Reg. no. 1959.21.553. Turipava, 4,100', Guadalcanal, Solomon Islands,

- 18 Aug. 1953. Collected by a native hunter for the Oxford University Expedition to the Solomon Islands.
- sloggetti—Fringillaria impetuani sloggetti Macdonald, 1957.
 Contributions to the ornithology of Western South Africa: 170.
 HOLTYPE, Adult male. Reg. no. 1901.9.5.47. Deelfontein, Cape Colony, 25 Jan. 1901. Collected by C. H. E. Seimund. Presented by Col. A. T. Sloggett.
- smithersi—Ortygospiza atricollis smithersi Benson, 1955 . . . 60 Bull. Brit. Orn. Cl., 75:106. Holotype, Adult male. Reg. no. 1955.41.1. Mpasa, Bangweulu 11°0′ S., 30°30′ E., north Northern Rhodesia, 19 Aug. 1953. Collected and presented by C. W. Benson.

There is a female syntype in the collection; four other syntypes are at the Academy of Natural Sciences, Philadelphia.

The male syntype is in the Academy of Natural Sciences, Philadelphia, vide Schauensee, Proc. Acad. nat. Sci. Philadl., 109 (1957): 197.

smithii—Muscipeta smithii Fraser, 1843	}
smithii—Ptilonorhynchus smithii Vigors and Horsfield, 1827 69 Trans. Linn. Soc. Lond., 15:264. Holotype, Adult. Reg. no. 1863.7.7.52. Watham River, New South Wales, Nov. 1804. Collected by R. Brown. Presented by the Linnean Society.	•
smithresi—Cisticola chiniana smithresi Hall, 1956 35 A mis-spelling, see smithersi. The subspecific name is also quoted correctly in the description.	5
sobatensis—Mirafra sobatensis Lynes, 1914	7
socialis—Prinia socialis Sykes, 1832	3
socotrana—Fringillaria socotrana Ogilvie-Grant and Forbes, 1899 49. Bull. Liverpool Museum, 12:2. SYNTYPE, Adult male. Reg. no. 1899.8.11.57. Adho Dimellus, Sokotra (4,500', 6 Feb. 1899., on label). Collected by W. R. Ogilvie-Grant and H. O. Forbes. Presented by the Royal Society. The female syntype is also in the collection.	•
socotranus—Rhynchostruthus socotranus P. L. Sclater and Hartlaub, 1881	9
There is another syntype in the collection. solitaria—Agriornis solitaria P. L. Sclater, 1859	9

solitaris—Muscicapa solitaris Müller, 1835
solstitialis—Troglodytes solstitialis P. L. Sclater, 1858 28 Proc. zool. Soc. Lond., 1858: 550. SYNTYPE, Adult male. Reg. no. 1886.9.15.193. Matos, near Riobamba, Ecuador, June 1858. Collected by L. Fraser. Sclater Collection. The second syntype is also in the collection.
somalica—Alaudula somalica Sharpe, 1895
somalica—Certhilauda somalica Witherby, 1903 17 Bull. Br. Orn. Cl., 14:29. HOLOTYPE, Adult male. Reg. no. 1908.5.28.111. Dibbit, about 6°50′ N., 47°30′ E., Somaliland, 11 March 1903. Collected by and purchased of A. E. Hamerton.
somalica—Cisticola somalica Sharpe, 1895
somalica—Saxicola somalica Sharpe, 1895
somaliensis—Muscicapa somaliensis Bannerman, 1909 37 Bull. Brit. Orn. Cl., 25: 20. SYNTYPE, Adult female. Reg. no. 1905. 11.27.241. Waghar Mountains, North Somaliland, (27 June 1905, on label). Collected by G. W. Bury. Purchased of E. Gerrard. The described male syntype is also in the collection. Both were labelled "type" by the author. The label of the above specimen gives the collector's data on bill, feet, etc., which also appears in the description.
Somaliensis—Parisoma blanfordi somaliensis W. L. Sclater and Mackworth-Praed, 1918

somervillei—Timalia somervillei Sykes, 1832 32B Proc. zool. Soc. Lond. 1832: 88. HOLOTYPE, Adult (relaxed mount). Reg. no. 1860.4.16.42a. Ghauts, Deccan, India. Collected by W. Sykes and presented by the Honourable East India Company Museum.
sondaica—Nitidula hodgsoni sondaica Robinson and Kloss, 1923 37 J. Fed. Malay Sts. Mus., 11:54. Holotype, Adult male. Reg. no. 1936.4.12.2134. Korinchi Peak, 7,300', West Sumatra, 10 May 1914. Collected by H. C. Robinson and C. B. Kloss. Presented by the Raffles Museum, Singapore.
songarus—Parus songarus Severtsoff, 1873
songeaensis—Apalis melanocephala songeaensis C. H. B. Grant and Mackworth-Praed, 1947
sonoroides—Ptilotis sonotoides G. R. Gray, 1862
sordida—Emberiza sordida Blyth, 1845
J. Asiat. Soc. Bengal., 13:958. HOLOTYPE, Adult female. Reg. no. 1845.1.9.333. Nepal. Collected and presented by B. H. Hodgson. Although Sharpe, Cat. Bds. Brit. Mus., 12 (1888):524, lists four "types", the description was taken from a single "presumed female" and Sharpe's other specimens are males.

- sordida". Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 267, lists three syntypes which are also in the collection. sordida—Glaucomvias sordida Walden, 1870 Ann. Mag. nat. Hist., (4) 5:218. SYNTYPE, Adult male. Reg. no. 1887,12,30,1209. (Rangwellitenuek, on label), Ceylon, 20 Jan. 1866. Collected by S. Chapman. Tweeddale Collection. The author refers to four syntypes, one other of which is also in the collection. sordida—Herpornis xantholeuca sordida Robinson & Kloss, 1919 . 32B Ibis, (11): 588. SYNTYPE, Adult male, Reg. no. 1919.12.20.326. Daban, 650', south Annam, 21 March 1918. Collected and presented by C. B. Kloss. This is the authors' selected male. Their selected female syntype is also in the collection. sordida—Siva sordida Hume, 1877 . 32B Stray Feathers, 5:104. HOLOTYPE, Adult male. Reg. no. 1886. 10.1.6842. Moolyit, Tenasserim, Burma, 23 Feb. 1877. Collected by W. Davison. Hume Collection. sordidior—Moupinia poecilotis sordidior Rothschild, 1921 Novit. Zool., 28:36. HOLOTYPE, Adult. Reg. no. 1921.7.15.249. Lichiang Range, Yunnan, 1918. Collected by G. Forrest and presented by Stephenson Clarke. sordidior—Proparus sordidior Rippon, 1903. Bull. Br. Orn. Cl., 13:60. HOLOTYPE, Adult. Reg. no. 1902. 8.8.180. Gyi-dzin-shān, 9,100', east of Talifu, west Yunnan, 2 April 1902. Collected and presented by G. Rippon. no. 1888.3.18.15. Batang Padang Mountains, Perak, Malaya. Collected and presented by L. Wray. The second syntype is also in the collection. sordidior—Yuhina gularis sordidior Kinnear, 1925. Bull. Br. Orn. Cl., 45: 74. HOLOTYPE, Adult male. Reg. no. 1924. 12.21.159. Ngoi-Tio, Tonkin, 24 June 1924. Collected by H. Stevens and presented by the Percy Sladen Trust and Salvin-Godman Exploration Fund. sordidulus—Contopus sordidulus P. L. Sclater, 1859

Proc. zool. Soc. Lond., 1859: 43. HOLOTYPE, Adult. Reg. no. 1888.1.13.1024. Orizaba, Mexico (vide, Ibis, (1) 1 (1859): 441,

sordidus—Cinclus sordidus Gould, 1859
sordidus—Turdus sordidus Latham, 1801
sordidus—Xanthiscus flavescens sordidus Robinson & Kloss, 1919 21 Ibis, (11) 1:569. SYNTYPE, Adult male. Reg. no. 1919.12.20.239. Arbre Broyé, 5,400', Langbian Mountains, south Annam, 8 May 1918. Collected by C. Boden Kloss and presented by the latter and H. C. Robinson. This is the authors' selected male. Their selected female syntype is also in the collection.
sorghophila—Calamodyta sorghophila Swinhoe, 1863 35 Proc. zool. Soc. Lond., 1863 : 92. HOLOTYPE, Adult. Reg. no. 1898.9.1.1079. Amoy, south-east China, 20 May 1861. Collected by R. Swinhoe. Seebohm Collection.
Soror—Minla soror Sharpe, 1887
soror—Pitta (Hydrornis) soror Wardlaw-Ramsay, 1881
sororia—Lochmias sororia P. L. Sclater & Salvin, 1873 3 Proc. zool. Soc. Lond., 1873: 511. Holotype, Adult. Reg. no. 1889.5.20.84. Venezuela, 1873. Collected by J. M. Spence. Sclater Collection.
soudanensis—Zonogastris soudanensis Sharpe, 1890 60 Cat. Bds. Brit. Mus., 13: 298. SYNTYPE, Adult male. Reg. no. 1877.7.11.211. Khartoum, (Chartum, on label; W. L. Sclater, Systema Avium Aethiopicarum, (1930): 788, gives the locality as "probably from the Upper Nile Valley"; and recent examinations show that the specimen matches birds from Mongalla, Upper Nile Valley, rather than those from further north). Obtained through L. Schaufuss of Dresden. Sharpe Collection. This is the described specimen and is labelled as the type by the author. There are three other syntypes in the collection.

- sousae—Apalis moreaui sousae Benson, 1945. 35

 Bull. Brit. Orn. Cl., 66: 19. Holotype, Adult male. Reg. no. 1945.25.3. Njesi Plateau, 10 miles north of Unangu, northern Portuguese East Africa, (12°45′ S., 35°20′ E., on label), 12 Aug. 1945. Collected by Jali Makawa for C. W. Benson. Presented by C. W. Benson.
- spaldingii—Orthonyx spaldingii Ramsay, 1868 32A Sydney Morning Herald, Mar. 21.1968 : 4, and Proc. zool. Soc. Lond., 1868 : 386. Syntype, Adult female. Reg. no. 1881.5.1. 4236. Dense brush about 35 miles inland from Rockingham Bay, Queensland, Oct. 1867. Collected by E. Spalding. Gould Collection.

The male syntype is also in the collection.

- speciosa—Dasycrotapha speciosa Tweeddale, 1878 32A Proc. zool. Soc. Lond., 1878: 114, pl. 9. HOLOTYPE, Adult male. Reg. no. 1888.4.20.1090. Valencia, Negros, Philippines, Aug. 1877. Collected by A. Everett. Tweeddale Collection.
- speciosa—Formicivora speciosa Salvin, 1876.

 Ibis, (3) 6: 494. Syntype, Adult male. Reg. no. 1889.7.10.469.

 Island of Puna, Ecuador. Collected by Villagomez for C.

 Buckley. Salvin-Godman Collection.

The female syntype is also in the collection.

See Horsfield, Zoological Researches in Java, 1824: 241–2. There is a second syntype in the collection.

(See also Hellmayr and Conover, Catalogue of birds of the Americas, 11 (1938): 437.)

spectabilis—Donacicola spectabilis P. L. Sclater, 1879 60 Proc. zool. Soc. Lond., 1879 : 449, pl. 37. HOLOTYPE, Adult. Reg. no. 1881.3.29.39. New Britain. Collected by the Rev. G. Brown. Purchased of E. Gerrard.
spectabilis—Saxicola spectabilis Hartlaub, 1865
Spectatrix—Cercomela spectatrix Clarke, 1919
specularis—Buarremon specularis Taczanowski, 1879
speculigera—Schistochlamys speculigera Gould, 1855 51 Proc. zool. Soc. Lond., 1855 : 69. HOLOTYPE, Adult. Reg. no. 1855.11.5.3. Ucayali River, Eastern Peru. (Aug. 1852. Collected by Hauxwell, vide Proc. zool. Soc. Lond., 1856 : 69). Purchased of Gould.
spekeoides—Ploceus spekeoides C. H. B. Grant and Mackworth-Praed, 1947
spencei—Turdus simillimus spencei Whistler and Kinnear, 1933 . 31 J. Bombay nat. Hist. Soc., 36:77. HOLOTYPE, Adult male. Reg. no. 1932.6.1.1. Vizaga Patam, 3,000', Jeypore Agency, Eastern Ghats, India, 19 April 1930. Collected and presented by V. S. La Personne of the Vernay Scientific Survey of the Eastern Ghats.
spilodera—Entomophila? spilodera G. R. Gray, 1859 36 Proc. zool. Soc. Lond., 1859: 155. HOLOTYPE, Adult (female, on label). Reg. no. 1873.5.12.215. North-west New Guinea (Dorey, 1858, on label). Collected by A. R. Wallace. Wallace Collection.
spilodera—Rhipidura spilodera G. R. Gray, 1870

Received and registered as the type. Tristram, Catalogue of a collection of birds. (1889): 206, claimed to have the types at that

date, but this syntype came to Seebohm later.

spodionota—Myrmotherula spodionota P. L. Sclater and Salvin, 1880 4
Proc. zool. Soc. Lond., 1880: 159. HOLOTYPE or SYNTYPE, Adult
male. Reg. no. 1889.7.10.351. Sarayacu, Ecuador. Collected by
C. Buckley. Salvin-Godman Collection.

There is another Buckley specimen from the same collection in the museum, also from Sarayacu, though the authors referred to a

"specimen unicum!"

spodionotus—Buarremon spodionotus P. L. Sclater and Salvin, 1879 49
Ibis, (4) 3: 425. Syntype, Adult male. Reg. no. 1885.6.12.927.
Guapula, West Ecuador, May 1859. Collected by L. Fraser. Sclater Collection.

Listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 264. The other syntypes are in the collection.

spodionotus—Dysithamnus spodionotus Salvin and Godman, 1883. 4

Ibis, (5) 1:211. Syntype, Adult male. Reg. no. 1889.7.10.263.

Roraima, British Guiana, 5 Jan. 1882. Collected by H. Whitely.

Salvin-Godman Collection.

There is another syntype in the collection.

spodioptila—Terenura spodioptila P. L. Sclater and Salvin, 1881 . 4

Ibis, (4) 5: 270, pl. 9. HOLOTYPE, Adult male. Rcg. no. 1889.

7.10.511. Bartica Grove, British Guiana, 7 June 1880. Collected by H. Whitely. Salvin-Godman Collection.

- spodiostethus—Attila spodiostethus Salvin and Godman, 1883. 7

 Ibis, (5) 1:209. Syntype, Adult male. Reg. no. 1888.1.20.644.
 Bartica Grove, British Guiana, (25 May 1880, on label). Collected by H. Whitely. Salvin-Godman Collection.

 The female syntype is also in the collection.
- spodiuros—Pachyrhamphus spodiuros P. L. Sclater, 1860 . . . 7

 Proc. zool. Soc. Lond., 1860: 279. SYNTYPE, Adult male. Reg.
 no. 1888.1.13.1469. Babahoyo, west Ecuador, July 1859. Collected by L. Fraser. Sclater Collection.

 There are two other syntypes in the collection.

- squamatum—Malacopteron squamatum Eyton, 1845 . . . 32B Ann. Mag. nat. Hist., (1) 16:228. SYNTYPE, Adult. Reg. no. 1846.3.4.33. Malacca. Collected and presented by A. Charlton. There is a second syntype in the collection.
- squamicollis—Uroloncha squamicollis Sharpe, 1890 60

 Cat. Bds. Brit. Mus., 13:359. SYNTYPE, Adult male. Reg. no. 1889.3.15.64. (Chungchow, on label), Szechuen, China, 4 May 1869. Collected by R. Swinhoe. Seebohm Collection.

 This is labelled as the described specimen. There are many other syntypes in the collection.
- squamifrons—Estrelda squamifrons Smith, 1836 6
 Report of the Expedition for exploring Central Africa of 1834: 49.
 SYNTYPE, Adult. Reg. no. 1872.10.4.92. South Africa (Latakoo, vide Smith, Illustrations of the zoology of South Africa, Aves (1844): pl. 95). Collected and bequeathed by A. Smith.

The use of the generic name "Estrelda" on the specimen label, in place of "Amadina" used in the Illustration of the Zoology of South Africa indicates that this is an early specimen. Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 408, listed it as the type. There is a female syntype in the Academy of Natural Sciences of Philadelphia.

- squamulatus—Crateropus squamulatus Shelley, 1884 32B Ibis, (5) 2:45. HOLOTYPE, Adult male. Reg. no. 1895.5.1.1328. Mombasa, East Africa, 23 July 1877. Collected by Dr. Fischer. Shelley Collection.
- squamulatus—Microcerculus squamulatus P. L. Sclater and Salvin, 1875 28 Proc. zool. Soc. Lond., 1875: 37, pl. 6. HOLOTYPE, Adult male. Reg. no. 1886.9.15.76. San Cristobal, Tachira Province, Venezuela. Collected by A. Goering. Sclater Collection.

Listed as the type by Mees, op. cit., p. 75; and labelled and registered as the type, but four specimens are referred to in the text. The other three syntypes are also in the collection.

- stanfordi—Alcippe nepalensis stanfordi Ticehurst, 1930 32B Bull. Br. Orn. Cl., 50:84. Holotype, Adult male. Reg. no. 1932.7.13.64. Taungup-Prome road, 2,900', Arakan Yoma, Burma, 20 March 1930. Collected and presented by J. K. Stanford.
- stanfordi—Pericrocotus stanfordi Vaughan and Jones, 1913 20 Ibis, (10) 1:383. Syntype, Adult male. Reg. no. 1910.5.2.264. Samshui, 110 miles from the sea, south-east China, 17 April 1906. Collected by and received in exchange from R. E. Vaughan and K. H. Jones.

There are fourteen other syntypes in the collection.

- stanfordi—Pomatorhimus ferruginosus stanfordi Ticehurst, 1935 . 32B Bull. Br. Orn. Cl., 55:178. HOLOTYPE, Adult male. Reg. no. 1936.7.14.1. Near Hkasang, Sadon-Sima road, 6,000', east of Myitkyina, Upper Burma, 24 Dec. 1934. Collected and presented by J. K. Stanford.

Although Shelley referred to the "type described" (i.e. the specimen selected above), he used two birds for his description:

this one for the plumage details and an Ayres specimen from Potchefstroom for the soft part colouration. This and a third syntype are in the collection.

- stejnegeri—Chrysomitris stejnegeri Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:210. SYNTYPE, Adult male. Reg. no. 1885.2.10.696. Sorata, Bolivia. Collected by C. Buckley. Sclater Collection.

This specimen was labelled as the type by the author. There are several other syntypes in the collection.

stejnegeri—Hemignathus stejnegeri S. B. Wilson, 1889 59

Ann. Mag. nat. Hist., (6) 4 (Nov. 1889): 400. Syntype, Adult male. Reg. no. 1898.10.20.1. Kauai, Sandwich Island, 1888. Collected and presented by S. B. Wilson.

Labelled "n. sp." by the author. There are two other syntypes in the University Museum of Zoology, Cambridge. This is a synonym of *H. procerus* Cabanis, 1889, but Peters, *Birds of the World*, 14 (1868): 97 does not refer to the name.

Labelled "n. sp." by the author. There are several other syntypes in the collection and another two in the University Museum of Zoology, Cambridge.

The other specimen from the original syntypical series is also in the collection.

- stellata—Hypocnemis stellata P. L. Sclater and Salvin, 1880 . . . 4

 Proc. zool. Soc. Lond., 1880: 160. SYNTYPE, Adult. Reg. no.
 1889.7.10.721. Sarayacu, Ecuador. Collected by C. Buckley.
 Salvin-Godman Collection.
 - The other syntype is in the collection.
- stellata—Margarornis stellata P. L. Sclater and Salvin, 1873 . . . 3
 Nomencl. Av. Neotrop.: 160. HOLOTYPE, Adult. Reg. no. 1889.5.
 14.530. Vicinity of Quito, Ecuador. Collected by Higgins.
 Salvin-Godman Collection.
- stellatus—Brachypteryx (Drymochares) stellatus Gould, 1868. . . 31

 Proc. zool. Soc. Lond., 1868: 218. SYNTYPE, Adult. Reg. no. 1881.5.1.308. Nepal, 10,000', (Sikkim, vide Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 31). Collected by C. V. Eccles. Gould Collection.

Listed as type by Sharpe, *loc. cit*. The second syntype is also in the collection.

- stenorhynchus—Pomatorhinus stenorhynchus Godwin-Austen, 1877 32B J. Asiat. Soc. Bengal, 46, pt. 2:43. Syntype, Adult male. Reg. no. 1888.4.20.727. Manbum Tila, 800', Tenga Pani River, near Saddya, Upper Assam, 21 Jan. 1877. Collected by M. J. Ogle. Tweeddale Collection; formerly in the Godwin-Austen Collection. There are several female syntypes in the collection.
- stephanophorus—Hyterhyphantes stephanophorus Sharpe, 1891 . 61 Ibis, (6) 3:117, pl. 6. HOLOTYPE, Adult male. Reg. no. 1893. 12.1.5. Mau, Uganda, 14 Sept. 1889. Collected and presented by F. J. Jackson.
- stevensi—Acrocephalus concinens stevensi E. C. S. Baker, 1922 . 35

 Bull. Brit. Orn. Cl., 43:16. HOLOTYPE, Adult male. Reg. no.
 1922.8.9.2. Hessamara, North Lakhimpur, Eastern Assam, 11

 April 1905. Collected and presented by H. Stevens.
- stevensi—Schoeniparus rufogularis stevensi Kinnear, 1924 . . . 32B Bull. Br. Orn. Cl., 45:10. Holotype, Adult male. Reg. no. 1924.12.21.165. Bao Ha, Tonkin, 22 Oct. 1923. Collected by H. Stevens and presented by the Percy Sladen Trust and Salvin Godman Exploration Fund.
- **stevensi**—Turdinulus brevicaudatus stevensi Kinnear, 1925 . . . **32B** Bull. Br. Orn. Cl., 45 : 74. HOLOTYPE, Adult male. Reg. no. 1924. 12.21.147. Ngoi-Tio, Tonkin, 26 June 1924. Collected by H.

Stevens and presented by the Percy Sladen Trust and Salvin-Godman Exploration Fund.
stewarti—Pseudosicalis stewarti Chubb, 1923 59 El Hornero, 3: 34. Pl. 1. Syntype, Adult male. Reg. no. 1922. 9.28.1. Gualfin, 2,200 metres, Datamarca, north-west Argentine, 29 Sept. 1917. Collected and presented by S. Shipton. This is the author's selected male syntype, the described female syntype is also in the collection.
stictigula—Alcippe stictigula Shelley, 1903
stictilaema—Crateropus stictilaema Boyd Alexander, 1901 32B Bull. Br. Orn. Cl., 12:10. SYNTYPE, Adult male. Reg. no. 1911. 12.23.1813. Volta River, Gold Coast, 6 April 1901. Collected and bequeathed by Boyd Alexander. There are several other syntypes in the collection.
stictoptera—Elainia stictoptera P. L. Sclater, 1859
stictoptera—Rhopoterpe stictoptera Salvin, 1893
stictothorax—Synallaxis stictothorax P. L. Sclater, 1859
sticturus—Herpsilochmus sticturus Salvin, 1885 4 Ibis, (5) 3:424. Syntype, Adult male. Reg. no. 1889.7.10.435. Bartica Grove, British Guiana, 16 Oct. 1879. Collected by H. Whitely. Salvin-Godman Collection. There are several other syntypes in the collection.
stigmatops—Buchanga stigmatops Sharpe, 1879
stochri—Cisticola stochri Neave, 1907

- Adult male. Reg. no. 1907.12.29.91. Luangwa Valley, Mterize River, north-east Rhodesia, 24 July 1905. Collected by S. A. Neave. Presented by the British South Africa Company. The described female syntype is also in the collection.

The male syntype is thought to be at Calcutta but was not listed among the other types there by Sclater, *Ibis*, (6) 4 (1892): 78.

This specimen appears to be the syntype referred to in the description (op. cit., 518).

The earlier label is inscribed "Type".

- stolidus—Serilophus hunatus stolidus Robinson and Kloss, 1919 . Bull. Br. Orn. Cl., 40: 16. Syntype, Adult male. Reg. no. 1936.

4.12.3151. Kao Nawng, Bandon, Siam, 19 May (June, on label) 1913. Collected by H. C. Robinson and E. Seimund and presented by the Raffles Museum, Singapore. There is a female syntype in the collection.
stolzmanni—Urothraupis stolzmanni Taczanowski and Berlepsch, 1885
stonii—Aeluroedus stonii Sharpe, 1881 69 Nature, 1876: 339. Holotype, Adult. Reg. no. 1877.2.9.4. Laloki River, south-east New Guinea. Collected by and purchased of O. C. Stone.
storeyi—Chloropeta storeyi Ogilvie-Grant, 1906 35 Bull. Brit. Orn. Cl., 19:32. HOLOTYPE, Adult. Reg. no. 1904. 4.14.15. (Nairobi River, 5,450', on label). British East Africa, 9 Oct. 1903. Collected and presented by C. B. Storey. See also Sclater, Systema Avium Æthiopicarum, (1930):414, for type locality and discussion of specimen.
Stracheyi—Emberiza stracheyi Moore, 1856
straminea—Locustella straminea Seebohm, 1881
strangei—Drymoica strangei Fraser, 1843
strenua—Geospiza strenua Gould, 1837

- strenuipes—Turdinus strenuipes Bannerman, 1920 . . . 32B Bull. Br. Orn. Cl., 41:5. HOLOTYPE, Adult male. Reg. no. 1920. 3.10.1. Iju Waterworks, near Lagos, south Nigeria, 15 Feb. 1920. Collected by W. P. Lowe and presented by Capt. Dane. R.N.

- streptophora—Rhipidura streptophora Ogilvie-Grant, 1911 . . . 37A Bull. Brit. Orn. Cl., 29:25. SYNTYPE, Adult male. Reg. no. 1911.12.20.1308. Mouth of the Mimika River, Dutch New Guinea, 18 March 1911. Collected by C. H. B. Grant. Presented by the subscribers to the British Ornithologists' Union Expedition to Dutch New Guinea.

This is the described male. The described female syntype is also in the collection.

- stresemanni—Mirafra stresemanni Bannerman, 1923 . . . 1'
 Bull. Br. Orn. Cl., 43:164. HOLOTYPE, Adult male. Reg. no. 1923.10.26.346. 15 miles north of Ngaundere, 3,700', Adamawa, north Cameroon, 3 Nov. 1922. Collected by and purchased of G. L. Bates.

This specimen was registered as the type and was figured in the description. The second syntype was sent in exchange to the American Museum of Natural History in 1921.

striata—Pachycephala striata Vigors and Horsfield, 1827 . . . 39

Trans. Linn. Soc. Lond., 15: 240. SYNTYPE. Adult (female, on label) (relaxed mount). Reg. no. 1863.7.7.39. Australia (=New South Wales, vide Mathews, Birds of Australia, 8 (1920): Collected by G. Caley. Presented by the Linnean Society.

Listed as the type by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 209. Two specimens were described but only this one came to the museum.

striata—Prinia striata Swinhoe, 1859
striata—Sittella striata Gould, 1869
striaticeps—Automolus striaticeps P. L. Sclater and Salvin, 1875. Proc. zool. Soc. Lond., 1875: 37. SYNTYPE, Adult. Reg. no. 1889.5.20.325. Bogota, Colombia, 1872. Sclater Collection; obtained by Sclater from Ward (a dealer). Automolus striaticeps Taczanowski, 1874, is a nomen nudum. Taczanowski used Sclater's MS. name. This bill is imperfect.
striaticeps—Hapalocercus striaticeps Salvin, 1897
striaticollis—Anabates striaticollis P. L. Sclater, 1857
striaticollis—Campylorhynchus striaticollis P. L. Sclater, 1857 . 28 Proc. zool. Soc. Lond., 1857: 272. HOLOTYPE, Adult. Reg. no. 1886.9.15.48. Bogota, Colombia. Sclater Collection; obtained by Sclater from Jamrach, a dealer.
striatigula—Neomixis striatigula Sharpe, 1881 321 Proc. zool. Soc. Lond., 1881 : 195, pl. 19. Holotype, Adult. Reg. no. 1880.5.1.23. Fianarantosa, Madagascar. Collected by the Rev. D. Cowan. Purchased of J. E. Harting to whom Cowan sent his Madagascar collection (vide Sharpe, in Hist. Coll. Brit. Mus. 2 (1906) : 381).
striatipectus—Poliospiza striatipectus Sharpe, 1891 59 Ibis, (6) 3:258. Syntype, Adult female. Reg. no. 1893.12.1.10. Elgeyo, Kenya, 8 July 1890. Collected and presented by F. J. Jackson. (cont.)

(cont.)

Both male and female syntypes were described but the male does not appear to be in the collection. Ann. Mus. Civ. Genoa. (2) 14:150. SYNTYPE, Adult male. Reg. no. 1897.8.29.18. Moroka, Owen Stanley Range, New Guinea, 26 July 1893. Collected by L. Loria. Purchased from the Genoa Museum as a syntype. striatus—Agriornis striatus Gould, 1839 Adult female (=male, vide Hellmayr, Catalogue of birds of the Americas, 5 (1927): 3). Reg. no. 1881.2.18.128. Santa Cruz, Patagonia, April. Collected by C. Darwin. Eyton Collection, received by him from the museum of the Zoological Society of London. striatus—Dicrurus striatus Tweeddale, 1877. Proc. zool. Soc. Lond., 1877: 545. SYNTYPE, Adult male. Reg. no. 1880.11.18.41. Pasananca, 7 miles from Zamboanga, south Mindanao, Philippines, Feb. 1875. Collected by J. Murray. Received from the *H.M.S. Challenger* Expedition. The female syntype is also in the collection. striatus—Megalurus striatus Jerdon, 1844 . Madras J. Lit. Sci., 13, pt. 1: 169. SYNTYPE, Adult (relaxed mount). Reg. no. 1845.1.10.24. Madras. Collected and presented by T. C. Jerdon. There are other syntypes in the collection. lected by Dr. Jerdon. Tweeddale Collection. There are two other syntypes in the collection. striatus—Zosterornis striatus Ogilvie-Grant, 1894. Bull. Br. Orn. Cl., 4:2. SYNTYPE, Adult male. Reg. no. 1897. 5.13.129. Molino, Isabella district, north-east Luzon, Philippines, 24 May 1894. Collected by J. Whitehead and presented by the Subscribers to the Whitehead Expedition. A female syntype is also in the collection. stricklandii—Copsychus stricklandii Motley and Dillwyn, 1855 . 31 Contributions to the natural history of Labuan and Borneo: 20, pl. 4. HOLOTYPE, Adult. Reg. no. 1888.4.20.207. Labuan, Borneo. Collected by J. Motley. Tweeddale Collection.

strictifrons—Euphonia strictifrons Strickland, 1851

Collection.

in Jardine, Contributions to Ornithology: 72, 84. SYNTYPE, Adult. Reg. no. 1885.6.12.108. Cayenne. Collected for Argent. Sclater

Strickland states (op. cit. pl. 71) that he had seen specimens in Sclater's Collection as well as his own.
stridulus—Pomatorhinus stridulus Swinhoe, 1861 32H Ibis, (1) 3:265. HOLOTYPE, Adult. Reg. no. 1898.10.3.605. Pehling Hills, Foochow, south China, 28 May 1857. Collected by R. Swinhoe. Seebohm Bequest.
striga—Ceblepyris striga Horsfield, 1821
strigatus—Lanius strigatus Eyton, 1839
strigatus—Turdus strigatus Horsfield, 1821
strigiceps—Zonotrichia strigiceps Gould, 1839
strigula—Cimyris strigula Hodgson, 1837
strigula—Siva strigula Hodgson, 1838
striolata—Cecropis striolata Rüppell, 1845

SYNTYPE, Adult. Reg. no. 1845.6.2.11. Shoa, Abyssinia. Collected by and purchased of E. Rüppell.

The other syntypes were in the Senckenberg Museum, Frankfurtam-Main.

striolata—Pyrrhula striolata Rüppell, 1840
styani—Pericrocotus brevirostris styani E. C. S. Baker, 1920 20 Bull. Br. Orn. Cl., 40:117. SYNTYPE, Adult male. Reg. no. 1914.5.6.358. Sechuen, western China, 29 May 1902. Collected and presented by F. W. Styan. The female syntype is also in the collection.
styani—Pomatorhinus styani Seebohm, 1884
styani—Suthora styani Rippon, 1903
suavis—Copsychus suavis P. L. Sclater, 1861
suavissima—Pipra suavissima Salvin and Godman, 1882 8 Ibis, (4) 6:79, pl. 1. SYNTYPE, Adult male. Reg. no. 1888.1.20. 163. Merume Mountains, British Guiana, 26 July 1881. Collected by H. Whitely. Salvin-Godman Collection. There are several other syntypes in the collection.
subadusta—Alseonax subadusta Shelley, 1897
subaffinis—Phylloscopus subaffinis Ogilvie-Grant, 1900
subalaris—Anabates subalaris P. L. Sclater, 1859

Listed	as type	by Sclater,	Cat. Bds.	Brit. A	Mus., 15	(1890):109
There ar	e several	other synt	ypes in th	e collect	tion.	

- subalaris—Bradyornis subalaris Sharpe, 1873

 Proc. zool. Soc. Lond., 1873: 713, pl. 58.
 no. 1873.9.9.3. Mombasa, East Africa. Collected and presented by the Rev. T. Wakefield.

- subcaerulatus—Garrulax subcaerulatus Hume, 1878 32B Stray Feathers, 7:140. Syntype, Adult. Reg. no. 1886.10.1. 4186. Near Shillong, Khasia Hills. Collected by J. Cockburn. Hume Collection.

There are several other syntypes in the collection.

- subcaeruleus—Sitta neumayer subcaeruleus Meinertzhagen, 1938 . 43

 Bull. Brit. Orn. Cl., 58:96. HOLOTYPE, Adult male. Reg. no. 1965.M.16. Haibak, 3,000', Afghan Turkestan, 17 May 1937. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- subcinerea—Tanagra subcinerea P. L. Sclater, 1861 51

 Proc. zool. Soc. Lond., 1861: 129. SYNTYPE, Adult. Reg. no. 1885.6.12.524. Venezuela. Collected for Verreaux. Sclater Collection.

Listed as the type by Sclater, Catalogue of a collection of American Birds. (1862): 77.

- subcinereus—Turdus subcinereus P. L. Sclater, 1866 39

 Proc. zool. Soc. Lond., 1866: 320. HOLOTYPE, Adult. Reg. no. 1873.8.6.2. Chile (=Australia, vide Sclater, Proc. zool. Soc. Lond., 1873: 555). 1863. Presented by P. L. Sclater who purchased it of Verreaux.
- subcinnamomea—Drymoica subcinnamomea Smith, 1847 35

 Illustrations to the Zoology of South Africa, Aves: pl. 111. HoloType, Adult (relaxed mount). Reg. no. 1845.7.6.149. Kamiesberg, Little Namaqualand. Collected by and purchased of A.
 Smith.

- subfasciatus—Thamnophilus subfasciatus P. L. Sclater and Salvin, 1876 4
 Proc. zool. Soc. Lond., 1876: 357, pl. 33. Syntype, Adult male.
 Reg. no. 1889.7.10.222. Tilotilo, Yungas Province, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.
 There is a female syntype in the collection.
- subflava—Serpophaga subflava P. L. Sclater and Salvin, 1873.
 Nomencl. Av. Neotrop., 1873: 158. HOLOTYPE, Reg. no. 1888.1.
 13.357. Pará, northern Brazil. Collected by A. R. Wallace. Sclater Collection.
- subflavum—Dicaeum minullum subflavum E. C. S. Baker, 1922 . 45
 Bull. Brit. Orn. Cl., 42:12. HOLOTYPE, Adult male. Reg. no. 1887.2.1.253. Belgaum, India, 19 Sept. 1879. (Collected by Capt. Butler, on label.) Hume Collection.

subfusa—Garrulax pectoralis subfusa Kinnear, 1924 32E Bull. Br. Orn. Cl., 44:103. HOLOTYPE, Adult male. Reg. no. 1888.4.20.778. Meitan, Tenasserim, 8 Nov. 1872. (Feb. on label.) Collected by O. Limborg. Tweeddale Collection.
subfusca—Hirundo subfusca Gould, 1856
subhemalayana—Prinia subhemalayana Blyth, 1844
subhemalayana—Tichodroma phaenicoptera subhemalayana Hodgson, 1844
subhemalayanus—Troglodytes subhemalayanus J. E. Gray, 1846 . 28 Catalogue of Mammalia and Birds of Nepal presented by B. H. Hodgson: 151. ?Syntype, Adult. Reg. no. 1846.1.9.791. Nepal. Collected and presented by B. H. Hodgson. There are several other specimens in the collection which are probable syntypes.
subhimachalus—Corythus? subhimachalus Hodgson, 1836 59 Asiat. Res., 19:152. SYNTYPE, Adult male (relaxed mount). Reg. no. 1843.1.13.836. Nepal. Collected and presented by B. H. Hodgson. There are other syntypes in the collection.
sublineatus—Graucalus sublineatus P. L. Sclater, 1879
submoniliger—Anthipes submoniliger Hume, 1877 Stray Feathers, 5:105. SYNTYPE, Adult male. Reg. no. 1886. 4.1.4090. Central Tenasserim, (near Meetan, 25 Feb. 1877, on label). Collected by W. Davison. Hume Collection. Labelled as the type on the original label. There are other syntypes in the collection.

subniger—Hypsipetes subniger Hume, 1877. Stray Feathers, 5: 109. SYNTYPE, Adult female. Reg. no. 1886. 9.1.937. Meetan, Burma, 11 Feb. 1877. Collected by W. Davison. Hume Collection

There is also a male syntype in the collection.

- subochraceum—Pellorneum subochraceum Swinhoe, 1871 . 32B Ann. Mag. nat. Hist., (4) 7:257. HOLOTYPE, Adult. Reg. no. 1898.10.3.720. Tenasserim. Collected by E. Blyth. Seebohm Bequest.
- subochraceus—Myiobius subochraceus P. L. Sclater, 1887 Proc. zool. Soc. Lond., 1887: 50. HOLOTYPE, Female (=immature, vide Berlepsch, Ornis, 14 (1907): 481). Reg. no. 1888.1.1.1246. Tilotilo, Yungas Province, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.
- subocularis—Glyciphila? subocularis Gould, 1838. Synopsis of the birds of Australia, (1838) pt. 4:6. (The description in Proc. zool. Soc. Lond, 1837: 154, was not published until Dec. 1838). HOLOTYPE, immature. Reg. no. 1881.5.1.4258. New South Wales (N.W. Australia, on label and vide Gould, Birds of Australia, 1 (1865): 502). Collected by Lieut. Emery of H.M.S. Beagle. Gould Collection.

That this was the type was confirmed by Gould, Birds of Australia, 1 (1865): 502, but Mathews, Birds of Australia, 11 (1924): 425, wrongly assumed that his comments on the type of G. ocularis referred to the type status of this specimen as regards the name G. subocularis. Stone, Aust. Av. Rec., 1 (1913): 173-4 claimed a juvenile from New South Wales in the collection at Philadelphia as the type, but this is not valid and has not been listed by de Schauensee, Proc. Acad. Nat. Sci. Philadelphia, 109 (1957): 123-246.

subpagana—Elainia subpagana P. L. Sclater and Salvin, 1860. Ibis, (1) 2:36. SYNTYPE, Adult female. Reg. no. 1888.1.13.516. Duenas, Guatamala, 31 July 1859. Collected by O. Salvin. Sclater Collection.

The male syntype is also in the collection.

subpallidus—Turdus subpallidus Hume, 1888 Stray Feathers, 11:132. SYNTYPE, Adult male. Reg. no. 1886. 7.8.3020. Shillong, Assam, 27 Oct. 1877. Collected by J. Cockburn. Hume Collection.

There are at least two other syntypes in the collection.

subpersonata - Motacilla subpersonata Meade-Waldo, 1901 Bull. Br. Orn. Cl., 12:27. SYNTYPE, Adult male. Reg. no. 1904. 4.12.30. Wad Moorbei, Rehamra, Morocco, 4 June 1901. Collected and presented by E. G. B. Meade-Waldo.

The female syntype is also in the collection.

subplacens—Elainea subplacens P. L. Sclater, 1862 9 Proc. zool. Soc. Lond., 1861 (publ. 1862): 407. SYNTYPE, Adult (female on label, male, in register). Reg. no. 1888.1.13.613. Pallatanga, Ecuador (Oct. 1859, on label). Collected by L. Fraser. Sclater Collection. There is another syntype in the collection.
subplumbeus—Dysithamnus subplumbeus P. L. Sclater & Salvin, 1880 4 Proc. zool. Soc. Lond., 1880: 158. SYNTYPE, Adult. Reg. no. 1889.7.10.292. Sarayacu, Ecuador. Collected by C. Buckley. Salvin-Godman Collection. Listed as type by Hellmayr, Catalogue of Birds of the Americas, 3 (1924): 256. There is another syntype in the collection.
subpudica—Synallaxis subpudica P. L. Sclater, 1874 3 Proc. zool. Soc. Lond., 1874: 10. SYNTYPE, Adult. Reg. no. 1889.5.20.152. Bogota, Colombia, 1868. Obtained through J. Verreaux by Sclater. Sclater Collection. There are two other syntypes in the collection. The bill is damaged.
subruficapilla—Drymoica subruficapilla Smith, 1843 35 Illustrations to the Zoology of South Africa, Aves: pl. 76. SYNTYPE, Adult female, (male, vide Lynes, Ibis (12) 6, Supplement, (1930): 666). Reg. no. 1843.2.28.96. South Africa. Collected by A. Smith. Purchased of the South African Society. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 288, and by Lynes, op. cit.
subrufus—Eupetes macrocercus subrufus Hachisuka, 1926 32A Bull. Br. Orn. Cl., 47:54. HOLOTYPE, Adult female. Reg. no. 1900.2.15.45. Mount Dulit, 3,000', Sarawak, Borneo, Oct. 1898. Collected and presented by C. Hose.
subsignata—Locustella subsignata Hume, 1873
subsoccata—Hirundo subsoccata Hodgson, 1844
subsquamicollis—Uroloncha striata subsquamicollis E. C. S. Baker, 1925

Substriata—Drymoica substriata Smith, 1842
substriolata—Lillia substriolata Hume, 1877
subundulata—Munia subundulata Godwin-Austen, 1874 60 Proc. zool. Soc. Lond., 1874: 48. SYNTYPE, Adult male. Reg. no. 1895.7.14.2861. Manipur, north-east India. Collected and presented by H. H. Godwin-Austen. This specimen, part of the syntypical material from the first collection, was originally selected by W. L. Sclater. There are other syntypes in the collection.
subunicolor—Trochalopteron subunicolor Blyth, 1843 32B J. Asiat. Soc. Bengal, 12:952. SYNTYPE, Immature (relaxed mount). Reg. no. 1843.1.13.1303. Nepal. Collected and presented by B. H. Hodgson. Blyth used Hodgson's MS. name.
subviridis—Phyllomyias subviridis Pelzeln, 1869 9 Ornithologie Brasiliens: 105, 175. SYNTYPE, Adult male (relaxed mount). Reg. no. 1888.1.13.431. Rio, Brazil, 20 July 1818. Collected by J. Natterer. Sclater Collection (received 1869). Pelzeln used Natterer's MS. name and original number. This specimen was listed by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 122 as an "authentic specimen".
subviridis—Reguloides subviridis, 1872
successor—Legatus albicollis successor Chubb, 1919 9 Bull. Brit. Orn. Cl., 40: 43. SYNTYPE, Adult male. Reg. no. 1922. 3.5.4647. British Guiana. Collected by F. V. McConnell. McConnell Collection. There are several other syntypes in the collection.
sudanac—Eremomela flaviventris sudanae Stoneham, 1925 35 Bull. Brit. Orn. Cl., 45:77. SYNTYPE, Adult male. Reg. no. 1915.12.24.1032. Sennar, Sudan, (Senga, Blue Nile, on label), 28 Dec. 1913. Collected by H. F. Stoneham. Presented by A. L. Butler. The female syntype is also in the collection.

sudanensis—Amadina sudanensis Boyd Alexander, 1907 60
Bull. Brit. Orn. Cl., 19:104. SYNTYPE, Adult male. Reg. no. 1911.12.23.3290. Near Kukawa, Northern Nigeria, 22 Nov. 1905, (in error, - 24 Nov. 1904, on label). Collected by Boyd
Alexander on the Alexander-Gosling Expedition. Boyd Alexander Bequest.
The female syntype is also in the collection.
sudanensis—Bradypterus baboecala sudanensis C. H. B. Grant and Mackworth-Praed, 1941
sudanensis—Cotile sudanensis Boyd Alexander, 1908
sudanensis—Myrmecocichla aethiops sudanensis Lynes, 1920 31 Bull. Br. Orn. Cl., 41:18. Holotype, Adult male. Reg. no. 1920.12.22.201. El Fasher, Darfur, Sudan, 13 March 1920. Collected and presented by H. Lynes.
sudanensis—Tschagra senegala sudanensis W. L. Sclater and Mack-
worth-Praed, 1918
suffusa—Suthora suffusa Swinhoe, 1871
sulaense—Dicaeum sulaense Sharpe, 1884
sulaensis—Calornis sulaensis Sharpe, 1890
sulfuriventer—Hyloterpe sulfuriventer Walden, 1872
Ann. Mag. Nat. Hist., (4) 9:399. SYNTYPE, Adult. Reg. no. 1888.5.30.341. Menado, north Celebes. Collected by B. Meyer. Tweeddale Bequest.
The second syntyne is also in the collection

- sulphurea—Stachyridopsis sulphurea Rippon, 1900 32B Bull. Brit. Orn. Cl., 11:11. HOLOTYPE, Adult. Reg. no. 1900. 12.20.287. Namehet, South Shan States, Feb. 1897. Collected and presented by G. Rippon.

 Bill broken.
- sulphureipygia—Tyrannula sulphureipygia P. L. Sclater, 1857. . . 9
 Proc. zool. Soc. Lond., 1856 (publ. 1857): 296. Holotype, Adult.
 Reg. no. 1857.7.30.19 (no. 84). Cordova, Vera Cruz, Mexico,
 Nov. 1855. Collected and presented by A. Sallé.
 Another specimen (Reg. no. 1888.1.13.825) is also marked "typus"

on the label and was listed as the type by Sclater, *Cat. Bds. Brit. Mus.*, 14 (1888): 201, but specimen no. 84 is quoted as type in the original description and the latter has no status.

There are three other syntypes in the collection.

- sumatrana—Bhringa remifer sumatrana Hachisuka, 1926 . . . 64
 Bull. Brit. Orn. Cl., 47: 57. HOLOTYPE, Adult male. Reg. no. 1920.6.29.542. Sungei Kumbang, Korinchi, Sumatra, 6 April 1914. Collected and presented by H. C. Robinson and C. B. Kloss.
- sumatrana—Cettia sumatrana Ogilvie-Grant, 1916 35

 Bull. Brit. Orn. Cl., 36:66. SYNTYPE, Adult male. Reg. no. 1920.6.29.463. Korinchi Peak, 10,000′, Sumatra, 3 May 1914. Collected and presented by H. C. Robinson and C. B. Kloss. The described female syntype is also in the collection.
- sumatrana—Notodela diana sumatrana Robinson and Kloss, 1918. 31 J.F.M.S. Mus., 8:215. SYNTYPE, Adult male. Reg. no. 1920.6. 29.459. Sungei Kumbang, 4,700', Korinchi, Sumatra, 2 April 1914. Collected and presented by H. C. Robinson and C. B. Kloss.

This is the authors' selected male. Their selected female syntype is also in the collection; other syntypes are in the Raffles Museum, Singapore.

Treubia, 13: 369. HOLOTYPE, Adult male. Reg. no. 1947.60.61. Siolak Dras, 900 metres, Korinchi Valley, Sumatra, 19 May 1914. Collected by C. B. Kloss and H. C. Robinson. Presented by the Raffles Museum, Singapore. sumatranus—Criniger sumatranus Wardlaw Ramsay, 1882 . . . 21 Ann. Mag. nat. Hist., (5) 10: 431. SYNTYPE, Adult male. Reg. no. 1888.4.1.433. Mount Sago, west Sumatra, 2 Sept. 1878. Collected by C. Bock. Tweeddale Collection. There are several other syntypes in the collection. sumatranus—Dendrobiastes hyperythrus sumatranus Hachisuka. 1920.6.29.228. Siolak Daras, 3,000', Korinchi, Sumatra, 25 March 1914. Collected and presented by H. C. Robinson and C. B. Kloss. sumatranus—Dicrurus sumatranus Wardlaw Ramsay, 1880 Proc. zool. Soc. Lond., 1880: 15. SYNTYPE, Adult male. Reg. no. 1887.11.20.742. West Sumatra. Collected by C. Bock. Tweeddale Collection. There are five other syntypes in the collection. sumatranus—Hemixus sumatranus Wardlaw Ramsay, 1882 . . . 21
Ann. Mag. nat. Hist., (5) 10:431. HOLOTYPE, Adult. Reg. no. 1888.4.1.305. Mount Singalan, west Sumatra, 28 June 1878. Collected by Beccari. Tweeddale Collection.
Obtained by Ramsay from Count Salvadori. sumatranus-Henicurus velatus sumatranus Robinson and Kloss, 1923

J.F.M.S. Mus., 5:56. SYNTYPE, Adult male. Reg. no. 1936.4. 12.145. Scolak Dras, 3,000', Korinchi, Sumatra, 27 March 1914. Collected by H. C. Robinson and C. B. Kloss and presented by the Raffles Museum, Singapore. This is the authors' selected male. Their selected female syntype

sumatrana—Zosterops aureiventer sumatrana Kloss, 1931 .

sumatrensis—Cryptolopha sumatrensis Robinson and Kloss, 1916. 35

J. Straits Branch roy. Asiat. Soc., 73: 277. Syntype, Adult male.
Reg. no. 1920.6.29.247. Sungei, Kumbang, Korinchi, West Sumatra, 1 April 1914. Collected and presented by H. C. Robinson and C. B. Kloss.

The female syntype is also in the collection.

is also in the collection.

superciliosa—Embernagra superciliosa Salvin, 1865

Collected by E. Arcé. Salvin-Godman Collection.

Proc. zool. Soc. Lond., 1864 (publ. 1865): 582. HOLOTYPE, Adult. Reg. no. 1885.12.14.1342. Bebedero, Nicoya, Costa Rica, 1864.

sumichrasti—Embernagra sumichrasti Sharpe, 1888

superciliosa—Gerygone superciliosa Wallace, 1864.	. 3:
Proc. zool. Soc. Lond., 1863 (publ. 1864): 491. SYNTYPE, A	Adult
male. Reg. no. 1873.5.12.212. East Timor. (Delhi, on 1	abel;
this is a Timor locality, vide Ticehurst, Systematic review of the	genus
Phylloscopus, (1938): 183), 1861. Collected by A. R. Wal	Ílace.
Wallace Collection.	
This specimen was registered as the type. There is an	other

This specimen was registered as the type. There is another syntype in the collection.

- superciliosa—Ochthoeca superciliosa P. L. Sclater and Salvin, 1871 Proc. zool. Soc. Lond., 1870 (publ. 1871): 786. SYNTYPE, Adult female. Reg. no. 1888.1.13.52. Paramo of Merida, Venezuela, July 1869. Collected by A. Goering. Sclater Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 20.

There is a translation of the description by F. C. Craemers in Rowley, *Ornithology Miscellany*, 2 (1877): 189.

superciliosus—Hyphantornis superciliosus Shelley, 1873. . . . 61

Ibis, (3) 3:140. Syntype, Adult (female, vide Sharpe, Cat. Bds.

Brit. Mus., 13 (1890):471). Reg. no. 1877.7.11.227. West Africa, March 1872. Collected by W. W. Saunders. Sharpe Collection.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 471. The other syntype is in the collection.

- superciliosus—Pomatorhimus superciliosus Vigors and Horsfield, 1826 32B Trans. Linn. Soc. Lond., 15:330. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.72. South coast of New Holland. Collected by R. Brown and presented by the Linnean Society.

C. r. Acad. Sci., Paris, 1854: 59. Holotype, (Immature female, vide, Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 62). Reg. no. 1852. 5.12.10. Sumatra. Collected by A. Delattre and purchased of Verreaux.
sushkini—Prunella fulvescens sushkini. See appendix, Corrections and Additions, vol. 2
swainsoni—Criniger swainsoni Neumann, 1914
swainsoni—Hippolais swainsoni Hodgson, 1844
swainsoni—Petrochelidon swainsoni P. L. Sclater, 1858
swanzii—Drymoeca swanzii Sharpe, 1870
swynnertoni—Erythracus swynnertoni Shelley, 1906 31 Bull. Br. Orn. Cl., 16: 125. SYNTYPE, Adult male. Reg. no. 1910.7.1.517. Chirinda Forest, 4,000', Gazaland, Southern Rhodesia, 17 June 1905. Collected by and purchased of C. F. M. Swynnerton. This is the author's selected syntype. There are several other syntypes in the collection.
swynnertoni—Trochocercus albonotatus swynnertoni Neumann, 1908 38 Bull. Brit. Orn. Cl., 23:46. HOLOTYPE, Adult male. Reg. no. 1910.7.1.361. Chirindi, 3,800', Southern Rhodesia, 12 July 1907. Collected for and purchased of C. F. M. Swynnerton.
swynnertoni—Turdus swynnertoni Bannerman, 1913 31 Bull. Br. Orn. Cl., 31:56. HOLOTYPE, Adult female. Reg. no. 1910.7.1.466. Chirinda Forest, Rhodesia, 17 June 1906. Collected by and purchased of C. F. M. Swynnerton.

- sykesi—Hypothymis azurea sykesi E. C. S. Baker, 1920 . . . 37 Bull. Brit. Orn. Cl., 41 (1920): 9. New name for Muscicapa caeruleocephala Sykes, 1832 (q.v.) on the grounds of its preoccupation by M. caeruleocephala Scopoli, 1786.

There is another syntype in the collection, and two more in the University Museum of Zoology, Cambridge.

There are several other syntypes in the collection. Vaurie, *The Birds of the Palearctic Fauna: Passeriformes*, (1959): 73, and Vaurie, in Peters' *Checklist of Birds of the World*, 9 (1960): 162, uses *Heterura sylvana* Blyth, *J. Asiat. Soc. Bengal*, 14 (1845): 556. This is incorrect, *vide* B. P. Hall, *Bull. Br. Mus. nat. Hist.* (*Zool.*), 7, no. 5 (1961): 289.

available at the time of the description.

This is the author's selected syntype. There are two other male syntypes in the collection.

- syriacus—Troglodytes troglodytes syriacus R. Meinertzhagen, 1933 28 Bull. Br. Orn. Cl., 54: 20. HOLOTYPE, Adult male. Reg. no. 1955.M.52. Zachle, 3,000', Lebanon, 30 Oct. 1919. Collected by R. Meinertzhagen. Meinertzhagen Collection.

- taczanowskii—Buarremon taczanowskii P. L. Sclater and Salvin, 1875 49 Proc. zool. Soc. Lond., 1875: 236. Since this is a new name for B. mystacalis Taczanowski, 1874, on the grounds of preoccupation, the type is that of the original description, and the type listed by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 268, is invalid.
- taczanowskii—Euphonia chlorotica taczanowskii P. L. Sclater, 1886 51 Cat. Bds. Brit. Mus., 11:65. SYNTYPE, Adult male. Reg. no. 1885.6.12.58. Callacate, Peru, 1879. Collected by J. Stolzmann. Sclater Collection.

There are five other syntypes in the collection.

There is another syntype in the Academy of Natural Science of Philadelphia.

- tahanensis—Mesia argentauris tahanensis Yen, 1934 32B Extr. Science J. (University of Sun Yatsen, Canton), 6, no. 2:379 (63 in separately paged reprint, vide Deignan, in Peters, Checklist of Birds of the World, 10 (1964):382). HOLOTYPE, Adult male. Reg. no. 1906.7.23.82. Gunong Tahan, 5,300-6,000', Malaya, 19 June 1905. Collected by H. C. Robinson and presented by the Selangore State Museum.

 Yen apparently used Robinson's MS, name (vide Chasen, Handlist)
- of Malayan Birds, (1935): 230).

 tahanensis—Zosterops tahanensis Ogilvie-Grant, 1906 47

 Bull. Brit. Orn. Cl., 19: 10. HOLOTYPE, Adult male. Reg. no. 1906.7.23.13. Gunong Tahan, 5,300-6,000′, north of Pahang,

Malaya, 7 July 1905. Collected by H. C. Robinson. Presented by the Selangore State Museum.

taiti—Acrocephalus taiti Ogilvie-Grant, 1913	5
taiti—Aegithalus caudatus taiti Ingram, 1913 4 Zoologist, (4) 17:137. HOLOTYPE, Adult. Reg. no. 1886.11.20.8. Coimbra, Portugal, Oct. 1886. Collected for and presented by Dr. Lopex Viera.	0
taivana—Budytes taivana Swinhoe, 1863	9
taivanicus—Dendrobiastes hyperythrus taivanicus Hachisuka, 1926 3 Bull. Brit. Orn. Cl., 46:52. HOLOTYPE, Adult male. Reg. no. 1897.7.2.494. (not 1899 as stated in the description). Tongapo, Formosa, 9 Jan. 1894. Collected by P. A. Holst. Seebohm Bequest.	7
taivanorum—Horornis canturians taivanorum Hachisuka, 1926. 3 Bull. Brit. Orn. Cl., 47:53. Holotype, Adult male. Reg. no. 1905.12.24.722. Hills near Tamsui, North Formosa, 24 Feb. 1895. Collected for J. de la Touche. Presented by C. B. Rickett.	5
taivanus—Garrulus taivanus Gould, 1862	
taivanus—Pycnonotus taivanus Styan, 1893	1
talacoma—Prionops talacoma Smith, 1836	3
talboti—Bathmedonia talboti Boyd Alexander, 1907 3 Bull. Brit. Orn. Cl., 19: 46. HOLOTYPE, Adult female. Reg. no. 1911.12.23.2103. Joga River, Libowka, Welle River, Belgian Congo, 7 March 1906. Collected by W. Boyd Alexander. Boyd Alexander Bequest.	5

talifuensis—Aegithaliscus talifuensis Rippon, 1903 Bull. Brit. Orn. Cl., 14:18. SYNTYPE, Adult. Reg. no. 1903.8.8. 20. Gyi-dzin-Shan, 7.000', east of Talifu, west Yunnan, 8 April 1902. Collected and presented by G. Rippon. This specimen is labelled as the type but a second syntype was collected at the same date. talifuensis—Anorthura talifuensis Sharpe, 1902 Bull. Br. Orn. Cl., 13:11. SYNTYPE, Adult. Reg. no. 1903.8.8. 485. Gyi-Dzin-Shan, 8,000', east of Talifu, north Yunnan, 4 March 1902. Collected and presented by Lt. Col. G. Rippon. There is a second syntype in the collection. tamucoensis—Upucerthia tamucoensis Chubb, 1911 Bull. Br. Orn. Cl., 27:101. SYNTYPE, Adult male. Reg. no. 1911.4.25.115. Pelal, Tamuco, southern Chile, 14 June 1910. Collected by and purchased of A. Saldana. There is another male syntype in the collection. tandae—Amblyospiza albifrons tandae Bannerman, 1921 Bull. Br. Orn. Cl., 41: 136. HOLOTYPE, Adult male. Reg. no. 1909.8.5.237. N'dalla Tando, northern Angola, 1 Jan. 1909. Collected by and purchased of W. J. Ansorge. tando—Nicator vireo tando Serle, 1952 23 Bull. Br Orn. Cl., 72:95. HOLOTYPE, Adult male. Reg. no. 1909.8.5.112. N'Dalla Tando, north Angola, 15 Oct. 1908. Collected by and purchased of W. J. Ansorge. tando-Serinus mozambicus tando W. L. Sclater and Mackworth-Ndala Tando, Northern Angola, 16 Sept. 1908. Collected by and purchased of W. J. Ansorge. tando—Sylvietta virens tando W. L. Sclater, 1927 Bull. Brit. Orn. Cl., 48:18. HOLOTYPE, Adult male. Reg. no. 1910.5.6.906. N'Dalla Tando, Northern Angola, 31 Oct. 1908. Collected by and purchased of W. L. Ansorge. tanganyicae—Cinnyris tanganyicae Ogilvie-Grant, 1907 Bull. Brit. Orn. Cl., 19:105. HOLOTYPE, Adult male. Reg. no. 1907.6.26.88. West of Lake Tanganyika, 4,000', (about 30 miles south-west of Baraka by date, vide Ibis, (9) 2 (1908): 264), 4 Jan. 1907. Collected by and purchased of D. Carruthers. tanganyika—Cisticola aridula tanganyika Lynes, 1930 35 Ibis, (12) 6, Supplement: 126, pl. 4. HOLOTYPE, Adult male. no. 1928.10.25.3. Morogoro, Tanganyika, 25 April 1927. Collected by Admiral H. Lynes and B. B. Osmaston. Presented of Admiral H. Lynes.

tardus—Rhinomvias tardus Robinson and Kloss, 1915 J. Fed. Malay Sts. Mus., 61:29. HOLOTYPE, Adult male, Reg. no. 1936.4.12.2033. Genting Bidai, 2,300', Selangor-Pahang Boundary, Malaya, 19 Sept. 1914. Collected for and presented by the Raffles Museum, Singapore. taupoensis—Anthus novaeseelandiae taupoensis Mathews, 1930 . 19 Bull. Br. Orn. Cl., 50:42. HOLOTYPE, Adult. Reg. no. 1910.7.7. 203. Lake Taupo, North Island, New Zealand, March 1910. Collected and presented by Miss A. Chirnside. taviunensis—Ptilotis procerior taviunensis Wiglesworth, 1891. Abh. Ber. K. zool. anthrop.-ethn. Mus. Dresden, 6:34. New name for Ptilotis similis Layard, 1876 (q.v.), on the grounds of its preoccupation by P. similis Pucheron, 1853. taylori—Calliste taylori Taczanowski and Berlepsch, 1885 Proc. zool. Soc. Lond., 1885: 78. SYNTYPE, Adult. Reg. no. 1886.5.12.370. Machay, Ecuador, 20 Dec. 1883. Collected by J. Stolzmann and A. Taylor. Sclater Collection. Tazcanowski and Berlepsch used a Stolzmann MS. name. This specimen was listed as one of the types by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 130. taylori—Hypacanthis spinoides taylori Kinnear, 1939 59 Ibis, (14) 3:752. HOLOTYPE, Adult male. Reg. no. 1938.12.13.1. Lilung, 9,800', Tsangpo Valley, south-east Tibet, 26 May 1938. Collected by F. Ludlow and G. Sherriff. Ludlow-Sherriff Expedition. Presented by the Godman-Thomas Fund. taylori—Pitangus taylori P. L. Sclater, 1864 Ibis, (1) 6:169. SYNTYPE, Adult. Reg. no. 1905.6.28.291. Porto Rico, West Indies, May 1863. Collected and bequeathed by E. Cavendish Taylor. There is another syntype in the collection. Sclater Ibis. (8) 5 (1905): 500 refers to the present specimen as the type but used both in the original description and lists the other as the type in Cat. Bds. Brit. Mus., 14 (1888): 180. tectirostris—Bhringa textirostris Hodgson, 1837. The Indian Review, 1 (1836, publ. 1837): 325. ?SYNTYPE, Adult. Reg. no. 1843.1.13.709. Nepal. Collected and presented by B. H. Hodgson. Received with his main collection, the specimen has Hodgson's drawing number on the label and is therefore likely to have been before him when the description was made. temmincki—Arachnothera temmincki Moore, 1858 in Horsfield and Moore, Catalogue of birds in the Museum of the

Honourable East India Company, 2:728. HOLOTYPE, Adult male. Reg. no. 1860.4.16.476. (Indian Archipelago, vide Gadow, Cat.

- Bds. Brit. Mus., 9 (1884): 103.) Collected by Middleton. Presented by the Honourable East India Company Museum.

 Marked as the type on the original label.
- temminckii—Orthonyx temminckii Vigors and Horsfield, 1827 . 32A Trans. Linn. Soc. Lond., 15. (1826, publ. 1827): 294. HOLOTYPE, Adult female. Reg. no. 1863.7.7.57. Hat Hill, New South Wales, 1804. Presented by the Linnean Society as the type.

The authors imply that theirs is the first description, but Mathews, Birds of Australia, 9 (1921): 175 has traced a reference to Ranzani, Elementi di Zoologia, 3 (1822) pt. 3:19, as the first author, although he does not appear to have been subsequently cited.

There is another syntype in the collection.

In Latham, Supplement 2 to the general synopsis of birds, (1801): 211 the author refers to several drawings. Vol. 2, no. 37 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2).

- temporalis—Pomatorhinus temporalis Vigors & Horsfield, 1827 . 32B Trans. Linn. Soc. Lond., 15 (1826, pubd. 1827): 330. HOLOTYPE, Adult (relaxed mount). Reg. no. 1863.7.7.71. Shoalwater Bay, Queensland, Aug. 1802. Collected by R. Brown and presented by the Linnean Society.
- tenebricosa—Apalis melanocephala tenebricosa Vincent, 1933 . 35
 Bull. Brit. Orn. Cl., 53: 141. HOLOTYPE, Adult male. Reg. no. 1933.3.1.10. Namuli Mountain, 4,800', Quelimane Province, Portuguese East Africa, 15°21' S., 37°04' E., 8 August 1932. Collected by J. Vincent. Presented by the Portuguese East Africa Expedition.
- tenebrosus—Pipilo fuscus tenebrosus van Rossem, 1938. . . . 49
 Bull. Brit. Orn. Cl., 58: 132. HOLOTYPE, Adult male. Reg. no.

1899.2.1.3505.	Zapotlan, (Jalisco	o, on label) Mexico	o, 23 Nov. 188	9
		Salvin-Godman C		

- tenellipes—Phylloscopus tenellipes Swinhoe, 1860. 35

 Ibis, (1) 2:53. HOLOTYPE, Adult, (female, on label). Reg. no. 1898.9.1.391. Amoy, China, 12 Oct. 1855. Collected by R. Swinhoe. Seebohm Collection.

No single type was designated, but this is one of the specimens available. A specimen collected after the date of publication was incorrectly labelled as the type.

- tennenti—Cossypha natalensis tennenti Williams, 1962. . . . 31

 Bull. Br. Orn. Cl., 82:137. HOLOTYPE, Adult male. Reg. no. 1963.20.1. Mount Endau, 3,400′, Kitui District, Kenya, 25 Jan. 1962. Collected and presented by J. R. M. Tennent.

Vol. 2, no. 80 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 11 (1924): 339.

- tenuirostris—Gracaulus tenuirostris Jardine & Selby, 1831 . . . 20 Edinburgh J. Nat. Geog. Sci., (n.s.) 4:211. (Illustrations of Ornithology, 3, pt. 8:pl. 114). HOLOTYPE, Adult. Reg. no. 1886.6. 24.124. Australia. Jardine Collection; purchased of E. Gerrard.

- tenuirostris—Lamprotornis tenuirostris Rüppell, 1836 62

 Neue Wirbelthiere zu der Fauna von Abyssinien, Vög.: 26, pl. 10.

 SYNTYPE, female or immature (relaxed mount). Reg. no. 1842.

 8.16.19. Abyssinia. Collected by and purchased of E. Rüppell.

 There is another syntype in the Senckenberg Museum, Frankfurten-Main.
- tenuirostris—Sittella tenuirostris Gould, 1865 43

 Handbook to the birds of Australia, 1:610. Holotype, (Immature male, vide Gadow, Cat. Bds. Brit. Mus., 8 (1883):363). Reg. no. 1881.5.1.4341. (?South Australia, vide Gadow, op. cit.). Collected by C. Sturt. Gould Collection.
- tephrogenys—Trichophorus tephrogenys Jardine & Selby, 1833 . 21 Illustrations of Ornithology, 3: pl. 127. Holotype, Adult. Reg. no. 1886.6.24.250. ". . . we suspect it to be Indian" (type locality restricted to Malacca by Hartert, Novit. Zool., 9 (1902): 558). Jardine Collection; purchased of E. Gerrard.
- tephrolaemus—Nectarinia tephrolaemus Jardine and Fraser, 1851. 46
 Contributions to Ornithology,: 154. HOLOTYPE, Adult male. Reg.
 no. 1886.6.24.315. Clarence, Fernando Po, 1850. Collected by
 L. Fraser. Jardine Collection.

We consider that Jardine's label inscribed "N. cyanolaimus" is an error and that this is the type specimen.

- tephropleurus—Zosterops tephropleurus Gould, 1855 47

 Proc. zool. Soc. Lond., 1855 : 166. HOLOTYPE, Adult male. Reg.
 no. 1855.10.23.1. Lord Howe's Island. Collected by J.

 Macgillivray. Presented by the Museum of Economic Geography,
 from the voyage of H.M.S. Herald.

terrestris—Drymoica terrestris Smith, 1842
terrestris—Synallaxis terrestris Jardine, 1847
terricolor—Iole terricolor Hume, 1878
terricolor—Synallaxis terricolor P. L. Sclater & Salvin, 1866 3 Proc. zool. Soc. Lond., 1866: 183. SYNTYPE, Adult. Reg. no. 1889.5.20.174. Upper Ucayali River, east Peru, Sept. 1865. Collected by E. Bartlett. Sclater Collection. There are three other syntypes in the collection.
tertale—Passer griseus tertale Benson, 1942
tertia—Alaemon hamertoni tertia Clarke, 1919
testacea—Pyranga testacea P. L. Sclater and Salvin, 1868 51 Proc. zool. Soc. Lond., 1868 : 388. Syntype, Adult male. Reg. no. 1885.6.7.367. Chitra, Veragua, 1869. Collected by E. Arcé. Salvin-Godman Collection. There are three other syntypes in the collection.
thai—Dicrurus macrocercus thai Kloss, 1921
thai—Tephrodornis pondicerianus thai Kloss & Chasen, 1926 20 Bull. Br. Orn. Cl., 46: 58. HOLOTYPE, Adult male. Reg. no. 1925. 7.12.1. Ta Chang Thai, 600', Raheng, west Siam, 24 July 1924. Collected for K. S. Gairdner and presented by the latter.

- thais—Molpastes aurigaster thais Kloss, 1924 21

 J. nat. Hist. Soc. Siam., 6: 291. SYNTYPE, Adult male. Reg. no. 1955.1.5678. Bangkok, central Siam, 4 March 1915 (in text; altered by the collector to April in his own copy). Collected and presented by Sir Walter Williamson.

 The other syntype is also in the collection.
- thais—Phyllergates cucullatus thais Robinson and Kloss, 1923 . 35 J. fed. Malay Sts. Mus., 11:56. HOLOTYPE, Adult male. Reg. no. 1936.4.12.142. Kao Luang, 5,000-5,800', Nakon Sri Tamarat, Siam, 30 March 1922. Collected by H. M. Pendlebury. Presented by the Raffles Museum, Singapore.
- theresae—Alauda arvensis theresae R. Meinertzhagen, 1947. . . . 17

 Bull. Br. Orn. Cl., 67:93. HOLOTYPE, Adult male. Reg. no. 1965.M.28. Corofin, Co. Clare, Ireland, 6 Sept. 1937. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Anthus pratensis theresae R. Meinertzhagen, 1953. . . . 19
 Bull. Br. Orn. Cl., 73:43. HOLOTYPE, Adult female. Reg. no. 1965.M.26. Achill Island, west Ireland, 12 Jan. 1947. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Argya caudatus theresae R. Meinertzhagen, 1930 . . . 32B Bull. Br. Orn. Cl., 50: 55. HOLOTYPE, Adult male. Reg. no. 1965. M.42. Baghdad, 28 Dec. 1922. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Calandrella sclateri theresae R. Meinertzhagen, 1949 . 17

 Bull. Br. Orn. Cl., 69: 106. HOLOTYPE, Adult female. Reg. no. 1965.M.32. 25 miles east of Pofadder, Namaqualand, 6 May 1949. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Emberiza striolata theresae R. Meinertzhagen, 1939 . 49
 Bull. Brit. Orn. Cl., 59: 68. HOLOTYPE, Adult male. Reg. no.
 1965-M-19. Andja, South-west Morocco, 5 Nov. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Erythrospiza githaginea theresae R. Meinertzhagen, 1939 59 Bull. Brit. Orn. Cl., 59:64. HOLOTYPE, Adult male. Reg. no.

- 1965.M.36. Taznakht, Anti Atlas, Ouarzazate District, Morocco, 22 Nov. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection
- theresae—Galerida theklae theresae R. Meinertzhagen, 1939 . . . 17

 Bull. Br. Orn. Cl., 59:65. HOLOTYPE, Adult male. Reg. no. 1965.M.34. Anti Atlas Mts, 30 km. south of Tiznit, southern Morocco, 5 Nov. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Garrulus glandarius theresae R. Meinertzhagen, 1939 . 71 Bull. Brit. Orn. Cl., 59:67. HOLOTYPE, Adult female. Reg. no. 1965.M.45. Azrou, Middle Atlas, Morocco, 4 Dec. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Lanius excubitor theresae R. Meinertzhagen, 1953 . . . 23

 Bull. Br. Orn. Cl., 73:72. HOLOTYPE, Adult male. Reg. no. 1965.M.48. Huleh Lake, north Palestine, 1 April 1953. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Montifringilla theresae R. Meinertzhagen, 1937 . . . 61
 Bull. Br. Orn. Cl., 58:10. HOLOTYPE, Adult male. Reg. no. 1965.M.22. Shibar Pass, 9,800', northern Afghanistan, 19 April 1937. Meinertzhagen Collection.
- theresae—Oenanthe moesta theresae R. Meinertzhagen, 1939 . . . 31

 Bull. Br. Orn. Cl., 59:66. HOLOTYPE, Adult male. Reg. no. 1965.M.44. Near Tiznit, south-west Morocco, 7 Nov. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection.

- theresae—Saxicola torquata theresae R. Meinertzhagen, 1934. . 31 Ibis, (13) 4:56. HOLOTYPE, Adult female. Reg. no. 1965.M.31. Grogarry, South Uist, Outer Hebrides, 11 Nov. 1932. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Scotocerca inquieta theresae R. Meinertzhagen, 1939 . 35

 Bull. Brit. Orn. Cl., 59:65. HOLOTYPE, Adult male. Reg. no. 1965.M.23. Izakarm, Moroccan Sahara, 5 Nov. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection.
- theresae—Serinus albogularis theresae R. Meinertzhagen, 1949 . 59
 Bull. Brit. Orn. Cl., 69:107. Holotype, Adult male. Reg. no.
 1965.M.35. 50 miles east of Springbok, Little Namaqualand,
 Cape Province, 7 May 1949. Collected by R. Meinertzhagen.
 Meinertzhagen Collection.

theresae—Sylvia nana theresae R. Meinertzhagen, 1937. Bull. Brit. Orn. Cl., 58:10. HOLOTYPE, Adult male. Reg. no. 1965.M.27. Rohri, Sind, India, 25 Jan. 1937. Collected by A. Jones. Meinertzhagen Collection. theresae—Turdus viscivorus theresae R. Meinertzhagen, 1939. Bull. Br. Orn. Cl., 59:67. HOLOTYPE, Adult male. Reg. no. 1965.M.41. Oulmes, Middle Atlas, Morocco, 7 Dec. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection. thermophilus—Cichlops thermophilus Hodgson, 1844 . 19 in J. E. Gray, Zoological Miscellany, 1844: 83 (nomen nudum). Four specimens in the collection were listed as types by Sharpe. Cat. Bds. Brit. Mus., 10 (1885): 569. 2333. Borders of Sikkim and Thibet, 1870. Collected by L. Mandelli. Hume Collection. thomassoni—Merula thomassoni Seebohm, 1894. Bull. Br. Orn. Cl., 3:51. SYNTYPE, Adult male. Reg. no. 1896. 6.1.647. North Luzon, Philippines, 21 Jan. 1894. Collected by J. Whitehead. Seebohm Collection. This specimen is figured in Seebohm's Monograph of the Turdidae. 2 (1902) : pl. 109. thomensis—Nectarinia thomensis Bocage, 1889 J. Sci. Math. Phys. Nat. Lisboa, (2) 1:143. SYNTYPE, Adult male. Reg. no. 1891.8.20.2. St. Miguel, St. Thomas Island, West Africa, Aug. 1889. (Collected by F. Newton, vide Shelley, Birds of Africa, 2 (1900): 120). Presented by J. V. Barboza du Bocage. Three specimens were referred to, of which this is one. thompsoni—Cerasophila thompsoni Bingham, 1900 21 Ann. Mag. nat. Hist., (7) 5:358. SYNTYPE, Adult male. Reg. no. 1900.9.2.1. Loi-San-Pa, 6,500', Möng-Köng State, South Shan States, Upper Burma, 1 Jan. 1900. Collected and presented by C. T. Bingham. There are two other syntypes in the collection. thompsoni—Suthora thompsoni Bingham, 1903 33 Bull. Br. Orn. Cl., 13:63. HOLOTYPE, Adult female. Reg. no. 1903.12.24.230. North-east of Kyatpyin, 2,500', near Paunglaung stream, Loilong State, South Shan States, Burma, 18 March 1902. Collected by H. N. Thompson and W. H. Craddock and purchased through E. Gerrard. thomsoni—Erythrocercus thomsoni Shelley, 1882 . Proc. zool. Soc. Lond., 1882: 307, pl. 16. ?HOLOTYPE, Adult. Reg. no. 1895.5.1.218. Rovuma River, Tanganyika. Collected by J. Thomson. Shelley Collection. (cont.)

Two specimens were received from this source, only one of which is an adult.
thomsoni—Manucodia thomsoni Tristram, 1889
thoracicus—Pteroptochus thoracicus P. L. Sclater, 1865
thoracicus—Thryothorus thoracicus Salvin, 1864
thoracicus—Xiphorhynchus thoracicus P. L. Sclater, 1860 2 Proc. zool. Soc. Lond., 1860: 277. HOLOTYPE, Adult male. Reg. no. 1889.5.20.602. Babahoyo, Ecuador, Aug. 1859. Collected by L. Fraser. Sclater Collection.
thruppi—Parus thruppi Shelley, 1885
tianschanica—Troglodytes tianschanica Sharpe, 1881
tibatiensis—Anomalospiza imberbis tibatiensis Bannerman, 1923 . 61 Bull. Br. Orn. Cl., 43: 143. HOLOTYPE, Adult male. Reg. no. 1923.10.26.433. Tibati, 2,850', Cameroons, 10 Oct. 1922. Collected by and purchased of G. L. Bates.
tibetana—Anorthura tiberana Walton, 1905
tibetana—Calandrella tibetana Brooks, 1880

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 585. There are several other syntypes in the collection.

tibetana—Pica tibetana Hodgson, 1849 Ann. Mag. nat. Hist., (2) 3:203. HOLOTYPE, Reg. no. 1879.11. 28.338. Tibet (1848, on label). Collected by B. H. Hodgson. Presented by the Honourable East India Company Museum.
tibetana—Prunella fulvescens tibetana Sushkin, 1925 30 Proc. Boston Soc., 38:53. HOLOTYPE, Adult male. Reg. no. 1905.12.31.231. Khambajong, base camp, 16,000', Gyangtze, south Tibet, 9 Oct. 1903. Collected and presented by H. J. Walton of the Mount Everest Expedition. (See also appendix, Corrections and Additions, vol. 2.)
tibetanus—Corvus tibetanus Hodgson, 1849
tibetanus—Passer montanus tiberanus E. C. S. Baker, 1925 61 Bull. Br. Orn. Cl., 45:92. HOLOTYPE, Adult male. Reg. no. 1905.12.31.289. Khumbajong, Tibet, 5 Oct. 1903. Collected and presented by H. J. Walton.
tibetanus—Phylloscopus tibetanus Ticehurst, 1937 35 Bull. Brit. Orn. Cl., 57: 109. Holotype, Adult male. Reg. no. 1937.1.17.622. Bimbi La, 13,500', Tsari, South Tibet, 9 June 1936. Collected by F. Ludlow. Ludlow-Sherriff Collection.
tibialis—Rectes tibialis Sharpe, 1877
tibicen—Coracius tibicen Latham, 1801
ticehursti—Alauda arvensis ticehursti Whistler, 1928 17 Bull. Br. Orn. Cl., 48:65. Holotype, Adult male. Reg. no. 1928.2.4.1. San Miguel de Oya, near Vigo, west Galicia, northwest Spain, 10 May 1927. Collected and presented by H. Witherby

Bull. Brit. Orn. Cl., 59: 69. HOLOTYPE, Adult female. Reg. no. 1965. M.29. Tinghir, Ouarzazate District, Moroccan Sahara, 24 Nov. 1938. Collected by R. Meinertzhagen. Meinertzhagen Collection.
tickelliae—Dicaeum tickelliae Blyth, 1843
tilemsiensis—Auripasser luteus tilemsiensis Bates, 1932 61 Bull. Br. Orn. Cl., 53:72. HOLOTYPE, Adult male. Reg. no. 1932.8.6.7. Taberréshat ("70 miles north-east of Burem", on label), French Sudan, 7 Sept. 1931. Collected by and purchased of G. L. Bates.
timbuktana—Tchagra senegala timbuktana Bates, 1932 23 Bull. Br. Orn. Cl., 53:74. HOLOTYPE, Adult female. Reg. no. 1932.8.6.6. Timbuktu, French Sudan, 18 Nov. 1931. Collected by and purchased of G. L. Bates.
timoriense—Edoliisoma timoriense Sharpe, 1878 20 Mitt. K. zool. Mus., Dresden, 1:369. SYNTYPE, Adult male. Reg. no. 1873.5.12.305. East Timor, 1861. Collected by A. R. Wallace. Wallace Collection. There are three other syntypes in the collection.
timoriensis—Megalurus timoriensis Wallace, 1864 35 Proc. zool. Soc. Lond., 1863 (publ. 1864): 489. SYNTYPE, Adult male. Reg. no. 1873.5.12.616. East Timor, 1861. Collected by A.R. Wallace. Wallace Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 129. A female syntype is also in the collection.
timoriensis—Petrochelidon timoriensis Sharpe, 1885
tingitanus—Corvus tingitanus Irby, 1874

- tinnabulans—Calamanthella tinnabulans Swinhoe, 1859 35 J. Roy. Asiat. Soc., North China Branch, 1859 : 225. HOLOTYPE, Adult (female, vide Lynes, Ibis, (12) 6, Supplement, (1930) : 668). Reg. no. 1888.4.20.423. Hongsan, North Formosa. Collected by R. Swinhoe. Tweeddale Collection.
- tionis—Stachyris nigriceps tionis Robinson & Kloss, 1927 . . . 32B J. F. M. S. Mus., 13:211. Holotype, Adult male. Reg. no. 1947.60.57. Juara Bay, Tioman Island, east coast of Malaya, 1 July 1916. Collected by C. B. Kloss and presented by the Raffles Museum, Singapore.
- tobagoensis—Pipromorpha oleaginea tobagoensis Chubb, 1919 . 9
 Ann. Mag. Nat. Hist., (9) 4:302. HOLOTYPE, Adult male. Reg.
 no. 1886.6.24.374. Tobago, West Indies. Collected by J. Kirk,
 Jardine Collection.
- tonkinensis—Garrulax moniliger tonkinensis Delacour, 1927. . . 32B Bull. Br. Orn. Cl., 47:158. Holotype, Adult male. Reg. no. 1927.6.5.10. Backan, north-east Tonkin, 12 Dec. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China.
- tonkinensis—Pericrocotus brevirostris tonkinensis Delacour, 1927. 20 Bull. Br. Orn. Cl., 47:156. Syntype, Adult male. Reg. no. 1927.6.5.7. Backan, north-east Tonkin, 31 Dec. 1926. Collected by J. Delacour, M. P. Jabouille and W. P. Lowe on the Third Franco-British Expedition to Indo-China.

This is the author's selected male. His selected female syntype is also in the collection.

- tonkinensis—Pitta douglasi tonkinensis Delacour, 1927. . . . 12

 Bull. Brit. Orn. Cl., 47:155. Holotype, Adult male. Reg. no.
 1927.6.5.6. Backan, north-east Tonkin, Indo-China, 23 Dec. 1926.
 Collected by J. Delacour, W. P. Lowe, and P. Jabouille, on the Third Anglo-French Expedition to Indo-China.
- tonkinensis Pomatorhimus tickelli tonkinensis Delacour, 1927 . 32B Bull. Br. Orn. Cl., 47:159. HOLOTYPE, Adult male. Reg. no. 1927.6.5.13. Lake Bahé, north-east Tonkin, 20 Dec. 1926. Collected by J. Delacour, P. Jabouille and W. P. Lowe on the Third Franço-British Expedition to Indo-China.

tonkinensis—Sitta castanea tonkinensis Kinnear, 1936 43 Bull. Brit. Orn. Cl., 56:71. Holotype, Adult male. Reg. no. 1928.6.26.1538. Napé, Laos, Indo-China, 28 Jan. 1928. Collected by J. Delacour and W. P. Lowe, from the Fourth Anglo-French Expedition to Indo-China.
tonsa—Diaphorophyia tonsa Bates, 1911
topela—Munia topela Swinhoe, 1863
toroensis—Sylviella toroensis Jackson, 1905 35 Bull. Brit. Orn. Cl., 15:38. HOLOTYPE, Adult female. Reg. no. 1905.1.10.2. Kibera River, Toro, West Uganda, 31 Jan. 1902. Collected and presented by F. J. Jackson.
toroensis—Trochocercus toroensis Jackson, 1906 38 Bull. Brit. Orn. Cl., 19: 20. Holotype, Adult female. Reg. no. 1907.6.19.1. Kibiran, Toro, Uganda (17 Sept. 1905, on label). Collected and presented as the type by J. F. Jackson.
torquata—Pachycephala torquata Layard, 1875 39 Proc. zool. Soc. Lond., 1895: 150. SYNTYPE, Adult male. Reg. no. 1879.4.5.1835, Mountains of Taviuni, Fiji Islands. Collected by Lieut. Liardet. Salvin-Godman Collection. There are several other syntypes in the collection.
torquatus—Acridotheres torquatus Davison, 1892 62 Ibis, (6) 4:102. HOLOTYPE, Adult (in moult: marked as female on label). Reg. no. 1892.8.1.9. Paulu Tawer, Panang River, Malaya, 6 Aug. 1891. Presented by W. R. Davison of the Raffles Museum, Singapore.
torquatus—Gampsorhynchus torquatus Hume, 1874 321 Proc. Asiat. Soc. Bengal, 5:107. SYNTYPE, Adult (male on label). Reg. no. 1886.10.1.3090. (Younzaleen River, below the pine forests, Salween district, Tenasserim, vide Hume, Stray Feathers, 2 (1874): 446), (16 Feb. 1874. On label). Collected by W. Davison. Hume Collection.
torquatus—Lanius torquatus Latham, 1801

- Vol. 3, no. 32 in Lambert drawings, in library of Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 10 (1923): 393–394.
- torquatus—Microbates torquatus P. L. Sclater and Salvin, 1873 . 34 Nomencl. Av. Neotrop.,: 161. HOLOTYPE, Adult (female, vide Sclater, Cat. Bds. Brit. Mus., 15 (1890): 263). Reg. no. 1889.9. 20.398. St. George d'Oyapock, Cayenne, 1869. Collected by Verdey (in text, collected by Jelski, vide Sclater, Cat. Bds. Brit. Mus., 15 (1890): 263). Sclater Collection, obtained by him from Verreaux.
- torrentium—Motacilla clara torrentium Ticehurst, 1940 19
 Bull. Br. Orn. Cl., 60:81. Holotype, Adult male. Reg. no. 1905.12.29.1542. Ngoye Forest, Zululand, 8 Oct. 1904. Collected by C. H. B. Grant and presented by C. D. Rudd.

- torridus—Furnarius torridus P. L. Sclater & Salvin, 1866 . . . 3

 Proc. zool. Soc. Lond., 1866: 183. SYNTYPE, Adult male. Reg. no. 1889.5.20.34. Lower Ucayali River, eastern Peru. Collected by E. Bartlett. Sclater Collection.

 Listed as type by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 15.

toxopei—Acrocephalus stentoreus toxopei Hartert, 1924 35 Treubia, 6:20. HOLOTYPE, Adult male. Reg. no. 1910.12.28.266. Kayeli, Buru, Molucca Islands, Aug. 1909. Collected by W. Stalker. Presented by the British Ornithologists' Union Dutch New Guinea Expedition Committee.
transandeanus—Thamnophilus transandeanus P. L. Sclater, 1855. 4 Proc. zool. Soc. Lond., 1855: 18. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1841.2.483. Guayaquil, south-west Ecuador. Collected and presented by G. Barclay.
transkeiensis—Anthus richardi transkeiensis Vincent, 1948 19 Bull. Br. Orn. Cl., 69:17. HOLOTYPE, Adult male. Reg. no. 1948.64.1. (not 1948.44.1. as in text). Quimbu, 3,400', Transkeian Territories, eastern Cape Province, 18 Feb. 1947. Collected and presented by J. Vincent.
transvaalensis—Sphenoeacus transvaalensis C. H. B. Grant, 1908. 35 Bull. Brit. Orn. Cl., 21:92. Syntype, Adult male. Reg. no. 1905.12.29.2179. Woodbush Hills, 4,500', (Zoutpannsberg District, on label), north-east Transvaal, 11 Nov. 1905. Collected by C. H. B. Grant. Presented by C. D. Rudd. The female syntype is also in the collection.
transvaalensis—Sylvietta rufescens transvaalensis W. L. Sclater and Mackworth-Praed, 1918
travancoreensis—Machlolophus xanthogenys travancoreensis Whistler and Kinnear, 1932
travancoreensis—Ploceus philippinus travancoreensis Whistler, 1936 61 J. Bombay nat. Hist. Soc., 38:504. HOLOTYPE, Adult male. Reg. no. 1936.6.18.2. Kottayam backwaters, Travancore, 19 Feb. 1933. Collected by Salim Ali. Presented by the Travancore State Ornithological Survey.
travancoreensis—Pomatorhinus horsfieldi travancoreensis Fry, 1914 32B J. Bombay nat. Hist. Soc., 23:333. HOLOTYPE, Adult. Reg. no. 1910.5.11.145. Peermall, Travancore, south India. Collected and presented by W. Fry.
traylori—Cisticola aridula traylori Benson and Irwin, 1966 35 Arnoldia (Rhodesia), 2:1. HOLOTYPE, Adult male. Reg. no. 1931.12.21.166. Mocussueze, Moxico Province, Eastern Angola, 11°6′ S., 21°56′ E., 19 Jan. 1931. Collected by Admiral H. Lynes. Lynes Collection.
traylori—Nectarinia seimundi traylori. See appendix, Corrections and Additions, vol. 2

triborhyncha—Alauda triborhyncha Hodgson, 1843	
tricolor—Collurio tricolor Hodgson, 1837	3
tricolor—Digenea tricolor Hodgson, 1845	
tricolor—Geocichla tricolor Hume, 1871	1
tricolor—Hyphantornis tricolor Hartlaub, 1854 65 J. Orn. 1854: 110. New name for Ploceus collaris J. E. Gray, 1831 (q.v.) which is preoccupied by P. collaris Vieillot, 1819.	1
tricolor—Ixos tricolor Hartlaub, 1862	1
Proc. zool. Soc. Lond., (July) 1843: 4 (also in Ann. Mag. nat. Hist. (1) 13 (Dec. 1843): 441). Holotype, Adult female. Reg. no. 1847.1.18.56. Clarence, Fernando Po, West Africa, June. Collected by and purchased of L. Fraser. Received with other types by Gray, but not labelled as such.	3
Trans. Linn. Soc. Lond., 13:155. HOLOTYPE, Adult male. (relaxed mount). Reg. no. 1880.1.1.2094. Java (restricted to East Java, vide Robinson and Kloss, Treubia, 5 (1924): 294). Presented by the Honourable East India Company Museum as the type.	<u> </u>
ricolor—Phoenicura tricolor Hodgson, 1844	
Proc. zool. Soc. Lond., 1864: 45. HOLOTYPE, Adult. Reg. no. 1860.12.31.44. Tette, Zambesi. Collected on Dr. Livingstone's expedition by Dr. Kirk and presented by the Foreign Office through Lord Russell.	

trifasciatus—Orpheus trifasciatus Gould, 1837
trinitatis—Coereba trinitatis P. R. Lowe, 1907
trinitatis—Euphonia trinitatis Strickland, 1851
trinitatis—Spermophila trinitatis Sharpe, 1888 59 Cat. Bds. Brit. Mus., 12:132, 133. SYNTYPE, Adult. Reg. no. 1885.2.10.162. Trinidad. Sclater Collection; obtained by him from Argent. Listed as the type by Hellmayr, Catalogue of birds of the Americas, 11 (1938):212. The two other syntypes are also in the collection.
tristis—Brachypodius tristis Blyth, 1845
tristis—Leptopogon tristis P. L. Sclater and Salvin, 1876 9 Proc. zool. Soc. Lond., 1876: 254. Holotype, Immature (vide, Hellmayr, Novit. Zool., 13 (1906): 320). Reg. no. 1888.1.1.561. Simacu, Bolivia. Collected by C. Buckley. Salvin-Godman Collection.
tristis—Myiobius tristis Gosse, 1847
tristis—Thamnophilus tristis P. L. Sclater & Salvin, 1873

- tsanae—Calamoecetor leptorlyncha tsanae Bannerman, 1936. . 35
 Bull. Brit. Orn. Cl., 57:71. HOLOTYPE, Adult male. Reg. no.
 1927.11.5.331. Achera Mariam, 6,000', Lake Tana, Abyssinia,
 31 May 1926. Collected by and purchased of R. E. Cheesman.

There are three other syntypes in the collection, all four being listed as "authentic specimens" by Sclater, Cat. Bds. Brit. Mus., 15 (1890): 191. Pelzeln used Natterer's MS. name.

- tuckeri—Dendrocitta sinensis tuckeri Harrisson and Hartley, 1934 71 Bull. Brit. Orn. Cl., 54:156. HOLOTYPE, Adult male. Reg. no. 1933.11.2.1. Mount Dulit, 4,400', Sarawak, 10 Oct. 1932 (13 Sept. 1932, on label). Collected by T. H. Harrisson and C. H. Hartley. From the Oxford University Borneo Expedition.

- turipavae—Cichlornis whitneyi turipavae Cain and Galbraith, 1955 35 Bull. Brit. Orn. Cl., 75:91. HOLOTYPE, Adult male. Reg. no. 1959.21.558. Turipava, over 3,500', Guadalcanal, Solomon Islands, 15 Aug. 1953. Collected by a native hunter for the Oxford University Expedition to the Solomon Islands.

 The bill is broken.

turipavae—Zoothera margaretae turipavae Cain & Galbraith, 1955. 31 Bull. Br. Orn. Cl., 75:92. HOLOTYPE, Adult male. Reg. no. 1959.21.557. Turipava, 4,100', Guadalcanal, Solomon Islands, 18 Aug. 1953. Collected by a native hunter for the Oxford University Expedition to the Solomon Islands.
tweeddalei—Megalurus tweeddalei. See appendix, Corrections and Additions, vol. 2
typica—Plocealauda typica Hodgson, 1843
tyrannina—Pyriglena tyrannina P. L. Sclater, 1855
tyrannulus—Herpornis tyrannulus Swinhoe, 1870
tytleri—Calornis tytleri Hume, 1873
tytleri—Hirundo tytleri Jerdon, 1864
tytleri—Phylloscopus tytleri Brooks, 1871
ubendeensis—Turdus pelios ubendeensis Moreau, 1944 31 Bull. Br. Orn. Cl., 64:65. Holotype, Adult male. Reg. no. 1945.34.769. Lukolansala River, tributary of the Upper Nyamanse River, Ushamba, Ubende, western Tanganyika, 21 Nov. 1943. Collected and presented by R. E. Moreau.
ubiquitarius—Cichlops ubiquitarius Hodgson, 1844 19 in J. E. Gray, Zoological Miscellany, 1844: 83. Eleven specimens in the collection were listed as types by Sharpe, Cat. Bds. Brit. Mus., 10 (1885): 578, but the name is a nomen nudum.

ufipae—Alethe poliocephala ufipae Moreau, 1942 31 Bull. Br. Orn. Cl., 62:54. HOLOTYPE, Adult female. Reg. no. 1944.8.1.6. Mbisi Forest, 8,200', east of Sumbawanga, Tanganyika, 16 June 1941. Collected by Salimu Asmani for R. E. Moreau and presented by the latter.
ufipae—Dioptrornis fisheri ufipae Moreau, 1942
ufipae—Sathrocercus cinnemomea ufipae C. H. B. Grant and Mackworth-Praed, 1941
ugandae—Burnesia ugandae Sharpe, 1897
ugandae—Camaroptera superciliaris ugandae Clarke, 1914 35 Bull. Brit. Orn. Cl., 33: 136. HOLOTYPE, Adult. Reg. no. 1914.5. 14.1. Uganda. Collected and presented by Col. R. Stephenson Clarke.
ukambensis—Erythropygia ukambensis Sharpe, 1900 31 Bull. Br. Orn. Cl., 11:28. HOLOTYPE, Adult. Reg. no. 1900.6. 22.5. Ukambani, north-east Africa, Oct. 1894. Collected and presented by F. J. Jackson.
ulietensis—Turdus ulietensis Gmelin, 1788
ultima—Cercomela melanura ultima Bates, 1933
Bull. Br. Orn. Cl., 53: 175. HOLOTYPE, Adult female. Reg. no. 1932.8.6.9. Niger River, near Gao, French Sudan, 21 Aug. 1931 (10 Aug., on label). Collected by and purchased of G. L. Bates.
ultima—Elainea obscura ultima Griscom, 1935
Ibis, (13) 5:550. HOLOTYPE, Adult male. Reg. no. 1888.1.1.795. San Pedro Martyr, Guatemala, 22 Oct. 1873. Collected by O. Salvin. Salvin-Godman Collection.

- umbraticus—Horeites brunnifrons umbraticus E. C. S. Baker, 1924. 35 Bull. Br. Orn. Cl., 44:63. Holotype, Adult male. Reg. no. 1921.7.15.396. Shweli-Salwin Divide, 7,000–8,000′, West Central Yunnan, 25°40′ N., June 1919. Collected by G. Forrest. Presented by Col. Stephenson R. Clarke.

There are several other syntypes in the collection; the selected specimen was fully described and might be regarded as the holotype.

umbrosa—Thwingorhina striolata umbrosa Kloss, 1921 32B J. F. M. S. Mus., 10:212. HOLOTYPE, Adult male. Reg. no. 1936.4.12.141. Bandar Baroe, 800 metres, Upper Deli, north-east Sumatra, 21 Aug. 1920. Collected by F. C. van Heurn and presented by the Raffles Museum, Singapore.

The specimen mentioned by Gibson-Hill, Bull. Raffles Mus., 19 (1949): 147-8, as bearing a type label is now in the Brit. Mus. (Nat. Hist.). It is in fact the holotype of Stachyris poliocephala

pulla Kloss (q.v.).

- uncinatus—Lanius uncinatus P. L. Sclater and Hartlaub, 1881. 23

 Proc. zool. Soc. Lond., 1881: 168. SYNTYPE, Adult male. Reg.
 no. 1881.3.21.10. Socotra, 18 Feb. 1880. Collected by I. Bayley
 Balfour and presented by the Socotran Committee of the British
 Association.
- undulatus—Scytalopus undulatus Jardine and Selby, 1851 . . . Contributions to Ornithology: 117, pl. 76. HOLOTYPE, (Immature, vide Hellmayr on specimen label). Reg. no. 1886.6.24.514. No locality; 1838. Jardine Collection. Obtained by Jardine from Carfrae, an Edinburgh dealer.

- unicolor—Dysithamnus unicolor P. L. Sclater, 1859 Proc. zool. Soc. Lond., 1859: 141. SYNTYPE, Adult male. Reg. no. 1889.9.20.216. Pallatanga, west Ecuador, Oct. 1858. Collected by L. Fraser. Sclater Collection.

The few ile syntype is in the collection.

* 1 M * 1
 unicolor—Myiadestes unicolor P. L. Sclater, 1857 Proc. zool. Soc. Lond., 1856 (publ. 1857): 299. SYNTYPE, Adult. Reg. no. 1884.5.15.168. Near Cordova, Vera Cruz, Mexico. Collected by A. Sallé and purchased of P. L. Sclater. Listed as type by Hellmayr, Catalogue of Birds of the Americas, 7 (1934): 443. See also Salvin and P. L. Sclater, Exotic Birds, (1867): 51, pl. 26.
unicolor—Pnoepyga unicolor Hodgson, 1845 32B Proc. zool. Soc. Lond., 1845 : 25. Syntype (Immature vide Sharpe, Cat. Bds. Brit. Mus., 6 (1881) : 303) (relaxed mount). Reg. no. 1845.1.9.797. Nepal. Collected and presented by B. H. Hodgson.
 unicolor—Ramphocelus unicolor P. L. Sclater, 1856 51 Proc. zool. Soc. Lond., 1856: 128. SYNTYPE, Adult male. Reg. no. 1885.6.12.578. Bogota, Colombia. (Collected by Parzudaki, on label.) Sclater Collection. There are two other syntypes in the collection.
 unicolor—Scytalopus unicolor Salvin, 1895
unicolor—Turdus unicolor Tickell, 1833
uniformis—Chloropipo uniformis Salvin and Godman, 1884 8 Ibis, (5) 2:447. SYNTYPE, Adult male. Reg. no. 1888.1.20.16. Roraima, 3,500', British Guiana, (25 Jan. 1884, on label). Collected by H. Whitely. Salvin-Godman Collection. There are several other syntypes in the collection.
uniformis—Haplospiza uniformis P. L. Sclater and Salvin, 1873 . 49 Nomencl. Av. Neotrop.: 29, 157. Holotype, Adult. Reg. no. 1885.12.14.687. Jalapa, southern Mexico, 1872. Collected by De Oca. Salvin-Godman Collection.
unimodus—Graucalus unimodus P. L. Sclater, 1883
unitatis—Hirundo puella unitatis W. L. Sclater and Mackworth-Praed, 1918

uropygialis—Arachnothera uropygialis G. R. Gray and Mitchell, 1847 46

The Genera of Birds, 1: pl. 33. HOLOTYPE, Adult male (relaxed mount). Reg. no. 1846.11.21.15. (Indian Archipelago, vide Gadow, Cat. Bds. Brit. Mus., 9 (1884): 103). Purchased of Warwick.

Shelley, *Monograph of the Nectariniidae*, (1878): 368 confirms that the type is in the collection.

- uropygialis—Basileuterus uropygialis P. L. Sclater, 1861 54

 Proc. zool. Soc. Lond., 1861: 128. HOLOTYPE, Adult. Reg. no. 1884.5.15.453. Brazil (Altered to 'Panama' on label) see Todd, Proc. U.S. Nat. Mus., 74, art. 7 (1929): 18. Collected for Verreaux. Sclater Collection.
- uropygialis—Chrysomitris uropygialis P. L. Sclater, 1862 . . . 59
 Catalogue of a collection of American Birds, : 125. SYNTYPE, Adult.
 Reg. no 1888.2.10.690. Chili. Collected by T. Bridges. Sclater Collection.

The second syntype is also in the collection.

- uropygialis—Lathria uropygialis P. L. Sclater and Salvin, 1876 . 7

 Proc. zool. Soc. Lond., 1876: 355, pl. 32. Syntype, Adult male.

 Reg. no. 1888.1.13.1523. Tilotilo, 8,000', Yungas of La Pas, Bolivia,

 1876. (Collected by C. Buckley, on label). Sclater Collection.

 The three other syntypes are in the collection.
- uropygialis—Pheucticus uropygialis P. L. Sclater and Salvin, 1871.
 50 Proc. zool. Soc. Lond., 1870 (publ. 1871): 840. SYNTYPE, Adult. Reg. no. 1885.2.10.4. Bogota, Colombia, 1870. Sclater Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 56. There are several other syntypes in the collection.
- uropygialis—Ramphocelus uropygialis Bonaparte, 1851. 51
 Rev. Zool., 1851: 178. HOLOTYPE, Adult. Reg. no. 1885.6.12.
 563. Guatemala. Collected for Verreaux. Sclater Collection.
- uropygialis—Rectes uropygialis G. R. Gray, 1862 39

 Proc. zool. Soc. Lond., 1861 (publ. 1862): 430. SYNTYPE, Adult male. Reg. no. 1873.5.12.1291. Mysol Island. 1860. Collected by A. R. Wallace. Wallace Collection.

Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 8 (1883): 285.

urosticta—Formicivora urosticta P. L. Sclater, 1857 4 Proc. zool. Soc. Lond., 1857 : 130, pl. 126. SYNTYPE, Adult male. Reg. no. 1889.9.20.311. Brazil. Collected by Parzudaki. Sclater Collection. The other syntype is in the collection.
ursulae—Cyanomitra ursulae Boyd Alexander, 1903
usambara—Hyliota australis usambara Sclater, 1932 35 Bull. Brit. Orn. Cl., 52:104. HOLOTYPE, Adult male. Reg. no. 1932.1.11.1. Amani, 3,000', Tanganyika, 2 Sept. 1931. Collected by and purchased of R. E. Moreau.
usambarae—Sheppardia sharpei usambarae Macdonald, 1940 . 31 Ibis, (14) 4:669. HOLOTYPE, Adult male. Reg. no. 1931.12.19. 136. Amani Forest, 3,500', Usambara Mountains, Tanganyika, 1 Feb. 1931. Collected by R. E. Moreau and presented by the Salvin-Godman Fund.
usheri—Asthenes dorbignyi usheri Morrison, 1946
usheri—Pomatorhinus ruficollis usheri Hall, 1954
ussheri—Andropadus indicator ussheri Bannerman, 1920 21 Bull. Br. Orn. Cl., 41:6. HOLOTYPF, Adult. Reg. no. 1876.5.23. 627. Fantee, Gold Coast. Collected by Ussher. Sharpe Collection.
ussheri—Artomyias ussheri Sharpe, 1871
ussheri—Laniarius ussheri Sharpe, 1882
in Layard, Birds of South Africa, (new ed.): 397. HOLOTYPE, Adult. Reg. no. 1875.4.9.306. No locality (Volta River, Gold Coast, Aug. 1870, on label). Collected by Ussher. Sharpe Collection.
ussheri—Pitta ussheri Sharpe, 1877
ussheri—Pitta ussheri Sharpe, 1877

0,0
ussurianus—Cettia ussurianus Seebohm, 1881
ustulatus—Microcerculus ustulatus Salvin and Godman, 1883 . 28 Ibis, (5) 1: 204, pl. 9. Syntype, Adult male. Reg. no. 1885.3.6. 194. Roraima, British Guiana, 17 Nov. 1881. Collected by H. Whitely. Salvin-Godman Collection. The two other syntypes are in the collection.
utakwensis—Edoliisoma utakwensis Ogilvie-Grant, 1913 20 Bull. Br. Orn. Cl., 31: 105. HOLOTYPE, Adult, 'marked female but almost certainly a male' (vide Ogilvie-Grant, loc. cit.). Reg. no. 1911.12.20.1528. Camp 9, 5,500', Utakwa River, Dutch New Guinea, 30 Jan. 1913. Collected by C. B. Kloss and presented by the Committee of the British Ornithologists' Union Expedition to Dutch New Guinea.
utakwensis—Ptilotis salvadorii utakwensis Ogilvie-Grant, 1915 Ibis, Jubilee Supplement, 2:71, pl. 11. SYNTYPE, Adult male. Reg. no. 1916.5.30.579. Camp 11, 8,000', Utakwa River, Dutch New Guinea, (9 Feb. 1913, on label). Collected by C. B. Kloss. Presented by the Committee of the British Ornithologists' Union Expedition to Dutch New Guinea. This specimen was figured and named as the type male in the description, but the author marked another pair of syntypes as the types. The other three syntypes are in the collection.
vaalensis—Anthus vaalensis Shelley, 1900
valia—Cyrtostomus frenatus valia Mathews, 1929 40 Bull. Brit. Orn. Cl., 50:11. New name for C. f. olivaceus Mathews, 1926 (q.v.), on the grounds of preoccupation by Cinnyris olivaceus Smith, 1839.
validirostris—Lanius validirostris Ogilvie-Grant, 1894
validirostris—Sylvia validirostris G. R. Gray, 1869 3: Handlist of birds, 1:216. This Jerdon MS. name is a nomen nudum. The type listed by Seebohm, Cat. Bds. Brit. Mus., 5 (1881):51, has therefore no type status.

Zur Ornithologie Brasiliens: 95, 169. SYNTYPE, Adult male. Reg. no. 1888.1.13.1573. Villa Maria, Matogrosso, Brazil, 28 Aug. Collected by J. Natterer. Sclater Collection.
Listed as 'authentic specimen' by P. L. Sclater, Cat. Bds. Brit. Mus., 14 (1888): 364.
vansomereni—Erythropygia leucophrys vansomereni W. L. Sclater, 1929
vanuensis—Merula vanuensis Layard, 1890
variegata—Elaenia variegata P. L. Sclater, 1857
variegata—Eopsaltria variegata G. R. Gray, 1859 39 Proc. zool. Soc. Lond., 1859: 162. SYNTYPE, Adult male. Reg. no. 1859.4.19.10. Island of Nu, Port de France, New Caledonia. (8 May 1858, on label). Collected by J. Macgillivray. Purchased of H. Cuming. The female syntype is also in the collection.
variegaticeps—Anabazenops variegaticeps P. L. Sclater, 1857 3 Proc. zool. Soc. Lond., 1856 (publ. 1857): 289. SYNTYPE, Adult. Reg. no. 1889.5.20.382. Cordova, Vera Cruz, southern Mexico. Collected by A. Sallé. Sclater Collection. Listed as type by Sclater, Catalogue of American Birds, (1862): 159. There is another syntype in the collection.
variegatus—Camarhynchus variegatus P. L. Sclater and Salvin, 1870. 53 Proc. zool. Soc. Lond., 1870: 324. Syntype, Adult male. Reg. no. 1885.12.14.542. Abingdon Island, Galapagos Archipelago, (30 Dec. 1868, on label). Collected by A. Habel. Salvin-Godman Collection. There are several other syntypes in the collection.
variegatus—Schoeniparus variegatus Styan, 1899

venningi—Turdinulus brevicaudatus venningi Harington, 1913 . 32B Bull. Br. Orn. Cl., 33:44. HOLOTYPE, Adult male. Reg. no. 1903.12.24.181. Pinlikum, Lorlong State, South Shan States, Burma, 23 March 1902. Collected by W. H. Craddock. Craddock and Thompson Collection; purchased of Gerrard.

ventralis—Thamnophilus ventralis P. L. Sclater, 1855 Edinb. New philos. J., (n.s.) 1:244. HOLOTYPE, Adult male. Reg. no. 1889.9.20.95. Southern Brazil, 1852. Collected for Argent. Sclater Collection. venusta—Calliste venusta P. L. Sclater, 1855 Proc. zool. Soc. Lond., 1854 (publ. 1855): 248. SYNTYPE, Adult. Reg. no. 1885.6.12.384. Bogota, Colombia, 1854. Collected for Argent. Sclater Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 133. There is another syntype in the collection. venustulus—Parus venustulus Swinhoe, 1870. Proc. zool. Soc. Lond., 1870:133. HOLOTYPE, Adult. Reg. no. 1898.9.20.157. Kweifoe, Szechuen, May 1869. Collected by R. Swinhoe. Seebohm Bequest. venustus—Artamus venustus Sharpe, 1878 in G. D. Rowley, Ornithological Miscellany, 3:198. SYNTYPE, Adult female. Reg. no. 1857.10.28.33. North-western Australia (Victoria River, Northern Territory, 1856, on label). Collected and presented by J. R. Elsev. There are several syntypes in the collection. vepres—Turdoides melanops vepres R. Meinertzhagen, 1936 Bull. Br. Orn. Cl., 57:69. HOLOTYPE, Adult male. Reg. no. 1965.M.40. Nanyuki, 7,000', Kenya, 29 Jan. 1936. Collected by R. Meinertzhagen. Meinertzhagen Collection. veraepacis—Automolus veraepacis Salvin and Godman, 1891. Biol. Centr.-Amer.: Aves, 2:156. SYNTYPE, Adult. Reg. no. 1889.5.14,329. Coban, Vera Paz, Guatemala, Nov. 1859. Collected by O. Salvin. Salvin-Godman Collection. There are several other syntypes in the collection. veraepacis—Heteropelma veraepacis P. L. Sclater and Salvin, 1860 . 8 Proc. zool. Soc. Lond., 1860: 300. SYNTYPE, Adult. Reg. no. 1888.1.20.310. (Choctum, on label), Vera Paz, Guatemala, (Jan. 1860, on label). Collected by O. Salvin. Salvin-Godman Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 321. There are several other syntypes in the collection. veraguensis—Basileuterus veraguensis Sharpe, 1885 Cat. Bds. Brit. Mus., 10: 403. HOLOTYPE, Adult. Reg. no. 1868.2, 17.55. Paraiso Station, Panama, 1867. Collected by A. Hughes. Purchased of O. Salvin. vermiculata-Stachyris larvata vermiculata T. H. Harrisson and Hartley, 1934 Bull. Br. Orn. Cl., 54:153. HOLOTYPE, Adult male. Reg. no.

1933.11.2.5. Mount Dulit, 4,000', Sarawak, Borneo, 29 Aug.

- 1932. Collected by T. H. Harrisson & C. H. Hartley and presented by the Oxford University Sarawak Expedition.
- vernayi—Cyornis poliogenys vernayi Whistler, 1931 37

 Bull. Brit. Orn. Cl., 52: 24. HOLOTYPE, Adult male. Reg. no. 1931.10.7.1. Sankrametta, 3,500', Vizagapatam District, India, 12 March 1930. Collected by H. Whistler. Presented by the Bombay Natural History Society.
- vernayi—Dendrocitta rufa vernayi Kinnear and Whistler, 1930 . 71 Bull. Brit. Orn. Cl., 51:17. HOLOTYPE, Adult male. Reg. no. 1937.12.21.46. Nallamalai Range, 2,000', south Kurnool, South India, 1 Nov. 1929. Collected by V. S. La Personne. Presented by the Bombay Natural History Society.
- vernayi—Leucocirca pectoralis vernayi Whistler, 1931 37A Bull. Brit. Orn. Cl., 52:40. Holotype, Adult male. Reg. no. 1931.10.7.2. Jeypore Agency, 3,000′, (2,000′, on label), Upper Eastern Ghats, India, 26 April 1930. Collected by V. S. La Personne. Presented by the Bombay Natural History Society.
- vernayi—Uroloncha kelaarti vernayi Whistler and Kinnear, 1933 . 60 J. Bombay nat. Hist. Soc., 36: 835. Holotype, Immature female. Reg. no. 1937.12.21.525. Vizagapatam, 3,500', Sankrametta, East Ghat Mountains, India, 17 March, 1930. Collected and presented by V. S. La Personne.
- verneyi—Tephrodornis pelvicus verneyi Kinnear, 1924 20 Bull. Br. Orn. Cl., 44: 102. HOLOTYPE, Adult male. Reg. no. 1924.12.22.91. Um Pang, 1,500', Siam, 25 Jan. 1924. Collected by W. P. Lowe and A. S. Vernay and presented by the latter.

of uncertain status (fide W. E. China, in litt., Z.N. (G) 34). The specimens described by Swainson (loc. cit.) are in the University Museum of Zoology, Cambridge (C. W. Benson, pers. comm.), and may be regarded as syntypes.

This specimen was labelled as the type by F. D. Godman and listed as such by Sharpe, *Cat. Bds. Brit. Mus.*, 10 (1885): 408, but other specimens were available and no type is specified in the description. There are other syntypes in the collection.

Vol. 1, no. 5 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). The author, in *General Synopsis of birds*, Suppl., 2 (1802): 117, refers to Lambert. See also Mathews, Birds of Australia, 10 (1923): 423.

There are four other syntypes at the Academy of Natural Sciences, Philadelphia.

- verticalis—Monarcha verticalis P. L. Sclater, 1877 38

 Proc. zool. Soc. Lond., 1877: 99, pl. 14. HOLOTYPE, Adult. Reg.
 no. 1887.12.30.1188. Duke of York Island, Oct. 1875. Collected
 by the Rev. G. Brown. Tweeddale Collection.

victoriae—Babax victoriae Rippon, 1905
victoriae—Certhia victoriae Rippon, 1906
victoriae—Ianthocincla victoriae Rippon, 1906 32B Bull. Br. Orn. Cl., 16: 47. SYNTYPE, Adult. Reg. no. 1905.9.10. 65. Mount Victoria, 8,000', South Chin Hills, Burma, 26 March 1904. Collected and presented by G. Rippon. There are several other syntypes in the collection.
victoriae—Menura victoriae Gould, 1865
victoriae—Ptiloris victoriae Gould, 1849
victoriae—Pyrrhula victoriae Rippon, 1906
victoriae—Sitta victoriae Rippon, 1904

victoriae—Siva strigula victoriae R. Meinertzhagen, 1926 32B Bull. Br. Orn. Cl., 46: 128. HOLOTYPE, Adult. Reg. no. 1905.9. 10.325. (not 1903.9.10.325. as in text). Mount Victoria, 7,000', Chin Hills, Burma, 26 March 1904. Collected and presented by Col. G. Rippon.
vieilloti—Calliste vieilloti P. L. Sclater, 1857
vieirae—Parus vieirae Nicholson, 1906
vigorsi—Ruticilla vigorsi Moore, 1854
vigorsii—Cinnyris vigorsii Sykes, 1832
vigorsii—Pitta vigorsii Gould, 1838
vilissima—Elainia vilissima P. L. Sclater and Salvin, 1859 9 Ibis, (1) 1:122, pl. 4. SYNTYPE, Adult. Reg. no. 1859.6.29.21. Guatemala. Collected by G. U. Skinner. Purchased of Gould. There are several other syntypes in the collection. The specimen listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 132, was collected after the date of the original description and has no type status.
villosus—Myiobius villosus P. L. Sclater, 1860 9 Proc. zool. Soc. Lond., 1860: 93. SYNTYPE, Adult (female, vide Sclater, Cat. Bds. Brit. Mus., 14 (1888): 201). Reg. no. 1888.1.13.

829. Nanegal, Ecuador, Feb. 1859. Collected by L. Fraser. Sclater Collection. The other syntype is also in the collection.
vincens—Prionchilus vincens P. L. Sclater, 1872
vincenti—Cyanomitra olivacea vincenti C. H. B. Grant and Mackworth-Praed, 1934
vincenti—Fringillaria capensis vincenti P. R. Lowe, 1932
vincenti—Phyllastrephus flavostriatus vincenti C. H. B. Grant and Mackworth-Praed, 1940
vincenti—Pirenestes vincenti Benson, 1955
vinipectus—Siva vinipectus Hodgson, 1838
vinitincta—Merula vinitincta Gould, 1855
violacea—Tchitrea plumbiceps violacea C. H. B. Grant and Mackworth-Praed, 1940

virata—Lagonosticta rubricata virata Bates, 1932.	60
Bull. Brit. Orn. Cl., 53:7. HOLOTYPE, Adult female.	Reg. no.
1930.3.4.331. Kulikoro, French Sudan, 25 June 1928.	Collected
by and purchased of G. L. Bates.	

- virens—Phylloscopus maculipennis virens Ticehurst, 1926 35 Bull. Brit. Orn. Cl., 64:61. HOLOTYPE, Adult male. Reg. no. 1926.1.20.1. Banjar, 4,500', Saraj, Punjab Himalayas, 15 Nov. 1923. Collected and presented by H. Whistler.
- virenticeps—Cycloris virenticeps P. L. Sclater, 1860 56 Proc. zool. Soc. Lond., 1860: 274, pl. 164. SYNTYPE, Adult. Reg. no. 1886.9.15.366. Babahoyo, 200 miles north east of Guayaquil, Ecuador, July 1859. Collected by L. Fraser. Sclater Collection. There is a second syntype in the collection.
- virescens—Calliste virescens P. L. Sclater, 1851 in Jardine, Contributions to Ornithology Jan. 1851: 22, 56, pl. 69. SYNTYPE, Adult male, Reg. no. 1885.6.12.256. Cavenne. Collected for Argent. Sclater Collection. Listed as the type by Sclater, Catalogue of a collection of American birds. (1862): 65.
- virescens—Dicaeum virescens Hume, 1873 45 Stray Feathers, 1:482. SYNTYPE, Adult male. Reg. no. 1887.2.1. 295. Port Blair, South Andaman Islands, 17 July 1873. Collected for A. O. Hume. Hume Collection. There are three other syntypes in the collection.
- virescens—Ptilotis virescens Wallace, 1864 Proc. zool. Soc. Lond., 1863 (publ. 1864): 494. SYNTYPE, Adult. Reg. no 1873.5.12.1036. Lombok, Lesser Sunda Islands, 1856. Collected by A. R. Wallace. Wallace Collection.

Renamed *Meliphaga I. lombokia* Mathews, 1926, on the grounds of preoccupation by Melithreptus virescens Vieillot, 1817. This specimen is listed as the type by Gadow, Cat. Bds. Brit. Mus., 9 (1884): 494. There are several other syntypes in the collection.

viridescens—Calornis viridescens G. R. Gray, 1858 62 Proc. zool. Soc. Lond., 1858: 181. SYNTYPE, Adult female. Reg. no. 1858.3.10.90. Aru Island. Collected by A. R. Wallace. Wallace Collection.

There are other syntypes in the collection. This is the only specimen labelled in Gray's handwriting.

viridescentior—Pycnonotus viridescentior Sharpe, 1904. *Ibis*, (8) 4:638. HOLOTYPE, Adult male. Reg. no. 1903.7.16.75. River Ja, Cameroons hinterland, 16 Jan. 1903. Collected by G. L. Bates and purchased of W. E. de Winton.

Labelled as the type by the author. There is a second syntype in the collection.

- viridiceps—Rhynchocyclus viridiceps P. L. Sclater and Salvin, 1873. 9

 Proc. zool. Soc. Lond., 1873: 280. HOLOTYPE, Adult male. Reg.
 no. 1888.1.13.722. Pebas, (Lareto), east Peru, 28 May 1866.
 Collected by J. Hauxwell. Sclater Collection.

Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 32. The

female syntype is also in the collection.

- virgatum—Trochalopteron virgatum Godwin-Austen, 1874 . . . 32B Proc. zool. Soc. Lond., 1874 : 46. HOLOTYPE, Adult male. Reg. no. 1895.7.14.2255. Rázami, 5,000', under the Kopamedza ridge, north-east frontier of Bengal. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.
- viridicollis—Calliste argentea viridicollis Taczanowski, 1884 . . 5:

 Ornithologie du Pérou, 2:468. Syntype, Adult male. Reg. no.
 1885.6.4.627. Huiro, 4,800′, Valley of Urubamba, Peru. (Collected by J. Orton, on label.) Salvin-Godman Collection.

 Taczanowski used a Salvin MS. name. The female syntype is also in the collection.
- viridinitens—Myiagra viridinitens G. R. Gray, 1859 38

 Proc. zool. Soc. Lond., 1859: 162. HOLOTYPE, Adult male. Reg.
 no. 1854.5.31.9. Loyalty Island, South West Pacific. Presented by Sir G. Grey.
- viridior—Dicrurus forficatus viridior Salomonsen, 1934. 64

 Proc. zool. Soc. Lond., 1934: 223. HOLOTYPE, Adult male. Reg.
 no. 1931.8.18.3055. Tabiky, western Madagascar, 29 Oct. 1913.
 Collected by A. L. Rand and R. A. Archbold. Archbold-Vernay Expedition.

lected by Sir S. Raffles and presented by the Honourable East India Company Museum.
viridis—Cochoa viridis Hodgson, 1836
viridis—Eroessa viridis Sharpe, 1883
viridis—Tatare? viridis Layard, 1875
viridis—Turdus viridis Horsfield, 1821
viridissimus—Tyranniscus viridissimus P. L. Sclater, 1874 9 Proc. zool. Soc. Lond., 1873 (publ. 1874): 782. SYNTYPE, Adult. Reg. no. 1888.1.13.500. Cosnipata, 2,300', Cuzco, south-east Peru, (July 1871, on label). Collected by H. Whitely. Sclater Collection. Listed as the type by Sclater, Cat. Bds. Brit. Mus., 14 (1888): 135. The second syntype is also in the collection.
viridogrisea—Aplonis viridogrisea G. R. Gray, 1859 62 Proc. zool. Soc. Lond., 1859: 164. SYNTYPE, Adult female. Reg. no. 1859.4.19.2. Port de France, Island of Nu, New Caledonia, 14 May 1858. Collected by J. McGillivray. Purchased of Cummings. Listed as type by Sharpe, Cat. Bds. Brit. Mus., 13 (1890): 128. There are two other syntypes in the collection.
virussu—Lipaugus virussu Pelzeln, 1869
vitellina—Pipra vitellina Gould, 1843

Reg. no. 1898.11.10.165. (Wakaia, on label), Fiji Islands, (13 Sept. 1875, on label). Collected by E. L. Layard. Seebohm Bequest.
The above specimen is labelled as a type. The described male, from Taviuni, cannot be traced; there are other syntypes in the collection.
vitiensis—Merula vitiensis E. L. Layard, 1876
vitiensis—Pachycephala vitiensis G. R. Gray, 1859 39 Catalogue of the birds of the tropical islands of the Pacific : 20. SYNTYPE, Adult male. Reg. no. 1856.8.30.35. Island of Ngau, Fijis. Collected by T. M. Rayner (queried by Mayr, Amer. Mus. Nov., no. 531 (1932) : 13). From the voyage of H.M.S. Herald. The female syntype is also in the collection.
vittata—Euphonia vittata P. L. Sclater, 1861
vivada—Cyornis vivada Swinhoe, 1864
vivada—Niltava vivada Hume, 1874
vivax—Trochocercus vivax Neave, 1909

vivida—Xanthiscus flavescens vivida E. C. S. Baker, 1917 . . . 21

Bull. Brit. Orn. Cl., 38:16. SYNTYPE, Adult male. Reg. no.
1886.9.1.2018. Salween, Burma, 11 Feb. 1874. Collected by W. Davison. Hume Collection.

This is the author's selected male. His selected female syntype is also in the collection.

- vividus—Pericrocotus peregrinus vividus E. C. S. Baker, 1920 . 20

 Bull. Brit. Orn. Cl., 40: 114. HOLOTYPE, Adult male. Reg. no. 1886.4.1.590. Attaran River, Burma, 9 Dec. 1878. Collected by W. Davison. Hume Collection

Presented as the type and listed as such by Lynes, *Ibis*, (12) 6, *Supplement*, (1930): 671, but other specimens are mentioned in the

description.

This is the described male syntype. The described female syntype

is also in the collection.

Purchased as the type. There is another syntype in the collection.

vulpecula—Calandrella rufescens vulpecula C. M. N. White, 1955 . 17 Bull. Br. Orn. Cl., 75:3. HOLOTYPE, Adult male. Reg. no. 1954. 56.1. Ten miles west of Bohotleh, 2,300', 8°12' N., 46°10' E., British Somaliland, 9 Feb. 1954. Collected and presented by J. G. Williams.

- vulpecula—Synallaxis vulpecula P. L. Sclater and Salvin, 1866 . 3

 Proc. zool. Soc. Lond., 1866: 184. SYNTYPE, Adult. Reg. no. 1889.5.20.190. Upper Ucayali River, east Peru, June 1865. Collected by E. Bartlett. Sclater Collection.

 There are two other syntypes in the collection.

 waddelli—Garralux waddelli Ogilvie-Grant 1894 . 32B
- waddelli—Garralux waddelli Ogilvie-Grant, 1894 32B Bull. Br. Orn. Cl., 3:29. HOLOTYPE, Adult female. Reg. no. 1895.6.30.1. Rangit River, 4,000′, Sikkim, April 1891. Collected and presented by L. A. Waddell.
- wagleri—Cacicus wagleri G. R. Gray and Mitchell, 1844 57 Genera of Birds, 2:342, pl. 85. HOLOTYPE, Adult male. Reg. no. 1845.6.13.16. (Coban, on label = lowlands of eastern Guatemala, vide Van Rossem, Bull. Mus. Comp. Zool. Harv., 77 (1934):405). Purchased of Leadbeater.

This specimen identified by W. L. Sclater (MS. notes) as one of a pair available for Gray's description and probably collected by Delattre who was in Guatemala in 1842. The only description is a plate and the identity of the holotype was established by Van Rossem (op. cit., 404-5).

Listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 373. There is another syntype in the collection, and others at the Smithsonian Institution.

- wakoloensis—Myzomela wakoloensis Forbes, 1883 . . . 4

 Proc. zool. Soc. Lond., 1883: 116. SYNTYPE, Adult male. Reg.
 no. 1884.5.12.24. Lake Wakolo, Bouru, Ceram group. Collected
 by H. O. Forbes. Purchased of E. W. Janson, from the Timor
 Laut Expedition (vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 402).
 Listed as the type by Sharpe, op. cit. The female and immature
 male syntypes are in the collection.
- waldeni—Actinodura waldeni Godwin-Austen, 1874 321

 Proc. zool. Soc. Lond., 1874 : 46, pl. 12. Syntype, Adult (female on label). Reg. no. 1895.7.14.1415. Japvo Peak, 9,000', Burrail Range, Naga Hills, north-east Bengal. Collected by H. H. Godwin-Austen. Godwin-Austen Collection.

There is a second syntype in the collection.

waldeni—Lanius waldeni Swinhoe, 1870
Reg. no. 1898.9.20.685. Fungtoo Hien, Yangtsze River, Szechuen, China, 5 May 1869. Collected by R. Swinhoe. Seebohm Bequest. There are several other syntypes in the collection.
waldroni—Anthoscopus flavifrons waldroni Bannerman, 1935. 41 Bull. Brit. Orn. Cl., 55: 131. HOLOTYPE, Adult (?male, on label). Reg. no. 1935.7.17.97. Goaso, Ashanti, Gold Coast. 15 Dec. 1934. Collected and presented by W. P. Lowe and Miss F. Waldron.
wallacei—Buchanga wallacei Walden, 1870
wallacei—Dendrocincla fuliginosa wallacei Chubb, 1919 2 Bull. Br. Orn. Cl., 39: 52. Syntype, Adult male. Reg. no. 1889. 5.20.628. Para, north-east Brazil, May 1849. Collected by A. R. Wallace. P. L. Sclater Collection. The female syntype described is also in the collection.
wallacei—Dissemurus paradiseus wallacei Hachisuka, 1926 64 Bull. Brit. Orn. Cl., 47:58. HOLOTYPE, Adult. Reg. no. 1873.5. 12.1984. Modjokerto, eastern Java, Aug. 1861. Collected by A. R. Wallace. Wallace Collection.
wallacei—Pipromorpha oleaginea wallacei Chubb, 1919 9 Ann. Mag. nat. Hist., (9) 4:301. Holotype, Adult. Reg. no. 1888.1.1.536. Para, Lower Amazon Valley, May 1849. Collected by A. R. Wallace. Salvin-Godman Collection.
wallacei—Zosterops wallacei. See appendix, Corrections and Additions, vol. 2
wallacii—Heteropelma wallacii P. L. Sclater and Salvin, 1867 8 Proc. zool. Soc. Lond., 1867 : 579. HOLOTYPE, Adult female. Reg. no. 1888.1.13.1364. Para, Lower Amazon, Brazil, May 1849. Collected for A. R. Wallace. Sclater Collection.
wallacii—Munia wallacii Sharpe, 1890
wallacii—Paradisea wallacii G. R. Gray, 1859
at that period, were published in groups of sheets, pages 129 to 352

allacii—Todopsis wallacii G. R. Gray, 1861
There are two other syntypes in the collection.
alleri—Amydrus walleri Shelley, 1880
altoni—Propasser waltoni Sharpe, 1905
ambera—Cisticola brunnescens wambera Lynes, 1930 35 Ibis, (12) 6, Supplement: 162. HOLOTYPE, Adult male. Reg. no. 1927.11.5.474. Wambera Town, 8,000', South-west Abyssinia, (error for North-West Abyssinia), 15 April 1927. Collected by R. E. Cheesman. Presented by Admiral Lynes.
ardi—Actinodura nipalensis wardi Kinnear, 1932
ardi—Pseudobias wardi Sharpe, 1870
ardii—Turdus wardii Blyth, 1842

- warsangliensis—Tschagra senegala warsangliensis Clarke, 1919 . 23
 Bull. Br. Orn. Cl., 40:50. HOLOTYPE, Adult male. Reg. no.
 1923.8.7.1268. Mush Haled, 4,000', Warsangli country, eastern
 British Somaliland, 4 March 1919. Collected by Sir G. F. Archer
 and presented by Col. Stephenson Clarke.

This is the author's selected male syntype. His selected female syntype and five other syntypes are also in the collection.

The above specimen is the one from which the colours of bill and feet, the locality, and the date were used for the type description. The collector's number and statement that the type was in the museum refer, however, to another specimen, reg. no. 1911.11.22.1., collected at Lake Ngeyi, Mufumbiro, on 14 Feb. 1911, and pre-

- sented by Fox; this specimen has a broken culmen. It would appear that Ogilvie-Grant used both specimens for the description, but listed only the syntype held by the museum at that time.
- wellsi—Parus dichrous wellsi E. C. S. Baker, 1917 42

 Bull. Brit. Orn. Cl., 38:8. HOLOTYPE, Adult. Reg. no. 1906.12.

 17.397. Yangtze River, big bend, 9-10,000', West Yunnan, March 1906. Collected and presented by G. Rippon.
- westermanni—Musicapula westermanni Sharpe, 1888 37

 Proc. zool. Soc. Lond., 1888 : 270. Holotype, Adult male. Reg.
 no. 1888.3.18.41. Gunong Ulu Batang Padang, 4,200', Perak,
 Malaya. Collected and presented by L. Wray.
- wetmorei—Passerculus sandwichensis wetmorei van Rossem, 1938 . 49
 Bull. Brit. Orn. Cl., 58: 129. HOLOTYPE, Adult male. Reg. no.
 1899.2.1.2893. Chancol, 10,000'., Huehuetenango, Guatamala,
 17 June 1897. Collected by W. B. Richardson. Salvin-Godman
 Collection.
- whistleri—Garrulax albogularis whistleri E. C. S. Baker, 1921 . 32B Bull. Br. Orn. Cl., 42:29. HOLOTYPE, Adult male. Adult male. Reg. no 1886.10.1.3841. Simla, India, 3 Nov. 1880. Hume Collection.
- whistleri—Horeites brunneifrons whistleri Ticehurst, 1923 . . . 35

 Bull. Brit. Orn. Cl., 44: 28. Holotype, Adult male. Reg. no. 1886.7.8.2196. Simla, north-west Himalayas, 24 Nov. 1880. Collected by W. Davison. Hume Collection.
- whistleri—Niltava sundara whistleri Ticehurst, 1926 37
 Bull. Brit. Orn. Cl., 46: 113. HOLOTYPE, Adult female. Reg. no. 1891.7.22.40. Murree, North-west Himalaya, 10 July 1873. Collected by F. Stoliczka. Received in exchange with the Indian Museum, Calcutta, as duplicates of the Second Yarkand Mission under Sir D. Forsyth, 1873-4.
- whistleri—Seicercus burkii whistleri Ticehurst, 1925 3

 Bull. Brit. Orn. Cl., 46: 22. HOLOTYPE, Adult male. Reg. no. 1926.1.4.1. Dharmsala, Punjab Himalaya, 26 March 1922. Collected and presented by H. Whistler.

whitakeri—Galerida cristata whitakeri Bannerman, 1922. Bull. Br. Orn. Cl., 42:124. HOLOTYPE, Adult male. Reg. no. 1922.5.1.19. Tatahouine, south-east Tunisia, Oct. 1899. Collected by M. Blanc and presented by J. I. S. Whitaker.
whiteheadi—Cinnyris whiteheadi Ogilvie-Grant, 1894 46 Bull. Brit. Orn. Cl., 3:50 Syntype, Adult male. Reg. no. 1897.5.13.44. La Trinadad, Benguet District, North-west Luzon, Philippines, 25 Feb. 1894. Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition. A female syntype is also in the collection.
whiteheadi—Dicaeum haematostictum whiteheadi Hachisuka, 1926 45 Bull. Brit. Orn. Cl., 47:55. HOLOTYPE, Adult male. Reg. no. 1897.5.13.441. Mt. Canloan, Negros, Philippines, 26 March 1896. Collected by J. Whitehead. Presented by the subscribers to the Whitehead Expedition.
whiteheadi—Merula whiteheadi Seebohm, 1893 31 Bull. Br. Orn. Cl. 1: 25. SYNTYPE, Adult male. Reg. no. 1896.6. 1.1012. Near Tozari, 7,000', east Java, 26 Sept. 1886. Collected by J. Whitehead. Seebohm Collection. The three other syntypes are also in the collection.
whiteheadi—Oreocincla whiteheadi E. C. S. Baker, 1913 31 Bull. Br. Orn. Cl., 31:79. SYNTYPE, Adult male. Reg. no. 1913.10.15.1. Khagan Valley, 13,000', Afridi Country, Northwest Frontier Province, 15 July 1912. Collected and presented by C. H. T. Whitehead. The juvenile syntypes are also in the collection.
whiteheadi—Sitta whiteheadi Sharpe, 1884
whiteheadi—Urocissa whiteheadi Ogilvie-Grant, 1899 71 Bull. Brit. Orn. Cl., 10:18. SYNTYPE, Adult male. Reg. no. 1900.1.2.141. Five Finger Mountains, Hainan, China, 17 April 1899. Collected by J. Whitehead. Purchased of E. Gerrard; from the Whitehead Expedition. The female syntype is also in the collection.
whiteheadi—Zosterornis whiteheadi Ogilvie-Grant, 1894 32B Bull. Br. Orn. Cl., 3:50. SYNTYPE, Adult male. Reg. no. 1897.5. 13.83. La Trinidad, Benguet District, North-west Luzon, Philippines, 11 Jan. 1894. Collected by J. Whitehead and presented by the Subscribers to the Whitehead Expedition. The two other syntypes are in the collection.

Bull. Brit. Orn. Cl., 57: 114. New name for A. m. bensoni C. H. B. Grant and Mackworth-Praed, 1937 (a.v.), on the grounds of preoccupation by A. bamerdae bensoni Vincent, 1935.

whitei—Cyornis whitei Harington, 1908 37 Ann. Mag. nat. Hist., (8) 2: 245. SYNTYPE, Adult male. Reg. no. 1908.8.2.26. Watan, Bhamo Hills, Burma, 3 April 1908. Collected and presented by H. H. Harington.

The two female syntypes are also in the collection.

whitelevana—Spermophila whitelevana Sharpe, 1888 Cat. Bds. Brit. Mus., 12:18. SYNTYPE, Adult male. Reg. no. 1885.12.14.523. Roraima (3,500', on label), British Guiana, 21 Nov. 1883. Collected by H. Whitely. Salvin-Godman Collec-

This specimen is marked on the label as the one described, and listed as the type by Hellmayr, Catalogue of Birds of the America, 11(1938): 179. The six other syntypes are also in the collection.

- whitelevi—Chelidon whitelevi Swinhoe, 1862 18 Proc. zool. Soc. Lond., 1862: 320. HOLOTYPE, Adult. 1898.10.20.40. Near Pekin. Collected by R. A. Fleming. ceived by Swinhoe from Whitely. Seebohm Bequest.
- whitelyanus—Atalotriccus griseiceps whitelyanus Chubb, 1919. Ann. Mag. nat. Hist., (9) 4:301. SYNTYPE, Adult male. Reg. no. 1892.1.16.259. Quonga, (1,000', on label), British Guiana, 22 Oct. 1887. Collected by H. Whitely Jnr. Salvin-Godman Collection.

There are two other syntypes in the collection.

whitelyi—Calliste whitelyi Salvin and Godman, 1884 51 Ibis. (5) 2:445, pl. 13. SYNTYPE, Adult male. Reg. no. 1885.6. 4.539. Forests on the slopes of Roraima, 5,000-6,000', British Guiana, 7 Sept. 1883. Collected by H. Whitely. Salvin-Godman Collection.

There are several other syntypes in the collection.

whitelyi—Elaenia cristata whitelyi Chubb, 1919 Ann. Mag. nat. Hist., (9) 4:304. SYNTYPE, Adult male. Reg. no. 1888.1.1.672. Mount Roraima, 28 Nov. 1881. Collected by H. Whitely. Salvin-Godman Collection.

The reference by Chubb, Birds of British Guiana, 2 (1921): 190, to specimens used in descriptions would appear to relate to that work

only. There are several other syntypes in the collection.

whitelyi-Pipreola whitelyi Salvin and Godman, 1884 Ibis. (5) 2:449. HOLOTYPE, Adult female. Reg. no. 1888.1.20.788. Roriama, 6,000', British Guiana, 7 Sept. 1883. Collected by H. Whitely. Salvin-Godman Collection.

whitii—Poospiza whitii P. L. Sclater, 1883
Proc. zool. Soc. Lond., 1883: 43, pl. 9. SYNTYPE, Adult male. Reg. no. 1885.2.10.321. Cosquin, 30 miles from Cordova, Argentine, 6 July 1882. Collected by E. W. White. Sclater Collection. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 12 (1888): 642. There are several other syntypes in the collection.
whitii—Zonotrichia whitii Sharpe, 1888
whitmeei — Lalage whitmeei Sharp, 1878
whymperi—Laiscopus collaris whymperi E. C. S. Baker, 1915 30 Bull. Br. Orn. Cl., 35:61. HOLOTYPE, Adult female. Reg. no. 1915.4.28.3. Nita Valley, 15,000', Garhwal, India, 4 July 1910. Collected by S. L. Whymper and presented by the Bombay Natural History Society.
whytei—Cinnyris afer whytei Benson, 1948
whytii—Serinus whytii Shelley, 1897
whytii—Sylviella whytii Shelley, 1894
wilkinsi—Graucalus papuensis wilkinsi Kinnear, 1924 20 Bull. Br. Orn. Cl., 44:68. HOLOTYPE, Adult male. Reg. no. 1925.11.1.771. Mount Driven, 600', south-central Queensland, 4 May 1923. Collected by G. H. Wilkins on the British Museum Expedition to Australia and the Islands.
wilkinsi—Nesospiza wilkinsi P. R. Lowe, 1923
williamsi—Geokichla piaggiae williamsi Macdonald, 1948 31 Bull. Br. Orn. Cl., 69:16. HOLOTYPE, Adult female. Reg. no.

- 1948.1.6. Mount Muhavura, 9.500', south-west Kigezi, Uganda, 24 Sept. 1946. Collected by J. G. Williams and presented by the Coryndon Museum, Nairobi.
- williamsi—Phylloscopus umbrovirens williamsi Clancey, 1956. . . 35

 Bull. Brit. Orn. Cl., 76: 10. HOLOTYPE, Adult female. Reg. no. 1956.23.3. 10 miles north of Erigavo, 7,000', British Somaliland, 27 Feb. 1954. Collected by J. G. Williams. Presented by the Coryndon Museum, Nairobi.
- williamsoni—Mirafra cantillans williamsoni E. C. S. Baker, 1915 . 17 Bull. Br. Orn. Cl., 36: 9. Syntype, Adult male. Reg. no. 1915.11. 11.1. Bangkok, Siam, 31 March 1915. Collected and presented Sir Walter J. F. Williamson.

This is the author's selected male. His selected female is also in the collection.

- williamsoni—Ploceus manyar williamsoni Hall, 1957 61

 Bull. Br. Orn. Cl., 77: 46. HOLOTYPE, Adult male. Reg. no. 1955.1.4192. Samkok, central Siam, 11 June 1921. Collected and presented by Sir Walter J. F. Williamson.
- williamsoni—Turdinus williamsoni Godwin-Austen, 1877 . . . 32B J. Asiat. Soc. Bengal, 46, pt. 2:44. HOLOTYPE, Adult male. Reg. no. 1895.7.14.1264. (Tenga Pani, on label), near Saddya, Assam, 18 Jan. 1877. Collected by M. J. Ogle. Godwin-Austen Collection.
- willoughbyi—Camaroptera superciliaris willoughbyi Bannerman, 1923 35 Bull. Brit. Orn. Cl., 43: 138. (and Ibis (11), 5, (1923): 700). HOLOTYPE, Adult male. Reg. no. 1923.11.12.211. Béoumi, Ivory Coast, West Africa, 30 Nov. 1922. Collected by W. P. Lowe and H. R. Hardy on the British Museum Expedition to the French Ivory Coast.
- wilsoni—Heterorhynchus wilsoni Rothschild, 1893. 55.

 Avifauna of Laysan . . . : 97, pl. 50. SYNTYPE, Adult male.

 Reg. no. 1939.12.9.138. (Collector's number 1351). Hawaii

 (29 Sept. 1891, on label). Collected by H. C. Palmer. Rothschild

 Bequest.

No holotype was designated. The specimen listed as the type by Hartert, *Novit. Zool.*, 26 (1919): 170, was one of the series listed by Rothschild (op. cit., : 98). The above specimen and a female syntype which is also in the collection are part of this series.

type which is also in the collection are part of this series.

wilsoni—Himatione wilsoni Rothschild, 1893
wingatei—Siva wingatei Ogilvie-Grant, 1900
winifredae—Artisornis winifredae Moreau, 1938 39 Bull. Brit. Orn. Cl., 58: 139. Holotype, Adult male. Reg. no. 1938.6.18.1. Kinole, Forest, 6,000', north Uulguru Forest, Tanganyika, 11 July 1937. Collected by and purchased of R. E. Moreau.
winifredae—Zosterops winifredae W. L. Sclater and Moreau, 1934 . 4 Bull. Brit. Orn. Cl., 55:14. HOLOTYPE, Adult male. Reg. no. 1934.11.21.137. Chome, 6,200', South Pare Mountains, Usambara District, Tanganyika, 9 July 1934. Collected by and purchased of R. E. Moreau.
winneba—Cisticola eximia winneba Lynes, 1931
wintoni—Macronyx wintoni Sharpe, 1891
witputzi—Tephrocorys cinerea witputzi Macdonald, 1952 1' Ann. Transvaal Mus., 22:31. HOLOTYPE, Adult male. Reg. no. 1950.50.1095. Witputs, 4,000', Great Namaqualand, 27°35'S., 16°42'E., South-West Africa, 25 Jan. 1950. Collected and presented by the British Museum (Natural History) South-West Africa Expedition.
wollastoni—Anthus wollastoni Ogilvie-Grant, 1913 19 Bull. Br. Orn. Cl., 31:105. HOLOTYPE, Adult male. Reg. no. 1916.5.30.844. Camp 11, 8,000′, Utakwa River, Dutch New Guinea, 6 Feb. 1913. Collected by C. B. Kloss and presented by the committee of the British Ornithologists' Union Expedition to Dutch New Guinea.
woodfordi—Macrocorax woodfordi Oligvie-Grant, 1887

woodi—Mixornis woodi Sharpe, 1877
woodi—Trichalopterum erythrolaema woodi E. C. S. Baker, 1914 . 32B Bull. Br. Orn. Cl., 35:17. HOLOTYPE, Adult. Reg. no. 1915.4.28. 4. Loi Song, Northern Shan States, 7 Jan. Collected by H. Wood and presented by the Bombay Natural History Society.
woodlarkensis — Piezorhynchus alecto woodlarkensis Mathews, 1928. 38 Bull. Brit. Orn. Cl., 48:93. HOLOTYPE, Adult female. Reg. no. 1898.4.30.30. Woodlark Island, Louisade Archipelago, 16 April 1897. Collected by A. S. Meek. Purchased of E. Gerrard.
woodwardi—Bradyornis woodwardi Sharpe, 1877
woolundra—Nesoptilotis leucotis woolundra Mathews, 1922
woosnami—Alethe woosnami Ogilvie-Grant, 1906 31 Bull. Br. Orn. Cl., 19: 24. HOLOTYPE, Adult male. Reg. no. 1906.12.23.1237. Forest near Urumu, 3,000′, North-east Belgian Congo, North-west of Ruwenzori, 18 Aug. 1906. Collected by R. B. Woosnam and presented by the subscribers to the Ruwenzori Expedition.
woosnami—Bleda woosnami Ogilvie-Grant, 1907
woosnami—Cisticola woosnami Ogilvie-Grant, 1908 35 Bull. Brit. Orn. Cl., 21:72. SYNTYPE, Adult male. Reg. no. 1906.12.23.927. Mokia, 3,400', South-east Ruwenzori, 9 May 1906. Collected by R. B. Woosnam and presented by the subscribers to the British Museum Ruwenzori Expedition.
wrayi—Aethopyga wrayi Sharpe, 1887

18.24. Gunong Batu Putch, Perak, Malaya, Collected and presented by L. Wray. A syntype collected by H. C. Robinson is also in the collection. wravi—Pericrocotus wravi Sharpe, 1888 Reg. no. 1888.3.18.57. Batang Padang Mountains, Perak, Malaya. Collected and presented by L. Wray. The female syntype is also in the collection. wravi—Pomatorhinus wravi Sharpe, 1887 Proc. zool. Soc. Lond., 1887: 437. SYNTYPE, Adult female. Reg. no. 1886.12.22.14. Larut Range, c. 4,400', Perak, Malaya, Sept.— Oct. 1886. Collected and presented by L. Wray. The male syntype is also in the collection. wravi—Sibia wravi Ogilvie-Grant, 1910 Bull. Br. Orn. Cl., 25:98. SYNTYPE, Adult male. Reg. no. 1888.3.18.29. (Batang Batang, on label), Perak Mountains, Malava. Collected and presented by L. Wray. This is the author's selected male. His selected female is also in the collection. wuchereri—Arremon wuchereri P. L. Sclater and Salvin, 1873. Nomencl. Av. Neotrop.: 25, 157. HOLOTYPE, Adult male. Reg. no. 1885.6.8.432. Near Bahia, Brazil. Collected by Wucherer. Salvin-Godman Collection. wuchereri—Euscarthmus wuchereri P. L. Sclater and Salvin, 1873 Nomencl. Av. Neotrop.: 158. HOLOTYPE, Adult. Reg. no. 1888. 1.13.274. Bahai, north-eastern Brazil, 1862. Collected by Wucherer. Sclater Collection. wyatti—Synallaxis wyatti P. L. Sclater and Salvin, 1870 Proc. zool. Soc. Lond., 1870: 840. HOLOTYPE, Adult male. Reg. no. 1889.5.20,267. Paramo of Pamplona, Colombia, 1870. Collected by C. Wyatt. Sclater Collection. xanthochlamys—Pyromelana xanthochlamys Sharpe, 1902 Bull. Br. Orn. Cl., 13:10. HOLOTYPE, Adult male. Reg. no. 1902.12.8.7. Hoima, western Uganda, 26 Oct. 1901. Collected and presented by F. J. Jackson. xanthochlora—Anthreptes xanthochlora Hume, 1875 . . 46 Stray Feathers, 3:320 (note). HOLOTYPE, Adult female. Reg. no. 1886.12.1.1254. Papyai, Mergui district, Tenasserim, Burma, (14 Nov. 1874, on label). Collected by W. Davison. Hume

Collection.

- xanthochloris—Pteruthius xanthochloris J. E. Gray, 1846

 Catalogue of . . . mammalia and birds of Nepal and Thibet presented by B. H. Hodgson to the British Museum: 95, 155. SYNTYPE, Adult (relaxed mount). Reg. no. 1843.1.13.411. Nepal. Collected and presented by B. H. Hodgson.

 Listed as type by Gadow, Cat. Bds. Brit. Mus., 8 (1883): 118.
- xanthochroa—Zosterops xanthochroa G. R. Gray, 1859. . . . 47

 Proc. zool. Soc. Lond., 1859: 161. HOLOTYPE, Adult (male, vide Gadow, Cat. Bds. Brit. Mus., 9 (1884): 174). Reg. no. 1859.4.19.7. Island of Nu, Port of France, New Caledonia. Collected by J. Macgillivray. Purchased of Cuming.
- xanthocnemis—Pachycephala xanthocnemis G. R. Gray, 1860 . 39

 Proc. zool. Soc. Lond., 1860: 353. Syntype, Adult (female, vide Sharpe, Cat. Bds. Brit. Mus., 8 (1883): 186, footnote). Reg. no. 1873.5.12.509. Amboyna, Moluccas. 1857. Collected by A. R. Wallace. Wallace Collection.
- xanthodryas—Phyllopneuste xanthodryas Swinhoe, 1863 . . . 35

 Proc. zool. Soc. Lond., 1863: 296. SYNTYPE, Adult. Reg. no. 1898.9.1.263. Amoy, China, May 1861. Collected by R. Swinhoe. Seebohm Bequest.

This specimen and another from the Seebohm Collections were labelled "Type". A specimen from the Hume Collection is also a syntype.

- xanthogaster—Atrornis xanthogaster Hodgson, 1844 35 in J. E. Gray, Zoological Miscellany: 82; and also J. E. Gray, List of mammalia and birds of Nepal and Thibet presented by B. H. Hodgson . . . to the British Museum, (1846): 66, 152. These are nomina nuda, and the specimen listed by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 66, has no type status.
- xanthogaster—Trichophorus xanthogaster Hodgson, 1844 . . . 2: in J. E. Gray, Zoological Miscellany, 1844 : 83. A specimen in the collection was listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881): 77 but the name is a nomen nudum.
- xanthogastra—Calliste xanthogastra P. L. Sclater, 1851. . . . 51 in Jardine, Contributions to Ornithology: 23, 55. HOLOTYPE, Adult male. Reg. no. 1885.6.12.254. Rio Negro, Upper Amazon, Brazil. Collected for Verreaux. Sclater Collection, received by him from E. Wilson.

- xanthogenys—Machaerirhynchus xanthogenys G. R. Gray, 1858 . 38 Proc. zool. Soc. Lond., 1858: 176. HOLOTYPE, Adult male. Reg. no. 1858.3.10.58. Aru Islands. Collected by A. R. Wallace. Wallace Collection.
- xanthogramma—Chlorospiza xanthogramma G. R. Gray, 1839 . 49 Zoology of the voyage of H.M.S. Beagle, 3, Birds: 96, pl. 33. Syntype, Immature. Reg. no. 1841.1.18.24. Tierra del Fuego, February. Collected and presented by C. Darwin.

This specimen appears to be part of the type material but was apparently overlooked by Sharpe and not listed in *Cat. Bds. Brit. Mus.* The specimen from Hermit Island listed as the type by Sharpe, op. cit., 12 (1888): 790, has no type status.

- xantholaemus—Brachypus xantholaemus Jerdon, 1844. . . . 21
 Madras J. Lit. Sci., 31, vol. 13:122. Syntype, Adult. Reg. no. 1846.4.30.33. West of Nellore, Eastern Ghats, 1842. Collected for T. C. Jerdon. Hume Collection.
 Listed as type by Sharpe, Cat. Bds. Brit. Mus., 6 (1881):146.
- xanthomelana—Cissopica whiteheadi xanthomelana Delacour, 1927 71 Bull. Brit. Orn. Cl., 47:164. Holotype, Adult male. Reg. no. 1927.6.5.19. Backan, north-east Tonkin, French Indo-China, 25 Dec. 1926. Collected by M. P. Jabouille, J. Delacour, and W. P. Lowe. From the Third Expedition to French Indo-China.
- xanthomus—Icterus xanthomus P. L. Sclater, 1862 57

 Catalogue of a collection of American Birds: 131. HOLOTYPE,

 Adult. Reg. no. 1885.7.12.133. Mexico (=Porto Rica, vide Sclater, Ibis, (1) 6 (1864): 168). Collected for Verreaux. Sclater Collection.
- xanthophrys—Chlorospingus xanthophrys P. L. Sclater, 1856. . . 54
 Proc. zool. Soc. Lond., 1856: 30, 93. Holotype, Adult. Reg. no.
 1884.5.15.470. Bogota, Colombia, 1854. (Collected by Parzudaki, on label.) Sclater Collection.
 See also Sclater, Proc. zool. Soc. Lond., 1875: 235.

- xanthophrys—Motacilla xanthophrys Sharpe, 1885 19 Cat. Bds. Brit. Mus., 10: 532, pl. 8. HOLOTYPE, Adult male. Reg. no. 1898.10.20.716. Lenkoran, Persia, March. Seebohm Bequest.
- xanthophrys—Oxylabes xanthophrys Sharpe, 1875 21

 Proc. zool. Soc. Lond., 1875: 76. HOLOTYPE, Adult. Reg. no. 1875.2.1.21. Central Madagascar. Collected by A. Crossley and purchased of W. Cutter.

No holotype was designated. The specimen listed as the type by Hartert, *Novit. Zool.*, 26 (1919): 170, was one of a series of syntypes

which also included the above specimen.

- xanthopterus—Hyphantornis xanthopterus Finsch and Hartlaub, 1870 61

 Die Vögel Ost-Afrikas: 399. (in Decken, Reisen in Ost-Afrikas,
 4). HOLOTYPE, Adult male. Reg. no. 1863.12.30.7. Shupanga,
 Lower Shiré River, Zambesi River. (Dec. 1861, on label). Collected by Sir J. Kirk. Presented by Earl Russell of the Foreign
 Office, from the Livingstone Zambesi Expedition.
- xanthopygium—Dicaeum xanthopygium Tweeddale, 1877 . . . 45
 Ann. Mag. nat. Hist., (4) 20:95. HOLOTYPE, Adult male. Reg.
 no. 1888.7.12.74. Monte Alban, Luzon Island, Philippines, (Feb.
 1877, on label). Tweeddale Collection.
- xanthopygius—Serimus xanthopygius Rüppell, 1840 59

 Neue Wirbelthiere zu der Fauna von Abyssinien gehorig, : 96, pl. 35,

 Adult. Reg. no. 1842.8.16.8. Simen, (Abyssinia, on label). Collected by and purchased of E. Rüppell.
- xanthopygius—Tachyphonus xanthopygius P. L. Sclater, 1855. . 5:

 Proc. zool. Soc. Lond., 1854 (publ. 1855): 158, pl. 69. SYNTYPE,

 Adult (female, vide Sclater, Cat. Bds. Brit. Mus., 11 (1886): 209).

 Bogota, New Grenada. Purchased of S. Stevens.

 Listed as the type by Sclater, Cat. Bds. Brit. Mus., 11 (1886): 209,

 but more than one specimen is referred to in the original description.

In the text (on, cit.), G. R. Grav uses the name A. manimbe, quoting A. xanthornus as a synonym. xanthoscelus—Turdus xanthoscelus Jardine, 1847. 31 Ann. Mag. nat. Hist., (1) 20:329. SYNTYPE, Adult male. Reg. no. 1896.6.1.485. Tobago, 1847. Collected by J. Kirk. Seebohm Bequest. The female syntype is also in the collection. xanthoschistos—Phyllopneuste xanthoschistos J. E. Grav. 1846 Catalogue of mammalia and birds of Nepal and Thibet presented by B. H. Hodgson . . . to the British Museum: 65, 151. SYNTYPE. Adult. Reg. no. 1843.1.13.389. Nepal. Collected and presented by B. H. Hodgson. Gray used a Hodgson MS. name. Listed as the type by Sharpe, Cat. Bds. Brit. Mus., 4 (1879): 398. There are several other syntypes in the collection. xanthostomus—Parus xanthostomus Shelley, 1892. Bull. Brit. Orn. Cl., 1:6. HOLOTYPE, Adult. Reg. no. 1895. 5.1.1402. Zambesia (=between Limpopo and Zambesi Rivers.) (Collected by Bradshaw vide Shelley, Birds of Africa, 2 (1900): 236.) Shelley Collection. xanthotis—Aethorhynchus xanthotis Sharpe, 1881. 22 Cat. Bds. Brit. Mus., 6:15. HOLOTYPE, Adult ?female. Reg. no. 1860.11.9.14. Cambodia. Collected by M. Mouhot and purchased of S. Stevens. yamathini—Anthus similis yamathini Hall, 1957 19 Bull. Br. Orn. Cl., 77:102. HOLOTYPE, Adult male. Reg. no. 1948.80.2754. Aingdo, Yamathin district, central Burma, 15 Nov. 1938. Collected by H. C. Smith and presented by Smith and P. F. Garthwaite. yangpiensis—Yuhina yangpiensis Sharpe, 1902 Bull. Br. Orn. Cl., 13:12. HOLOTYPE, Adult. Reg. no. 1903.8.8. 132. Yangpi, 5,300', Tali-fu road, west Yunnan, 22 March 1902. Collected and presented by G. Rippon. vatii—Passer vatii Sharpe 1888 Cat. Bds. Brit. Mus., 13: 322. HOLOTYPE, Adult male. Reg. no. 1886.9.16.178. De-dadi, Western Afghanistan. Collected and presented by J. Aitchison. yavelloensis—Aethocorys personata yavelloensis Benson, 1947. Bull. Br. Orn. Cl., 68:9. HOLOTYPE, Adult male. Reg. no. 1946.5.2056. 10 miles west of Yavello, south Abyssinia, 23 July 1941. Collected and presented by C. W. Benson. yemenensis—Cryptolopha umbrovicens yemenensis Ogilvie-Grant, 1913 35 Bull. Brit. Orn. Cl., 31:90. SYNTYPE, Adult male. Reg. no.

1913.7.18.9. Menchaa, 8,000', Yemen, South Arabia, 20 Dec. 1912. Collected by and purchased of G. W. Bury. This is the described male syntype. The female syntype is also in the collection.
yemenensis—Lanius yemenensis Ogilvie-Grant, 1914 23 Bull. Br. Orn. Cl., 33:91. Holotype, Adult female. Reg. no. 1916.1.28.66. Sanaa, 7,600', Yemen, South Arabia, 3 Sept. 1913. Collected by and purchased of G. W. Bury.
yemenensis—Oenanthe yemenensis Ogilvie-Grant, 1913 31 Bull. Br. Orn. Cl., 31:87. SYNTYPE, Adult male. Reg. no. 1913. 7.18.3. Menacha, 7,000', Yemen, Arabia, 25 Jan. 1913. Collected by and purchased of G. W. Bury. The female syntype is also in the collection.
yemenensis—Pseudacanthis yemenensis Ogilvie-Grant, 1931 59 Bull. Brit. Orn. Cl., 31: 89. Syntype, Adult male. Reg. no. 1913. 7.18.7. Menacha, 8,500', Yemen, Southern Arabia, 16 Jan. 1913. Collected by and purchased of G. W. Bury. The female syntype is also in the collection.
yemenensis—Rhynchostruthus percivali yemenensis Ogilvie-Grant 1913
yerburyi—Myrmecocichla yerburyi Sharpe, 1895
yersini—Trochalopteron yersini Robinson and Kloss, 1919 32B Ibis, (11) 1:575, pl. 13. SYNTYPE, Adult male. Reg. no. 1919. 12.20.269. Langbian Peaks, 5,500-7,500', south Annam, 22 April 1918. Collected and presented by C. B. Kloss. This is the authors' selected male. Their selected female syntype is also in the collection.
yessoensis—Schoenicola yessoensis Swinhoe, 1874
younghusbandi—Cinclus younghusbandi Walton, 1905 27 Bull. Br. Orn. Cl., 15:92. HOLOTYPE, Adult female. Reg. no. 1905.12.21.212. Khamba Jong, Tibet, 6 Nov. 1903. Collected and presented by H. J. Walton. (cont.)

- The collector's number of this specimen is 1770, not 153 as printed. See Kinnear, *Ibis*, (14) 1 (1937): 262.

- yunnanense—Trochalopterum yunnanense Rippon, 1906. . . . 32B Bull. Brit. Orn. Cl., 19:32. HOLOTYPE, Adult. Reg. no. 1906. 12.17.155. Yangtze River (big bend), 10,000', Western Yunnan, 22 March 1906. Collected and presented by G. Rippon.
- yunnanensis—Alcippe fratercula yunnanensis Harington, 1913. . 32B Bull. Br. Orn. Cl., 33:63. Holotype, Adult. Reg. no. 1903. 8.8.162. Gyi-dzin-shan, 8,900', east of Talifu, Yunnan Mountains, 2 April 1902. Collected and presented by Col. G. Rippon.

- yunnanensis—Leiothrix luteus yunnanensis Rothschild, 1921 . . . 32B Novit. Zool., 28:36. Syntype, Adult male. Reg. no. 1921.7.15. 245. Shweli-Salwin Divide, 7,000-10,000', Yunnan, May 1919. Collected by G. Forrest and presented by Stephenson Clarke. There are three other syntypes in the collection.

- yunnanensis—Parus monticolus yunnanensis La Touche, 1921. 4.

 Bull. Brit. Orn. Cl., 42:51. SYNTYPE, Adult male. Reg. no.
 1923.3.10.83. Milati, south-east Yunnan, 20 Jan. 1921. Collected and presented by J. D. La Touche.

 There are a male and a female syntype at Harvard, vide Bangs,

Bull. Mus. Comp. Zool., Harvard, 70 (1930): 350.

yunnanensis—Pteruthius erythropterus yunnanensis Ticehurst, 1937 32B

- yunnanensis—Pteruthius erythropterus yunnanensis Ticehurst, 1937 32B Bull. Br. Orn. Cl., 57: 147. HOLOTYPE, Adult male. Reg. no. 1921.7.15.331. Shweli-Salween Divide, 8,000', north-west Yunnan, Dec. 1919. Collected by G. Forrest and presented by Col. Stephenson Clarke.
- yunnanensis—Sitta yunnanensis Ogilvie-Grant, 1900 43

 Bull. Brit. Orn. Cl., 10:37. HOLOTYPE, Adult (male, on label).

 Reg. no. 1900.3.5.53. Wei-Yuan, East of Mekong River, Southern Yunnan (12 March 1899, on label). Collected and presented by A. M. S. Wingate.

There are seven other syntypes in the collection.

- yunnanensis—Tarsiger indicus yunnanensis Rothschild, 1922. . . 31

 Bull. Br. Orn. Cl., 43:10. Holotype, Adult male. Reg. no. 1922.12.7.98. Lichiang, 10,000', north-west Yunnan, 13 Nov. 1921. Collected by G. Forrest and presented by Col. Stephenson Clarke.

Ibis, (12) 6, Supplement: 390. HOLOTYPE, Adult male. Reg. no. 1922.12.8.1067. Zalingei, Darfur Province, Sudan, 7 Oct. 1920. Collected by Admiral H. Lynes and W. P. Lowe. Presented by Admiral H. Lynes.
zalingei—Galerida cristata zalingei C. H. B. Grant and Mackworth-Praed, 1939
zambesiae—Chlorocichla zambesiae Shelley, 1896 21 The Birds of Africa, 1:64. HOLOTYPE, Adult. Reg. no. 1895. 5.1.733. Makalaka, south Zambesia. Collected by Bradshaw. Shelley Collection.
zambesiana—Anthodiaeta zambesiana Shelley, 1876 46 Monograph of the Nectariniidae, : 343, pl. 111. HOLOTYPE, Adult male. Reg. no. 1863.12.30.21. Shupanga, Zambezi. Collected by Dr. Kirk. Presented by Earl Russell, Secretary of State for the Colonies; from Livingstone's Expedition, 1858–1864.
zambesiana—Erythropygia zambeziana Sharpe, 1882 31 Proc. zool. Soc. Lond., 1882: 588, pl. 45. HOLOTYPE, Adult. Reg. no. 1860.12.31.103. Tete, Lower Zambezi. Collected by Sir J. Kirk on the Livingstone Expedition and presented by Lord Russell of the Foreign Office.
zantholeuca—Erpornis zantholeuca Blyth, 1844
zanzibarica—Chalcomitra veroxii zanzibarica Grote, 1932
zanzibarica—Urobrachya zanzibarica Shelley, 1881
zanzibaricus—Eurillas virens zanzibaricus Pakenham, 1935 21 Bull. Br. Orn. Cl., 55: 111. HOLOTYPE, Adult male. Reg. no.

1935.2.6.1. Jozani Forest, Zanzibar, 7 Nov. 1934. Collected and presented by R. H. W. Pakenham.

2 Lapinion
zaphiroi—Sycobrotus zaphiroi Ogilvie-Grant, 1902 6 Bull. Br. Orn. Cl., 13:22. HOLOTYPE, Adult female. Reg. no. 1902.11.8.25. Abou Beker, Harar, South Abyssinia, 10 June 1902. Collected by and purchased of P. Zaphiro.
zetlandicus—Sturnus vulgaris zetlandicus Hartert, 1918 6 Novit. Zool., 25: 329. HOLOTYPE, Adult male. Reg. no. 1913. 11.7.3. Gutcher, north Yell, Shetland Islands, 22 Nov. 1913. Collected and presented by R. H. Nicholson.
ziaratensis—Ianthocincla lineatum ziaratensis Ticehurst, 1920 . 32 Bull. Br. Orn. Cl., 41:55. HOLOTYPE, Adult male. Reg. no. 1920.12.4.1. Juniper forests of Ziarat, 8,000', near Quetta, north Baluchistan, 25 Sept. 1919. Collected and presented by C. B. Ticehurst.
ziaratensis—Parus major ziaratenis Whistler, 1929 4 Bull. Brit. Orn. Cl., 50: 7. HOLOTYPE, Adult male. Reg. no. 1941.5.30.622. Ziarat, 8,500', Baluchistan. 30 Sept. 1919. Collected by C. B. Ticehurst. Ticehurst Bequest.
zita—Dicaeum sulaense zita Harrisson and Hartley, 1934 4 Bull. Brit. Orn. Cl., 54:159. HOLOTYPE, Adult male. Reg. no. 1933.11.2.8. Mt. Dulit, 4,500', central Borneo, 15 Sept. 1932. Collected by T. H. Harrisson and C. H. Hartley. From the Oxford University Expedition to Borneo.
zombae—Mirafra zombae Ogilvie-Grant, 1902
zombensis—Andropadus zombensis Shelley, 1894
Zonarius—Cinnyris chalybeus zonarius Vincent, 1933 4 Bull. Brit. Orn. Cl., 53: 146. HOLOTYPE, Adult male. Reg. no. 1933.3.1.7. Near Zobué, 2,900', Portuguese East Africa – Nyasaland border, 15°36' S., 34°27' E., 10 April 1910. Collected by and purchased of J. Vincent.
Proc. zool. Soc. Lond., 1875: 76. Syntype, Adult (supposed female, vide Sharpe, Cat. Bds. Brit. Mus., 7 (1883): 571). Reg. no. 1875. 2.1.22. Central Madagascar. Collected by A. Crossley and purchased of W. Cutter. Listed as type by Sharpe, 1883, loc. cit. The male syntype is also in the collection.



CORRECTIONS AND ADDITIONS

TO

VOLUME 1

In the following list of amendments and additions to volume one the entries are prefixed by the page number in the first volume on which they appear or would have appeared. Since the decision to include the type water-colour drawings in the library of the Zoology Department of the British Museum (Natural History) was not taken until the first volume had already been prepared, the list is larger than it might otherwise have been.

p. 11.	alfredi—Crypselus alfredi
p. 11.	alleni—Selaphorus alleni Henshaw, 1877

The above collection incorporated the Henshaw Collection, vide Grinnell, Univ. Calif. Publ. Zool., 38 (1932): 280. The author's selected syntype cannot be traced but this specimen is referred to in the description and has "type" on the label, a female specimen in the Museum is also marked "type".

- p. 13. amethystinus—Lampornis amethystinus. 25
 Phillips, Bull. Br. Orn. Cl., 86 (1966): 103-4. when discussing the subspecies of L. amethystinus, gives the locality of O. henricus as "Coatepac, near Xalapa".

Mathews, Systema Avium Australasianarum, 1 (1927): 259, note, considers that the drawing is indeterminable.

Vol. 2, no. 8 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 5 (1916): 323.

p. 39. boothi—Petrophassa albipennis boothi Goodwin, 1969 . 19
Bull. Br. Orn. Cl., 89: 132. The holotype was registered in the Brit. Mus. register as no. 1969. 4. 112. It has, however, been presented to the Arid Zone Research Institute, Alice Springs, Northern Territory, Australia.

- p. 47. caerulescens—Rallus caerulescens Gmelin 1779. 15

 Systema Naturae, 1, pt. 2:716. HOLOTYPE, signed and completed plate. Seen in a menagerie at Cape of Good Hope. No. 129 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 302.
- p. 53. **catherina**—Eriocnemis catherina. For catherina read catharina.
- p. 54. cautacuta—Hirundo cautacuta Latham, 1801 24

 Index Ornithologicus, Suppl.: lvii. Holotype, Signed watercolour drawing. New South Wales. No. 217 in Watling,
 Original water-colour drawings . . . Library of the Zoology
 Department, British Museum (Natural History). Vide Sharpe,
 Hist. Coll. Brit. Mus., 2 (1906): 144, who reversed this and no.
 216.
 - Vol. 2, no. 88 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue, vol. 2). See also, Mathews, Systema Avium Australasianarum, 1 (1927): 404.
- p. 65. **cobanese**—Glaucidium cobanese. For cobanese read cobanense.

Gibson-Hill, Bull. Raffles Mus., 19 (1949): 156, comments on the sex of the specimen and confirms that the holotype is in the collection.

Vol. 2, no. 95 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also

Mathews, Birds of Australia, 5 (1916): 338.

- p. 70. cozumelae—Coccyzus minor cozumelae Van Rossem, 1934. 12

 Bull. Mus. comp. Zool., Harv., 77: 390. Holotype, Adult.

 Reg. no. 1897.4.1.49. Cozumel Island, off coast of Quintana Roo, Mexico, Jan. 1886. Collected by G. F. Gaumer. Salvin-Godman Collection.
- p. 74. cyanocephalus—Cuculus cyanocephalus Latham, 1801 . 21

 Index Ornithologicus, Suppl.: xxx. ? HOLOTYPE, Signed water-colour drawing. New South Wales. No. 72 in Watling, Original water-colour drawings Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 121.

Vol. 1, no. 34 in Lambert drawings, in Library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). *See also* Mathews, *Systema Avium Australasianarum*, 1 (1927): 418.

- p. 86. ecaudata—Rallus ecaudata J. F. Miller, 1783 15

 Icones Animalium: pl. 47. HOLOTYPE, Unsigned painting.

 Tonga, 1 July 1774. No. 127 in Forster, unpublished drawings
 . . . made during Cook's second voyage. Library of Zoology
 Department, British Museum (Natural History). Vide Lysaght,
 Bull. Brit. Mus. (Nat. Hist), Hist. Ser., 1 (1959): 302.
- p. 95. fasciata—Porzana fasciata P. L. Sclater and Salvin, 1868 . 15

 Proc. zool. Soc. Lond., 1867 (Publ. early 1868): 981. SYNTYPE, Adult male. Reg. no. 1889.11.20.138. Pebas, Upper
 Amazon, Peru, 27 June 1866. Collected by J. Hauxwell.
 Salvin-Godman Collection.
 This was renamed Porzana hauxwelli, P. L. Sclater and

Salvin, 1869 (see entry in Warren, Type specimens of birds, 1 (1966): 126.—The authors considering it preoccupied by Rallina fasciata Raffles, 1822). See Hellmayr, Catalogue of birds of the Americas, 1 (1942): 382.

- p. 97. flabelliformis—Cuculus flabelliformis Latham, 1801 . . . 21

 Index Ornithologicus, Suppl.: xxx. Sharpe, Hist. Coll. Brit.

 Mus., 2 (1906): 121, lists a drawing, no. 75 in Watling, Original water-colour drawings . . ., as the type, but this is not regarded as referable to the type description.
- p. 110. glanvillei—Apus barbatus glanvillei Benson, 1967 . . . 24

 Bull. Brit. Orn. Cl., 87: 125. HOLOTYPE, Adult female.

 Reg. no. 1938.6.5.14 (not 15 as printed). Rokupr, Sierra
 Leone, 9°01′ N., 12°57′ W., 28 Sept. 1937. Collected and
 presented by R. R. Glanville.

22

p. 117.	Order number should be 22, not 25.
p. 124.	harmani—Crossoptilon harmani. Altitude should be "not less than 10,000 feet," vide Ludlow, Ibis, 86 (1944): 378.
p. 126.	hawaiiensis—Anous hawaiiensis Rothschild, 1893 17 Bull. Brit. Orn. Cl., 1: 57. SYNTYPE, Adult (male, on label). Reg. no. 1968.88.1 (subsequently assigned). Sandwich Group, (Kauai Island, 21 April 1891, on label). Collected by H. C. Palmer. Presented by the Hon. W. Rothschild. The author stated that types were at both Tring and the British Museum, and the specimen listed as the type by Hartert, Novit. Zool., 34 (1927): 18 must be a syntype.
p. 150.	kathleenae—Guttera edouardi kathleenae C. M. N. White, 1943
p. 156.	layardi—Ptilopus layardi D. G. Elliot, 1878 19 Proc. zool. Soc. Lond., 1878 : 567. New name for Chrysoena viridis Layard, 1875 (q.v.) on the grounds of preoccupation, by Columba viridis Linnaeus, 1766.
p. 167.	lucidus—Cuculus lucidus Gmelin, 1788

morroronsis Otus trichonsis quarraransis

p. 168. lunulatus—Falco lunulatus Latham, 1801 13

Index Ornithologicus, Suppl.: xiii. ? Holotype, Watercolour drawing. New South Wales. No. 12 in Watling,
Original water-colour drawings Library of the Zoology
Department, British Museum (Natural History). Vide Sharpe,
Hist. Coll. Brit. Mus., 2 (1906): 111, who refers to it in error
as lunatus.

Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1, (1959): 286.

Vol. 2, no. 3 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 5 (1916): 228.

during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght,

- p. 177. marina—Procellaria marina Latham, 1790 9

 Index Ornithologicus, 2:826. Holotype, unfinished and unsigned painting. Lat. 37° S., 23 Dec. 1768. No. 12 in Parkinson, Unpublished water-colour drawings of animals made during Cook's first voyage. Library of the Zoology Department, British Museum (Natural History), vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 274.
- p. 182. meinertzhageni—Tringa totanus meinertzhageni Thiede, MS? 16

 A specimen was selected and labelled, c. 1964, as the holotype of this then unpublished name by W. Thiede, who has since used it in print (e.g. Zool. Anz., 177 (1966): 171). We are unable to discover whether this name has been validly published; the entry here does not make it nomenclatorially available. The selected specimen is an adult male, reg. no. 1965. M. 127, from Shushal, E. Ladak, 10 June 1925, collected and presented by R. Meinertzhagen.
- p. 195. mollis—Procellaria mollis Gould, 1844 9

 Ann. Mag. nat. Hist., 13:363. Syntype, Adult male, (relaxed mount). Reg. no. 1841.6.1307. South Atlantic, 29°45′ S., 15°3′W., 27 July 1838. Collected by and purchased of J. Gould.

Listed with a female, now untraced, as types by Salvin, Cat. Bds. Brit. Mus., 25 (1896): 406. Stone and Mathews, Austral Avian Record, 1 (1913): 137, state that the type is in Philadelphia; but it is not listed by de Schauensee, Proc. Acad. nat. Sci. Philad., 109 (1957).

- p. 202. newtoni—Tinnunculus newtoni Gurney, 1863 13

 Ibis, (1) 5:34, pl. 2. Syntype, Adult male. Reg. no. 1955.6.N.20.1365. Madagascar, (Buravin, Madagascar, 1 Oct. 1861, on label). Collected by E. Newton and presented by him to J. H. Gurney. Purchased of Norwich Castle Museum. The second syntype, a male in the dark phase, is also in the

collection.

- p. 209. novaehollandiae—Falco novaehollandiae Gmelin, 1788 . 13

 Systema Naturae, 1, pt. 1:264. HOLOTYPE, Unsigned pencil sketch. Tasmania, late summer of 1773. No. 35 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 281.
- p. 209. novaeseelandiae—Anas novaeseelandiae Gmelin, 1789 . 12

 Systema Naturae, 1, pt. 2:541. Holotype, Unsigned painting. Dusky Sound, South Island, New Zealand, 1773. No. 79 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 290.

Vol. 2, no. 85 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue, vol. 2).

p. 218. pacifica—Coracius pacifica Latham, 1801 30

Index Ornithologicus, Suppl.: xxvii. ? Holotype, Watercolour drawing. New South Wales. No. 64 in Watling,
Original water-colour drawings . . . Library of the Zoology
Department, British Museum (Natural History). Vide Sharpe,
Hist. Coll. Brit. Mus., 2 (1906): 120.

Vol. 2, no. 83 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews, Systema Avium Australasianarum, 1 (1927): 361.

- p. 218. pacifica—Hirundo pacifica Latham, 1801 . . . 24

 Index Ornithologicus, Suppl.: Iviii. ? HOLOTYPE, Signed water-colour drawing. New South Wales. No. 216 in Watling, Original water-colour drawings . . . Library of the Zoology Department, British Museum (Natural History). Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 144, who reversed this and no 217.
 - Vol. 2, no. 89 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2). See also Mathews, Systema Avium Australasianarum, I (1927): 405.

(cont.)

Zoology Department, British Museum (Natural History).

Vide Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 151.

Vol. 2, no. 92 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue, vol. 2). The description is of an immature bird, vide Mathews, Birds of Australia, 2 (1913): 477.

- p. 219. palliolatus—Cuculus palliolatus Latham, 1801. . . . 21

 Index Ornithologicus, Suppl.: xxx. Sharpe, Hist. Coll. Brit.

 Mus., 2 (1906): 121, lists a drawing, no. 74 in Watling,

 Original water-colour drawings . . . , as the type, but this is
 not regarded as referable to the type description.
- p. 226. phasianinus—Cuculus phasianinus Latham, 1801 . . . 21

 Index Ornithologicus, Suppl.: xxx. ? Holotype, Watercolour drawing. New South Wales. No. 73 in Watling,
 Original water-colour drawings . . . Library of the Zoology
 Department, (British Museum (Natural History). Vide Sharpe,
 Hist. Coll. Brit. Mus., 2 (1906): 112.

Vol. 1, no. 36 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews, Systema Avium Australasianarum, 1 (1927): 421.

- p. 229. plagosus—Cuculus plagosus Latham, 1801 . . . 21

 Index Ornithologicus, Suppl.: xxxi. ? Holotype, Watercolour drawing. New South Wales. No. 76 in Watling,
 Original water-colour drawings Library of the Zoological Department, British Museum (Natural History). Vide
 Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 122.

Vol. 3, no. 22 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). The author, in General Synopsis of birds, Suppl., 2; (1802): 139, refers to Lambert.

p. 229. plancus—Falco plancus J. F. Miller, 1777 13

Icones Animalium, pt. 3: pl. 17. HOLOTYPE, Unsigned painting. Tierra del Fuego, 26 Dec. 1774. No. 33 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 281.

Vol. 1, no. 38; vol. 3, no. 24 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue vol. 2). See also Mathews, Birds of Australia, 5 (1915): 89.

- p. 268. serlei—Apus barbatus serlei De Roo, 1970 . . . 24
 Rev. Zool. Bot. Afr., 81:157. Holotype, Adult male.
 Reg. no. 1966.16.120. Sagpa, 5,200', Bamenda Division,
 Southern Cameroons, 6° N., 10° 17' E., 12 Jan. 1957. Collected by and purchased of Dr. W. Serle.
- p. 268. serresiana—Eudyptula serresiana Oustalet, 1878

 Annls. Sci. nat., (6) 8: art. no. 4. An immature specimen from the type locality, Tierra del Fuego, was received in exchange with P. L. Sclater. Reg. no. 1897.5.28.1. It is referred to by Ogilvie-Grant, Cat. Bds. Brit. Mus., 26 (1898): 638, as a "co-type". An early label bears the names "M. Roux, M. Willems" and also "C. G. 1892, no. 797", but there is no evidence to show whether this refers to Admiral Serres's expedition or a later one. The specimen has not been selected.
- p. 284. strigoides—Caprimulgus strigoides Latham, 1801 . 23

 Index Ornithologicus, Suppl.: lviii. ? Holotype, Watercolour drawing. New South Wales. No. 220 in Watling,
 Original water-colour drawings . . . Library of the Zoology
 Department, British Museum (Natural History). Vide Sharpe,
 Hist. Coll. Brit. Mus., 2 (1906): 145.

Vol. 1, no. 67 in Lambert drawings, in library at Knowsley, Lancashire, (see introduction to this catalogue vol. 2). See also Mathews, Systema Avium Australasianarum, I (1927): 355.

- p. 288. superciliosa—Anas superciliosa Gmelin, 1789 12

 Systema Naturae, 1, pt. 2:537. Holotype, Unsigned painting. New Zealand, 1773. No. 77 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History.) Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 290.
- p. 291. tannensis—Columba tannensis Latham, 1790 19

 Index Ornithologicus, 2: 600. Holotype, Finished unsigned painting. Tanna, New Hebrides, 1774. No. 138 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 304.

Vol. 1, no. 60 in Lambert drawings, in library at Knowsley, Lancashire (see introduction to this catalogue, vol. 2). Sharpe, Hist. Coll. Brit. Mus., 2 (1906): 145 lists this drawing, in error, as the type of New Holland Quail, P. australis Latham, 1801,

but Perdix varia is the New Holland Partridge.

- p. 303. variegata—Anas variegata Gmelin, 1789. 12

 Systema Naturae, 1, pt. 2:505. Holotype, Unsigned painting. Dusky Bay, New Zealand, 7 April 1773. No. 67 in Forster, Unpublished water-colour . . . drawings made during Cook's second voyage. Library of the Zoology Department, British Museum (Natural History). Vide Lysaght, Bull. Brit. Mus. (Nat. Hist.), Hist. Ser., 1 (1959): 288.

CORRECTIONS AND ADDITIONS

TO

VOLUME 2

In the following list of amendments and additions to the present volume the entries are prefixed by the page number in the first volume on which they appear or would have appeared. Brief references are present at the appropriate places in the main text.

- p. 13. albiventris—Loxia albiventris Swinhoe, 1870 59

 Renamed Loxia curvirostra japonica Ridgway, 1885, on the grounds of its preoccupation by Loxia albiventris Hermann, 1804 (unidentified).
- p. 22. amicorum—Turdus torquatus amicorum Hartert, 1923 . 31 Vögel der paläarktischen Fauna, Nachtrag: 57. New name for Merula torquata orientalis Seebohm, 1888, (q.v.), on the grounds of its preoccupation by Turdus orientalis Gmelin, 1789.
- p. 27. annamarulae—Melaenornis annamarulae Forbes-Watson, 1970 37

 Bull. Brit. Orn. Cl., 90: 146. HOLOTYPE, Adult male. Reg. no. 1969.25.1. Grassfield, 550 metres, Mount Nimba, Liberia, 7° 30′ N, 8° 35′ W, 12 Aug. 1967. Collected by A. D. Forbes-Watson. Presented by the IUCN Nimba Research Committee.
- p. 44. aureifrons—Zosterops aureifrons Wallace, 1864. . . . 47
 Renamed Zosterops wallacei Finsch, 1901, on the grounds of its preoccupation by Z. aureifrons Heuglin, 1863.
- p. 66. bicolor—Chimarrhornis bicolor Ogilvie-Grant, 1894 . . . 31

 Bull. Br. Orn. Cl., 3: 49. Holotype, Adult male. Reg. no. 1897.5.13.82. North west Luzon, Philippines, 9 Jan. 1894. Collected and presented by J. Whitehead.

 There was only a single specimen, vide Ogilvie-Grant, Ibis, (6) 6 (1894): 509.
- p. 85. cacabata—Muscicapa sibirica cacabata Penard, 1919. . . 37

 Proc. New England zool. Cl., 7:22. New name for Hemichelidon fuliginosa Hodgson, 1845 (q.v.) on the grounds of its
 preoccupation by Muscicapa fuliginosa Sparrman, 1787 and
 Gmelin, 1789.
- p. 96. castaneiventris—Monarcha castaneiventris Verreaux, 1858. 38
 Hartert, Novit. Zool. 15 (1908): 363, states that the type belongs to the form found on Guadalcanar, Santa Isabel and Choiseul, and not on Navigators Islands. Mayr, Amer. Mus. Novit., 504 (1931): 23, refers only to Guadalcanar as Hartert's revised locality.

. 37

	Renamed Alseonax cinereus nigrorum Collin and Hartert, 1927, on the grounds of its preoccupation by Muscicapa cinerascens Spix, 1825.
p. 169.	everetti—Dicaeum everetti Tweeddale, 1877
p. 175.	fuliginosa—Hemichelidon fuliginosa Hodgson, 1845 37 Renamed Muscicapa sibirica cacabata Penard, 1919 on the grounds of its preoccupation by Muscicapa fuliginosa Sparrman, 1787 and Gmelin, 1789.
p. 210.	gouldii—Meliphaga gouldii Schlegel, 1872
p. 253.	incerta—Zosterops incerta Salvadori, 1892
p. 261.	insulana—Cossypho insulana Grote, 1935 31 Orn. Monatsb., 43: 95. New name for Callene poensis Boyd Alexander, 1903, (q.v.) on the grounds of its preoccupation by Cossypha poensis Strickland, 1844.
p. 271.	japonica—Loxia curvirostra japonica Ridgway, 1885 59 Proc. Biol. Soc. Washington, 2:103. New name for Loxia albiventris Swinhoe, 1870, (q.v.), on the grounds of its preoccupation by Loxia albiventris Hermann, 1804 (unidentified).
p. 338.	melanocephalus—Lanius collurioides melanocephalus Delacour, 1926
p. 349.	meridionalis—Parus meridionalis Sclater, 1857 42 Renamed Parus sclateri Kleinschmidt, 1897, on the grounds of its preoccupation by Parus meridionalis Lilljeborg, 1852.
p. 355.	minor—Anthreptes seimundi minor Bates, 1926 46 Renamed Nectarinia seimundi traylori Wolters, 1965, on the grounds of its preoccupation by Cinnyris angolensis minor Oustalet, 1893.
p. 362.	montana—Saxicola montana Gould, 1865 31 Renamed Saxicola oreophila Oberholser, 1900, on the grounds of its preoccupation by S. montana Koch, 1816.
p. 369.	mystacalis—Meliphaga mystacalis Gould, 1841 48 Renamed Meliphaga gouldii Schlegel, 1872, on the grounds of its preoccupation by M. mystacalis Temmine kand Laugier, 1825.

p. 116. cinerascens—Muscicapa cinerascens Sharpe, 1879 .

- p. 380. **nigricapillus**—Lanius collurioides nigricapillus Delacour, 1926 **23**Bull. Br. Orn. Cl., 47: 70. New name for, Lanius collurioidesm elanocephalus Delacour, 1926, (q.v.), on the grounds of its preoccupation by Lanius melanocephalus Gmelin, 1788.
- p. 389. niveiceps—Planesticus niveiceps Hellmayr, 1919 . . . 31

 Verh. Orn. Ges. Bayern, 14: 133. New name for Turdus albiceps Swinhoe, 1964 (q.v.), on the grounds of its preoccupation by Turdus albiceps Pucheran, 1854.
- p. 394. **nyikensis**—Nectarinia johnstoni nyikensis Delacour, 1944. **46**Zoologica, New York, 29: 32. New name for Nectarinia salvadorii Shelley, 1903 (q.v.) on the grounds of its preoccupation by Cinnyris salvadorii Shelley, 1877.
- p. 406. oreophila—Saxicola oreophila Oberholser, 1900 . . . 31

 Proc. U.S. Nat. Mus., 22: 221. New name for Saxicola montana Gould, 1865, (q.v.), on the grounds of its preoccupation by S. montana Koch, 1816.
- p. 407. **orientalis**—Merula torquata orientalis Seebohm, 1888 . **31**Renamed Turdus torquatus amicorum Hartert, 1923 on the grounds of its preoccupation by Turdus orientalis Gmelin, 1789. Also renamed T. torquatus caucasicus Collin and Hartert, 1927, for the same reason.
- p. 420. parvula—Colluricincla parvula Gould, 1845 39

 Proc. zool. Soc. Lond., (Sept.) 1845: 62, and Birds of Australia, 2 (Dec. 1845): pl. 78. ? Syntype, Adult (male, on label). Reg. no. 1881.5.1.446. Port Essington, (25 Jan. 1841 on label). Collected by J. Gilbert (Gilbert was in the locality until March 1841, vide Chisholm, Emu, 44 (1944): 131). Purchased of J. Gould.

The specimen lacks a tail. Gould did not define a holotype and the plate showed both sexes. Gould quotes from Gilbert. de Schauensee, *Proc. Acad. Nat. Sci. Philadelphia*, 109 (1957): 218, claims a holotype, together with a female, but is mistaken in stating that there is no specimen from Gould in the British Museum. The present specimen, marked by P. L. Sclater in the B.M. register as "type" appears to be a syntype.

p. 441. poensis—Callene poensis Boyd Alexander, 1903 . . . 31

Renamed Cossypha insulana Grote, 1935, on the grounds of its preoccupation by Cossypha poensis Strickland, 1844.

- p. 444. poliosoma—Pachycephalopsis poliosoma Sharpe, 1882 . 37 J. Linn. Soc. Lond., Zool., 16: 318. HOLOTYPE, Adult male. Reg. no. 1882.3.8.49. Astrolabe Mountains, New Guinea. Collected by A. Goldie. Purchased of Gerrard.
- p. 445. pontifex—Dicaeum hypoleucum pontifex Mayr, 1946. 45

 Zoologica, New York, 31:8. New name for Dicaeum everetti Tweeddale, (Dec.) 1877, (q.v.), on the grounds of its preoccupation by Prionochilus everetti Sharpe, (Jan.) 1877.
- p. 472. rubrocanus—Turdus rubrocanus Gray, 1846 31

 According to the amended article 11d incorporated into the second edition of the International Code of Zoological Nomenclature, (1963), this name is available because it was used by E. C. Stuart Baker, Fauna of British India, Birds, 8 (1930): 624, as a valid name. However, this availability is derived from its publication as a synonym of Merula castanea Gould, Proc. Zool. Soc., Lond., 1835: 185, so that the type-specimen of M. castanea, and not the specimen listed by Seebohm, Cat. Bds. Brit. Mus., 5 (1881): 260, is the type of rubrocanus. This specimen no longer exists.
- p. 475. ruficeps—Megalurus ruficeps Tweeddale, 1877. . . . 35
 Renamed Megalurus tweeddalei McGregor, 1908 on the grounds of its preoccupation by M. ? ruficeps Sykes, 1832.
- p. 486. salvadorii—Nectarinia salvadorii Shelley, 1903. . . . 46
 Renamed Nectarinia johnstoni nyikensis Delacour, 1944, on the grounds of its preoccupation by Cinnyris salvadorii Shelley, 1877.
- p. 497. sclateri—Parus sclateri Kleinschmidt, 1897 42

 J. Orn., 45: 133. New name for Parus meridionalis Sclater, 1857, (q.v.), on the grounds of its preoccupation by Parus meridionalis Lilljeborg, 1852.
- p. 561. tibetana—Prunella fulvescens tibetana Sushkin, 1925. 30
 Renamed Prunella fulvescens sushkini Collin and Hartert,
 1927 on the grounds of its preoccupation by Accentor collaris tibetanus Bianchi, 1904.

- p. 566. **traylori**—Nectarinia seimundi traylori Wolters, 1965 . **46**J. Orn., 106: 357. New name for Anthreptes seimundi minor
 Bates, 1926 (q.v.), on the grounds of its preoccupation by
 Cinnyris angolensis minor Oustalet, 1893.
- p. 570. tweeddalei—Megalurus tweeddalei McGregor, 1908. . . 35

 Philippine J. Sci., (A) 3:283. New name for Megalurus ruficeps Tweeddale, 1877 (q.v.), on the grounds of its preoccupation by M. ? ruficeps Sykes, 1832.

